

PTO/PCT HSE 1 FEB 2001

Nucleic acids encoding human TBC-1 protein and polymorphic markers thereof.

## FIELD OF THE INVENTION

The invention concerns genomic and cDNA sequences of the human *TBC-1* gene. The invention also concerns polypeptides encoded by the *TBC-1* gene. The invention also deals with antibodies directed specifically against such polypeptides that are useful as diagnostic reagents. The invention further encompasses biallelic markers of the *TBC-1* gene useful in genetic analysis.

## BACKGROUND OF THE INVENTION

The incidence of prostate cancer has dramatically increased over the last decades. It averages 30-50/100,000 males in Western European countries as well as within the US White male population. In these countries, it has recently become the most commonly diagnosed malignancy, being one of every four cancers diagnosed in American males. Prostate cancer's incidence is very much population specific, since it varies from 2/100,000 in China, to over 80/100,000 among African-American males.

In France, the incidence of prostate cancer is 35/100,000 males and it is increasing by 10/100,000 per decade. Mortality due to prostate cancer is also growing accordingly. It is the second cause of cancer death among French males, and the first one among French males aged over 70. This makes prostate cancer a serious burden in terms of public health.

Prostate cancer is a latent disease. Many men carry prostate cancer cells without overt signs of disease. Autopsies of individuals dying of other causes show prostate cancer cells in 30 % of men at age 50 and in 60 % of men at age 80. Furthermore, prostate cancer can take up to 10 years to kill a patient after the initial diagnosis.

The progression of the disease usually goes from a well-defined mass within the prostate to a breakdown and invasion of the lateral margins of the prostate, followed by metastasis to regional lymph nodes, and metastasis to the bone marrow. Cancer metastasis to bone is common and often associated with uncontrollable pain.

Unfortunately, in 80 % of cases, diagnosis of prostate cancer is established when the disease has already metastasized to the bones. Of special interest is the observation that prostate cancers frequently grow more rapidly in sites of metastasis than within the prostate itself.

Early-stage diagnosis of prostate cancer mainly relies today on Prostate Specific Antigen (PSA) dosage, and allows the detection of prostate cancer seven years before clinical symptoms become apparent. The effectiveness of PSA dosage diagnosis is however limited, due to its inability to discriminate between malignant and non-malignant affections of the organ and because not all prostate cancers give rise to an elevated serum PSA concentration. Furthermore, PSA dosage and

other currently available approaches such as physical examination, tissue biopsy and bone scans are of limited value in predicting disease progression.

Therefore, there is a strong need for a reliable diagnostic procedure which would enable a more systematic early-stage prostate cancer prognosis.

5 Although an early-stage prostate cancer prognosis is important, the possibility of measuring the period of time during which treatment can be deferred is also interesting as currently available medicaments are expensive and generate important adverse effects. However, the aggressiveness of prostate tumors varies widely. Some tumors are relatively aggressive, doubling every six months whereas others are slow-growing, doubling once every five years. In fact, the majority of prostate  
10 cancers grows relatively slowly and never becomes clinically manifest. Very often, affected patients are among the elderly and die from another disease before prostate cancer actually develops. Thus, a significant question in treating prostate carcinoma is how to discriminate between tumors that will progress and those that will not progress during the expected lifetime of the patient.

Hence, there is also a strong need for detection means which may be used to evaluate the  
15 aggressiveness or the development potential of prostate cancer tumors once diagnosed.

Furthermore, at the present time, there is no means to predict prostate cancer susceptibility. It would also be very beneficial to detect individual susceptibility to prostate cancer. This could allow preventive treatment and a careful follow up of the development of the tumor.

A further consequence of the slow growth rate of prostate cancer is that few cancer cells are  
20 actively dividing at any one time, rendering prostate cancer generally resistant to radiation and chemotherapy. Surgery is the mainstay of treatment but it is largely ineffective and removes the ejaculatory ducts, resulting in impotence. Oral oestrogens and luteinizing releasing hormone analogs are also used for treatment of prostate cancer. These hormonal treatments provide marked improvement for many patients, but they only provide temporary relief. Indeed, most of these  
25 cancers soon relapse with the development of hormone-resistant tumor cells and the oestrogen treatment can lead to serious cardiovascular complications. Consequently, there is a strong need for preventive and curative treatment of prostate cancer.

Efficacy/tolerance prognosis could be precious in prostate cancer therapy. Indeed, hormonal therapy, the main treatment currently available, presents important side effects. The use of  
30 chemotherapy is limited because of the small number of patients with chemosensitive tumors. Furthermore the age profile of the prostate cancer patient and intolerance to chemotherapy make the systematic use of this treatment very difficult.

Therefore, a valuable assessment of the eventual efficacy of a medicament to be administered to a prostate cancer patient as well as the patient's eventual tolerance to it may permit to  
35 enhance the benefit/risk ratio of prostate cancer treatment.

It is known today that there is a familial risk of prostate cancer. Clinical studies in the 1950s had already demonstrated a familial aggregation in prostate cancer. Control-case clinical studies

have been conducted more recently to attempt to evaluate the incidence of the genetic risk factors in the disease. Thus Steinberg et al., 1990, and McWhorter et al., 1992 confirm that the risk of prostate cancer is increased in subjects having one or more relatives already affected by the disease and when forms of early diagnosis in the relatives exist.

5 It is now well established that cancer is a disease caused by the deregulation of the expression of certain genes. In fact, the development of a tumor necessitates an important succession of steps. Each of these steps comprises the deregulation of an important gene intervening in the normal metabolism of the cell and the emergence of an abnormal cellular sub-clone which overwhelms the other cell types because of a proliferative advantage. The genetic origin of this  
10 concept has found confirmation in the isolation and the characterization of genes which could be responsible. These genes, commonly called "cancer genes", have an important role in the normal metabolism of the cell and are capable of intervening in carcinogenesis following a change.

Recent studies have identified three groups of genes which are frequently mutated in cancer. The first group of genes, called oncogenes, are genes whose products activate cell  
15 proliferation. The normal non-mutant versions are called protooncogenes. The mutated forms are excessively or inappropriately active in promoting cell proliferation, and act in the cell in a dominant way in that a single mutant allele is enough to affect the cell phenotype. Activated oncogenes are rarely transmitted as germline mutations since they may probably be lethal when expressed in all the cells. Therefore oncogenes can only be investigated in tumor tissues.

20 The second group of genes which are frequently mutated in cancer, called tumor suppressor genes, are genes whose products inhibit cell growth. Mutant versions in cancer cells have lost their normal function, and act in the cell in a recessive way in that both copies of the gene must be inactivated in order to change the cell phenotype. Most importantly, the tumor phenotype can be rescued by the wild type allele, as shown by cell fusion experiments first described by Harris and  
25 colleagues (1969). Germline mutations of tumor suppressor genes may be transmitted and thus studied in both constitutional and tumor DNA from familial or sporadic cases. The current family of tumor suppressors includes DNA-binding transcription factors (i.e., p53, WT1), transcription regulators (i.e., RB, APC, probably BRCA1), protein kinase inhibitors (i.e., p16), among others (for review, see Haber D & Harlow E, 1997).

30 The third group of genes which are frequently mutated in cancer, called mutator genes, are responsible for maintaining genome integrity and/or low mutation rates. Loss of function of both alleles increases cell mutation rates, and as a consequence, proto-oncogenes and tumor suppressor genes may be mutated. Mutator genes can also be classified as tumor suppressor genes, except for the fact that tumorigenesis caused by this class of genes cannot be suppressed simply by restoration  
35 of a wild-type allele, as described above. Genes whose inactivation may lead to a mutator phenotype include mismatch repair genes (i.e., MLH1, MSH2), DNA helicases (i.e., BLM, WRN)

or other genes involved in DNA repair and genomic stability (i.e., p53, possibly BRCA1 and BRCA2) (For review see Haber D & Harlow E, 1997; Fishel R & Wilson T. 1997; Ellis NA, 1997).

There is growing evidence that a critical event in the progression of a tumor cell from a non-metastatic to metastatic phenotype is the loss of function of metastasis-suppressor genes. These genes specifically suppress the ability of a cell to metastasize. Work from several groups has demonstrated that human chromosomes 8, 10, 11 and 17 encode prostate cancer metastasis suppressor activities. However, other human chromosomes such as chromosomes 1, 7, 13, 16, and 18 may also be associated to prostate cancer.

It thus remains to localize and to identify the genes specifically involved in the development and the progression of prostate cancers starting from the genetic analysis of the hereditary and the non-hereditary forms and to define their clinical implications in terms of prognosis and therapeutic innovations.

## SUMMARY OF THE INVENTION

The present invention pertains to nucleic acid molecules comprising the genomic sequence of a novel human gene which encodes a TBC-1 protein. The *TBC-1* genomic sequences comprise regulatory sequence located upstream (5'-end) and downstream (3'-end) of the transcribed portion of said gene, these regulatory sequences being also part of the invention. The human *TBC-1* genomic sequence is included in a previously unknown candidate region of prostate cancer located on chromosome 4.

The invention also deals with the two complete cDNA sequences encoding the TBC-1 protein, as well as with the corresponding translation product.

Oligonucleotide probes or primers hybridizing specifically with a *TBC-1* genomic or cDNA sequence are also part of the present invention, as well as DNA amplification and detection methods using said primers and probes.

A further object of the invention consists of recombinant vectors comprising any of the nucleic acid sequences described above, and in particular of recombinant vectors comprising a *TBC-1* regulatory sequence or a sequence encoding a TBC-1 protein, as well as of cell hosts and transgenic non human animals comprising said nucleic acid sequences or recombinant vectors.

The invention also concerns a *TBC-1*-related biallelic marker and the use thereof.

Finally, the invention is directed to methods for the screening of substances or molecules that inhibit the expression of *TBC-1*, as well as with methods for the screening of substances or molecules that interact with a TBC-1 polypeptide.



## BRIEF DESCRIPTION OF THE DRAWINGS

**Figure 1 :** An amino acid alignment of a portion of the amino acid sequence of the TBC-1 protein of SEQ ID No 5 with other proteins sharing amino acid homology with TBC-1. The amino acid numbering refers to the murine TBC-1.

### 5 Brief Description of the sequences provided in the Sequence Listing

SEQ ID No 1 contains a first part of the *TBC-1* genomic sequence comprising the 5' regulatory sequence and the exons 1, 1bis, and 2.

SEQ ID No 2 contains a second part of the *TBC-1* genomic sequence comprising the 12 last exons of the *TBC-1* gene and the 3' regulatory sequence.

10 SEQ ID No 3 contains a first cDNA sequence of the *TBC-1* gene.

SEQ ID No 4 contains a second cDNA sequence of the *TBC-1* gene.

SEQ ID No 5 contains the amino acid sequence encoded by the cDNAs of SEQ ID Nos 3 and 4.

15 SEQ ID No 6 contains a primer containing the additional PU 5' sequence described further in Example 3.

SEQ ID No 7 contains a primer containing the additional RP 5' sequence described further in Example 3.

In accordance with the regulations relating to Sequence Listings, the following codes have been used in the Sequence Listing to indicate the locations of biallelic markers within the sequences and to identify each of the alleles present at the polymorphic base. The code "r" in the sequences indicates that one allele of the polymorphic base is a guanine, while the other allele is an adenine. The code "y" in the sequences indicates that one allele of the polymorphic base is a thymine, while the other allele is a cytosine. The code "m" in the sequences indicates that one allele of the polymorphic base is an adenine, while the other allele is an cytosine. The code "k" in the sequences indicates that one allele of the polymorphic base is a guanine, while the other allele is a thymine. The code "s" in the sequences indicates that one allele of the polymorphic base is a guanine, while the other allele is a cytosine. The code "w" in the sequences indicates that one allele of the polymorphic base is an adenine, while the other allele is an thymine. The nucleotide code of the original allele for each biallelic marker is the following:

30	Biallelic marker	Original allele
	99-430-352	G
	99-20508-456	C
	99-20469-213	C
	5-254-227	A
35	5-257-353	C
	99-20511-32	T

	99-20511-221	A
	99-20504-90	G
	99-20493-238	A
	99-20499-221	G
5	99-20499-364	A
	99-20499-399	A
	5-249-304	G
	99-20485-269	A
	99-20481-131	G
10	99-20481-419	T
	99-20480-233	A

## DETAILED DESCRIPTION OF THE INVENTION

The present invention concerns polynucleotides and polypeptides related to the human *TBC-1* gene (also termed "*TBC-1* gene" throughout the present specification), which is potentially involved in the regulation of the differentiation of various cell types in mammals. A deregulation or an alteration of *TBC-1* expression, or alternatively an alteration in the amino acid sequence of the *TBC-1* protein may be involved in the generation of a pathological state related to cell differentiation in a patient, more particularly to abnormal cell proliferation leading to cancer states, such as prostate cancer.

### Definitions

Before describing the invention in greater detail, the following definitions are set forth to illustrate and define the meaning and scope of the terms used to describe the invention herein.

The term "*TBC-1* gene", when used herein, encompasses mRNA and cDNA sequences encoding the *TBC-1* protein. In the case of a genomic sequence, the *TBC-1* gene also includes native regulatory regions which control the expression of the coding sequence of the *TBC-1* gene.

The term "functionally active fragment" of the *TBC-1* protein is intended to designate a polypeptide carrying at least one of the structural features of the *TBC-1* protein involved in at least one of the biological functions and/or activity of the *TBC-1* protein.

A "heterologous" or "exogenous" polynucleotide designates a purified or isolated nucleic acid that has been placed, by genetic engineering techniques, in the environment of unrelated nucleotide sequences, such as the final polynucleotide construct does not occur naturally. An illustrative, but not limitative, embodiment of such a polynucleotide construct may be represented by a polynucleotide comprising (1) a regulatory polynucleotide derived from the *TBC-1* gene sequence and (2) a polynucleotide encoding a cytokine, for example GM-CSF. The polypeptide

encoded by the heterologous polynucleotide will be termed an heterologous polypeptide for the purpose of the present invention.

By a "biologically active fragment or variant" of a regulatory polynucleotide according to the present invention is intended a polynucleotide comprising or alternatively consisting in a  
5 fragment of said polynucleotide which is functional as a regulatory region for expressing a recombinant polypeptide or a recombinant polynucleotide in a recombinant cell host.

For the purpose of the invention, a nucleic acid or polynucleotide is "functional" as a regulatory region for expressing a recombinant polypeptide or a recombinant polynucleotide if said regulatory polynucleotide contains nucleotide sequences which contain transcriptional and  
10 translational regulatory information, and such sequences are "operatively linked" to nucleotide sequences which encode the desired polypeptide or the desired polynucleotide. An operable linkage is a linkage in which the regulatory nucleic acid and the DNA sequence sought to be expressed are linked in such a way as to permit gene expression.

A "promoter" refers to a DNA sequence recognized by the synthetic machinery of the cell  
15 required to initiate the specific transcription of a gene.

A sequence which is "operably linked" to a regulatory sequence such as a promoter means that said regulatory element is in the correct location and orientation in relation to the nucleic acid to control RNA polymerase initiation and expression of the nucleic acid of interest.

As used herein, the term "operably linked" refers to a linkage of polynucleotide elements in  
20 a functional relationship. For instance, a promoter or enhancer is operably linked to a coding sequence if it affects the transcription of the coding sequence. More precisely, two DNA molecules (such as a polynucleotide containing a promoter region and a polynucleotide encoding a desired polypeptide or polynucleotide) are said to be "operably linked" if the nature of the linkage between the two polynucleotides does not (1) result in the introduction of a frame-shift mutation or (2)  
25 interfere with the ability of the polynucleotide containing the promoter to direct the transcription of the coding polynucleotide. The promoter polynucleotide would be operably linked to a polynucleotide encoding a desired polypeptide or a desired polynucleotide if the promoter is capable of effecting transcription of the polynucleotide of interest.

The term "primer" denotes a specific oligonucleotide sequence which is complementary to  
30 a target nucleotide sequence and used to hybridize to the target nucleotide sequence. A primer serves as an initiation point for nucleotide polymerization catalyzed by either DNA polymerase, RNA polymerase or reverse transcriptase.

The term "probe" denotes a defined nucleic acid segment (or nucleotide analog segment, e.g., polynucleotide as defined hereinbelow) which can be used to identify a specific polynucleotide  
35 sequence present in samples, said nucleic acid segment comprising a nucleotide sequence complementary of the specific polynucleotide sequence to be identified.

007621.062501  
T05290.1T227600

The terms "sample" or "material sample" are used herein to designate a solid or a liquid material suspected to contain a polynucleotide or a polypeptide of the invention. A solid material may be, for example, a tissue slice or biopsy within which is searched the presence of a polynucleotide encoding a TBC-1 protein, either a DNA or RNA molecule or within which is searched the presence of a native or a mutated TBC-1 protein, or alternatively the presence of a desired protein of interest the expression of which has been placed under the control of a *TBC-1* regulatory polynucleotide. A liquid material may be, for example, any body fluid like serum, urine etc., or a liquid solution resulting from the extraction of nucleic acid or protein material of interest from a cell suspension or from cells in a tissue slice or biopsy. The term "biological sample" is also used and is more precisely defined within the Section dealing with DNA extraction.

As used herein, the term "purified" does not require absolute purity; rather, it is intended as a relative definition. Purification of starting material or natural material to at least one order of magnitude, preferably two or three orders, and more preferably four or five orders of magnitude is expressly contemplated. As an example, purification from 0.1% concentration to 10% concentration is two orders of magnitude.

The term "isolated" requires that the material be removed from its original environment (e.g. the natural environment if it is naturally occurring). For example, a naturally-occurring polynucleotide or polypeptide present in a living animal is not isolated, but the same polynucleotide or DNA or polypeptide, separated from some or all of the coexisting materials in the natural system, is isolated. Such polynucleotide could be part of a vector and/or such polynucleotide or polypeptide could be part of a composition and still be isolated in that the vector or composition is not part of its natural environment.

The term "polypeptide" refers to a polymer of amino acids without regard to the length of the polymer; thus, peptides, oligopeptides, and proteins are included within the definition of polypeptide. This term also does not specify or exclude post-expression modifications of polypeptides, for example, polypeptides which include the covalent attachment of glycosyl groups, acetyl groups, phosphate groups, lipid groups and the like are expressly encompassed by the term polypeptide. Also included within the definition are polypeptides which contain one or more analogs of an amino acid (including, for example, non-naturally occurring amino acids, amino acids which only occur naturally in an unrelated biological system, modified amino acids from mammalian systems etc.), polypeptides with substituted linkages, as well as other modifications known in the art, both naturally occurring and non-naturally occurring.

The term "recombinant polypeptide" is used herein to refer to polypeptides that have been artificially designed and which comprise at least two polypeptide sequences that are not found as contiguous polypeptide sequences in their initial natural environment, or to refer to polypeptides which have been expressed from a recombinant polynucleotide.

The term "purified" is used herein to describe a polypeptide of the invention which has been separated from other compounds including, but not limited to nucleic acids, lipids, carbohydrates and other proteins. A polypeptide is substantially pure when at least about 50%, preferably 60 to 75% of a sample exhibits a single polypeptide sequence. A substantially pure polypeptide typically comprises about 50%, preferably 60 to 90% weight/weight of a protein sample, more usually about 95%, and preferably is over about 99% pure. Polypeptide purity or homogeneity is indicated by a number of means well known in the art, such as polyacrylamide gel electrophoresis of a sample, followed by visualizing a single polypeptide band upon staining the gel. For certain purposes higher resolution can be provided by using HPLC or other means well known in the art.

As used herein, the term "non-human animal" refers to any non-human vertebrate, birds and more usually mammals, preferably primates, farm animals such as swine, goats, sheep, donkeys, and horses, rabbits or rodents, more preferably rats or mice. As used herein, the term "animal" is used to refer to any vertebrate, preferable a mammal. Both the terms "animal" and "mammal" expressly embrace human subjects unless preceded with the term "non-human".

As used herein, the term "antibody" refers to a polypeptide or group of polypeptides which are comprised of at least one binding domain, where an antibody binding domain is formed from the folding of variable domains of an antibody molecule to form three-dimensional binding spaces with an internal surface shape and charge distribution complementary to the features of an antigenic determinant of an antigen, which allows an immunological reaction with the antigen. Antibodies include recombinant proteins comprising the binding domains, as wells as fragments, including Fab, Fab', F(ab)<sub>2</sub>, and F(ab')<sub>2</sub> fragments.

As used herein, an "antigenic determinant" is the portion of an antigen molecule, in this case a TBC-1 polypeptide, that determines the specificity of the antigen-antibody reaction. An "epitope" refers to an antigenic determinant of a polypeptide. An epitope can comprise as few as 3 amino acids in a spatial conformation which is unique to the epitope. Generally an epitope consists of at least 6 such amino acids, and more usually at least 8-10 such amino acids. Methods for determining the amino acids which make up an epitope include x-ray crystallography, 2-dimensional nuclear magnetic resonance, and epitope mapping e.g. the Pepscan method described by Geysen et al. 1984; PCT Publication No. WO 84/03564; and PCT Publication No. WO 84/03506.

Throughout the present specification, the expression "nucleotide sequence" may be employed to designate indifferently a polynucleotide or an oligonucleotide or a nucleic acid. More precisely, the expression "nucleotide sequence" encompasses the nucleic material itself and is thus not restricted to the sequence information (i.e. the succession of letters chosen among the four base letters) that biochemically characterizes a specific DNA or RNA molecule.

As used interchangeably herein, the term "oligonucleotides", and "polynucleotides" include RNA, DNA, or RNA/DNA hybrid sequences of more than one nucleotide in either single chain or

5

10

15

20

25

30

Typically, between different genomes or between different individuals, the polymorphic site may be occupied by two different nucleotides.

The term "biallelic polymorphism" and "biallelic marker" are used interchangeably herein to refer to a single nucleotide polymorphism having two alleles at a fairly high frequency in the population. A "biallelic marker allele" refers to the nucleotide variants present at a biallelic marker site. Typically, the frequency of the less common allele of the biallelic markers of the present invention has been validated to be greater than 1%, preferably the frequency is greater than 10%, more preferably the frequency is at least 20% (i.e. heterozygosity rate of at least 0.32), even more preferably the frequency is at least 30% (i.e. heterozygosity rate of at least 0.42). A biallelic marker wherein the frequency of the less common allele is 30% or more is termed a "high quality biallelic marker".

The location of nucleotides in a polynucleotide with respect to the center of the polynucleotide are described herein in the following manner. When a polynucleotide has an odd number of nucleotides, the nucleotide at an equal distance from the 3' and 5' ends of the polynucleotide is considered to be "at the center" of the polynucleotide, and any nucleotide immediately adjacent to the nucleotide at the center, or the nucleotide at the center itself is considered to be "within 1 nucleotide of the center." With an odd number of nucleotides in a polynucleotide any of the five nucleotides positions in the middle of the polynucleotide would be considered to be within 2 nucleotides of the center, and so on. When a polynucleotide has an even number of nucleotides, there would be a bond and not a nucleotide at the center of the polynucleotide. Thus, either of the two central nucleotides would be considered to be "within 1 nucleotide of the center" and any of the four nucleotides in the middle of the polynucleotide would be considered to be "within 2 nucleotides of the center", and so on. For polymorphisms which involve the substitution, insertion or deletion of 1 or more nucleotides, the polymorphism, allele or biallelic marker is "at the center" of a polynucleotide if the difference between the distance from the substituted, inserted, or deleted polynucleotides of the polymorphism and the 3' end of the polynucleotide, and the distance from the substituted, inserted, or deleted polynucleotides of the polymorphism and the 5' end of the polynucleotide is zero or one nucleotide. If this difference is 0 to 3, then the polymorphism is considered to be "within 1 nucleotide of the center." If the difference is 0 to 5, the polymorphism is considered to be "within 2 nucleotides of the center." If the difference is 0 to 7, the polymorphism is considered to be "within 3 nucleotides of the center," and so on.

As used herein the terminology "defining a biallelic marker" means that a sequence includes a polymorphic base from a biallelic marker. The sequences defining a biallelic marker may be of any length consistent with their intended use, provided that they contain a polymorphic base from a biallelic marker. The sequence has between 1 and 500 nucleotides in length, preferably between 5, 10, 15, 20, 25, or 40 and 200 nucleotides and more preferably between 30 and 50 nucleotides in length. Each biallelic marker therefore corresponds to two forms of a polynucleotide sequence included in a gene, which, when compared with one another, present a nucleotide

09762344.062501  
T05290"TE29260

modification at one position. Preferably, the sequences defining a biallelic marker include a polymorphic base selected from the group consisting of the biallelic markers A1 to A19 and the complements thereof. In some embodiments the sequences defining a biallelic marker comprise one of the sequences selected from the group consisting of P1 to P7, P9 to P13, P15 to P19 and the complementary sequences thereto. Likewise, the term "marker" or "biallelic marker" requires that the sequence is of sufficient length to practically (although not necessarily unambiguously) identify the polymorphic allele, which usually implies a length of at least 4, 5, 6, 10, 15, 20, 25, or 40 nucleotides.

The term "upstream" is used herein to refer to a location which is toward the 5' end of the polynucleotide from a specific reference point.

The terms "base paired" and "Watson & Crick base paired" are used interchangeably herein to refer to nucleotides which can be hydrogen bonded to one another by virtue of their sequence identities in a manner like that found in double-helical DNA with thymine or uracil residues linked to adenine residues by two hydrogen bonds and cytosine and guanine residues linked by three hydrogen bonds (See Stryer, L., *Biochemistry*, 4<sup>th</sup> edition, 1995).

The terms "complementary" or "complement thereof" are used herein to refer to the sequences of polynucleotides which is capable of forming Watson & Crick base pairing with another specified polynucleotide throughout the entirety of the complementary region. For the purpose of the present invention, a first polynucleotide is deemed to be complementary to a second polynucleotide when each base in the first polynucleotide is paired with its complementary base. Complementary bases are, generally, A and T (or A and U), or C and G. "Complement" is used herein as a synonym from "complementary polynucleotide", "complementary nucleic acid" and "complementary nucleotide sequence". These terms are applied to pairs of polynucleotides based solely upon their sequences and not any particular set of conditions under which the two polynucleotides would actually bind.

## **Variants and fragments**

### **1. Polynucleotides**

The invention also relates to variants and fragments of the polynucleotides described herein, particularly of a *TBC-1* gene containing one or more biallelic markers according to the invention.

Variants of polynucleotides, as the term is used herein, are polynucleotides that differ from a reference polynucleotide. A variant of a polynucleotide may be a naturally occurring variant such as a naturally occurring allelic variant, or it may be a variant that is not known to occur naturally. Such non-naturally occurring variants of the polynucleotide may be made by mutagenesis techniques, including those applied to polynucleotides, cells or organisms. Generally, differences are limited so that the nucleotide sequences of the reference and the variant are closely similar overall and, in many regions, identical.



Variants of polynucleotides according to the invention include, without being limited to, nucleotide sequences that are at least 95% identical to any of SEQ ID Nos 1-4 or the sequences complementary thereto or to any polynucleotide fragment of at least 8 consecutive nucleotides of any of SEQ ID Nos 1-4 or the sequences complementary thereto, and preferably at least 98% identical, more particularly at least 99.5% identical, and most preferably at least 99.9% identical to any of SEQ ID Nos 1-4 or the sequences complementary thereto or to any polynucleotide fragment of at least 8 consecutive nucleotides of any of SEQ ID Nos 1-4 or the sequences complementary thereto.

Changes in the nucleotide of a variant may be silent, which means that they do not alter the amino acids encoded by the polynucleotide.

However, nucleotide changes may also result in amino acid substitutions, additions, deletions, fusions and truncations in the polypeptide encoded by the reference sequence. The substitutions, deletions or additions may involve one or more nucleotides. The variants may be altered in coding or non-coding regions or both. Alterations in the coding regions may produce conservative or non-conservative amino acid substitutions, deletions or additions.

In the context of the present invention, particularly preferred embodiments are those in which the polynucleotides encode polypeptides which retain substantially the same biological function or activity as the mature TBC-1 protein.

A polynucleotide fragment is a polynucleotide having a sequence that entirely is the same as part but not all of a given nucleotide sequence, preferably the nucleotide sequence of a *TBC-1* gene, and variants thereof. The fragment can be a portion of an exon or of an intron of a *TBC-1* gene. It can also be a portion of the regulatory sequences of the *TBC-1* gene. Preferably, such fragments comprise the polymorphic base of a biallelic marker selected from the group consisting of the biallelic markers A1 to A19 and the complements thereof.

Such fragments may be "free-standing", i.e. not part of or fused to other polynucleotides, or they may be comprised within a single larger polynucleotide of which they form a part or region. However, several fragments may be comprised within a single larger polynucleotide.

As representative examples of polynucleotide fragments of the invention, there may be mentioned those which have from about 4, 6, 8, 15, 20, 25, 40, 10 to 20, 10 to 30, 30 to 55, 50 to 100, 75 to 100 or 100 to 200 nucleotides in length. Preferred are those fragments having about 49 nucleotides in length, such as those of P1 to P7, P9 to P13, P15 to P19 or the sequences complementary thereto and containing at least one of the biallelic markers of a *TBC-1* gene which are described herein.

## 2. Polypeptides.

The invention also relates to variants, fragments, analogs and derivatives of the polypeptides described herein, including mutated TBC-1 proteins.

The variant may be 1) one in which one or more of the amino acid residues are substituted with a conserved or non-conserved amino acid residue (preferably a conserved amino acid residue) and such substituted amino acid residue may or may not be one encoded by the genetic code, or 2) one in which one or more of the amino acid residues includes a substituent group, or 3) one in which the mutated TBC-1 is fused with another compound, such as a compound to increase the half-life of the polypeptide (for example, polyethylene glycol), or 4) one in which the additional amino acids are fused to the mutated TBC-1, such as a leader or secretory sequence or a sequence which is employed for purification of the mutated TBC-1 or a preprotein sequence. Such variants are deemed to be within the scope of those skilled in the art.

More particularly, a variant TBC-1 polypeptide comprises amino acid changes ranging from 1, 2, 3, 4, 5, 10 to 20 substitutions, additions or deletions of one amino acid, preferably from 1 to 10, more preferably from 1 to 5 and most preferably from 1 to 3 substitutions, additions or deletions of one amino acid. The preferred amino acid changes are those which have little or no influence on the biological activity or the capacity of the variant TBC-1 polypeptide to be recognized by antibodies raised against a native TBC-1 protein.

By homologous peptide according to the present invention is meant a polypeptide containing one or several amino acid additions, deletions and/or substitutions in the amino acid sequence of a TBC-1 polypeptide. In the case of an amino acid substitution, one or several consecutive or non-consecutive amino acids are replaced by « equivalent » amino acids.

The expression "equivalent" amino acid is used herein to designate any amino acid that may be substituted for one of the amino acids having similar properties, such that one skilled in the art of peptide chemistry would expect the secondary structure and hydrophobic nature of the polypeptide to be substantially unchanged. Generally, the following groups of amino acids represent equivalent changes: (1) Ala, Pro, Gly, Glu, Asp, Gln, Asn, Ser, Thr; (2) Cys, Ser, Tyr, Thr; (3) Val, Ile, Leu, Met, Ala, Phe; (4) Lys, Arg, His; (5) Phe, Tyr, Trp, His.

By an equivalent amino acid according to the present invention is also meant the replacement of a residue in the L-form by a residue in the D form or the replacement of a Glutamic acid (E) residue by a Pyro-glutamic acid compound. The synthesis of peptides containing at least one residue in the D-form is, for example, described by Koch (1977).

A specific, but not restrictive, embodiment of a modified peptide molecule of interest according to the present invention, which consists in a peptide molecule which is resistant to proteolysis, is a peptide in which the -CONH- peptide bond is modified and replaced by a (CH<sub>2</sub>NH) reduced bond, a (NHCO) retro inverso bond, a (CH<sub>2</sub>-O) methylene-oxy bond, a (CH<sub>2</sub>-S) thiomethylene bond, a (CH<sub>2</sub>CH<sub>2</sub>) carba bond, a (CO-CH<sub>2</sub>) cetomethylene bond, a (CHOH-CH<sub>2</sub>) hydroxyethylene bond, a (N-N) bound, a E-alcene bond or also a -CH=CH- bond.

The polypeptide according to the invention could have post-translational modifications. For example, it can present the following modifications: acylation, disulfide bond formation, prenylation, carboxymethylation and phosphorylation.

A polypeptide fragment is a polypeptide having a sequence that entirely is the same as part  
5 but not all of a given polypeptide sequence, preferably a polypeptide encoded by a *TBC-1* gene and variants thereof. Preferred fragments include those regions possessing antigenic properties and which can be used to raise antibodies against the TBC-1 protein.

Such fragments may be "free-standing", i.e. not part of or fused to other polypeptides, or they may be comprised within a single larger polypeptide of which they form a part or region.

10 However, several fragments may be comprised within a single larger polypeptide.

As representative examples of polypeptide fragments of the invention, there may be mentioned those which comprise at least about 5, 6, 7, 8, 9 or 10 to 15, 10 to 20, 15 to 40, or 30 to 55 amino acids of the TBC-1. In some embodiments, the fragments contain at least one amino acid mutation in the TBC-1 protein.

## 15 Identity Between Nucleic Acids Or Polypeptides

The terms "percentage of sequence identity" and "percentage homology" are used interchangeably herein to refer to comparisons among polynucleotides and polypeptides, and are determined by comparing two optimally aligned sequences over a comparison window, wherein the portion of the polynucleotide or polypeptide sequence in the comparison window may comprise  
20 additions or deletions (i.e., gaps) as compared to the reference sequence (which does not comprise additions or deletions) for optimal alignment of the two sequences. The percentage is calculated by determining the number of positions at which the identical nucleic acid base or amino acid residue occurs in both sequences to yield the number of matched positions, dividing the number of matched positions by the total number of positions in the window of comparison and multiplying the result  
25 by 100 to yield the percentage of sequence identity. Homology is evaluated using any of the variety of sequence comparison algorithms and programs known in the art. Such algorithms and programs include, but are by no means limited to, TBLASTN, BLASTP, FASTA, TFASTA, and CLUSTALW (Pearson and Lipman, 1988; Altschul et al., 1990; Thompson et al., 1994; Higgins et al., 1996; Altschul et al., 1993). In a particularly preferred embodiment, protein and nucleic acid  
30 sequence homologies are evaluated using the Basic Local Alignment Search Tool ("BLAST") which is well known in the art (see, e.g., Karlin and Altschul, 1990; Altschul et al., 1990, 1993, 1997). In particular, five specific BLAST programs are used to perform the following task:

- (1) BLASTP and BLAST3 compare an amino acid query sequence against a protein sequence database;
- 35 (2) BLASTN compares a nucleotide query sequence against a nucleotide sequence database;

09763341.062501

(3) BLASTX compares the six-frame conceptual translation products of a query nucleotide sequence (both strands) against a protein sequence database;

(4) TBLASTN compares a query protein sequence against a nucleotide sequence database translated in all six reading frames (both strands); and

5 (5) TBLASTX compares the six-frame translations of a nucleotide query sequence against the six-frame translations of a nucleotide sequence database.

The BLAST programs identify homologous sequences by identifying similar segments, which are referred to herein as "high-scoring segment pairs," between a query amino or nucleic acid sequence and a test sequence which is preferably obtained from a protein or nucleic acid sequence database.

10 High-scoring segment pairs are preferably identified (i.e., aligned) by means of a scoring matrix, many of which are known in the art. Preferably, the scoring matrix used is the BLOSUM62 matrix (Gonnet et al., 1992; Henikoff and Henikoff, 1993). Less preferably, the PAM or PAM250 matrices may also be used (see, e.g., Schwartz and Dayhoff, eds., 1978). The BLAST programs evaluate the statistical significance of all high-scoring segment pairs identified, and preferably  
15 selects those segments which satisfy a user-specified threshold of significance, such as a user-specified percent homology. Preferably, the statistical significance of a high-scoring segment pair is evaluated using the statistical significance formula of Karlin (see, e.g., Karlin and Altschul, 1990).

#### Stringent Hybridization Conditions

20 By way of example and not limitation, procedures using conditions of high stringency are as follows: Prehybridization of filters containing DNA is carried out for 8 h to overnight at 65°C in buffer composed of 6X SSC, 50 mM Tris-HCl (pH 7.5), 1 mM EDTA, 0.02% PVP, 0.02% Ficoll, 0.02% BSA, and 500 µg/ml denatured salmon sperm DNA. Filters are hybridized for 48 h at 65°C, the preferred hybridization temperature, in prehybridization mixture containing 100 µg/ml  
25 denatured salmon sperm DNA and 5-20 X 10<sup>6</sup> cpm of <sup>32</sup>P-labeled probe. Alternatively, the hybridization step can be performed at 65°C in the presence of SSC buffer, 1 x SSC corresponding to 0.15M NaCl and 0.05 M Na citrate. Subsequently, filter washes can be done at 37°C for 1 h in a solution containing 2 x SSC, 0.01% PVP, 0.01% Ficoll, and 0.01% BSA, followed by a wash in 0.1 X SSC at 50°C for 45 min. Alternatively, filter washes can be performed in a solution containing 2  
30 x SSC and 0.1% SDS, or 0.5 x SSC and 0.1% SDS, or 0.1 x SSC and 0.1% SDS at 68°C for 15 minute intervals. Following the wash steps, the hybridized probes are detectable by autoradiography. Other conditions of high stringency which may be used are well known in the art and as cited in Sambrook et al., 1989; and Ausubel et al., 1989, are incorporated herein in their entirety. These hybridization conditions are suitable for a nucleic acid molecule of about 20  
35 nucleotides in length. There is no need to say that the hybridization conditions described above are to be adapted according to the length of the desired nucleic acid, following techniques well known to the one skilled in the art. The suitable hybridization conditions may for example be adapted

097634 06504  
T05390 TEE9460

according to the teachings disclosed in the book of Hames and Higgins (1985) or in Sambrook et al.(1989).

#### Candidate Region On The Chromosome 4 (Linkage Analysis).

In order to localize the prostate cancer gene(s) starting from families, a systematic familial  
5 study of genetic link research is carried out using markers of the microsatellite type described at the Genethon laboratory by the Jean Weissenbach team (Dib et al., 1996).

The studies of genetic link or of "linkage" are based on the principle according to which  
two neighboring sequences on a chromosome do not present (or very rarely present) recombinations  
by crossing-over during meiosis. To do this, microsatellite DNA sequences (chromosomal markers)  
10 constantly co-inherited with the disease studied are searched for in a family having a predisposition  
for this disease. These DNA sequences organized in the form of a repetition of di-, tri- or  
tetranucleotides are systematically present along the genome, and thus allow the identification of  
chromosomal fragments harboring them. More than 5000 microsatellite markers, have been  
localized with precision on the genome as a result of the first studies on the genetic map carried out  
15 at Genethon under the supervision of Jean Weissenbach, and on the physical map (using the "Yeast  
Artificial Chromosomes"), work conducted by Daniel Cohen at C.E.P.H. and at Genethon  
(Chumakov et al., 1995). Genetic link analysis calculates the probabilities of recombinations of the  
target gene with the chromosomal markers used, according to the genealogical tree, the transmission  
of the disease, and the transmission of the markers. Thus if a particular allele of a given marker is  
20 transmitted with the disease more often than chance would have it (recombination level of between  
0 and 0.5), it is possible to deduce that the target gene in question is found in the neighborhood of  
the marker. Using this technique, it has been possible to localize several genes of genetic  
predisposition to familial cancers. In order to be able to be included in a genetic link study, the  
families affected by a hereditary form of the disease must satisfy the "informativeness" criteria:  
25 several affected subjects (and whose constitutional DNA is available) per generation, and at best  
having a large number of siblings.

By linkage analysis, the inventors have identified a candidate region for prostate cancer on  
chromosome 4. Indeed, the LOD scores at 2 points between the disease and the markers on a total  
population of approximately fifty families present a value of 2.49 for marker D4S398 which  
30 indicates a probable genetic link with this marker. The curve of the variation of the LOD score on a  
map of 5 markers is centered on D4S398 and the value higher than 3.3 indicates that a gene  
involved in familial prostate cancer is probably found in the region located between markers  
D4S2978 and D4S3018, or a space of approximately 9.7 cM.

### Homologies Of The Novel Human Gene Translation Product With A Known Murine Protein.

A novel human gene was found in this candidate region. It presents a good probability to be involved in cancer. Database homology searches have allowed the inventors to determine that the translation product of this novel human gene has significant identity with a murine protein called *tbc1*. Therefore, the novel human gene of the invention has thus been called *TBC-1* throughout the present specification. *TBC-1* comprises an open Reading frame that encodes a novel protein, the TBC-1 protein. Based on sequence similarity, an alignment of a portion of the TBC-1 amino acid sequence with the known *tbc1* murine protein, it is expected that TBC1 protein may play a role in the cell cycle and in differentiation of various tissues. Indeed, the TBC1 protein contains a 200 amino acid domain called the TBC domain that is homologous to regions in the *trc2*-oncogene and in the yeast regulators of mitosis BUB2 and *cdc16*.

The cDNA of the murine *tbc1* gene has been described in US Patent No US 5,700,927 and it encodes a putative protein product of 1141 amino acids. The N-terminus of the murine *tbc1* protein contains stretches of cysteines and histidines which may form zinc finger structures in the mature polypeptides. The N-terminus also comprises short stretches of basic amino acids which may be involved in a nuclear localization signal. The TBC domain of the murine *tbc1* protein contains several tyrosine residues which are conserved in BUB2 and *cdc16*. The C-terminus of the murine *tbc1* protein contains a long stretch of evenly spaced leucine residues which are susceptible to form a leucine zipper motif.

The murine *tbc1* gene has been shown to be highly expressed in testis and kidney. However, lower levels of expression have also been identified in lung, spleen, brain, and heart. Moreover, murine *tbc1* is a nuclear protein which is expressed in a cell- and stage-specific manner.

Studies of murine bone marrow have demonstrated that erythroid cells and megakaryocytes expressed substantial levels of the murine *tbc1* protein, but none was detected in mature neutrophils.

Similarly, spermatogonia do not express murine *tbc1*, but primary and secondary spermatocytes express abundant *tbc1*. Later in the differentiation of the germ cells, the *tbc1* levels appear to decrease in spermatids and active sperm. The differentiation program of spermatogonia to spermatocytes therefore involves a significant upregulation of murine *tbc1* expression.

The general distribution of murine *tbc1* is not tissue-specific, but is cell-specific within individual tissues and intimately linked to tissue differentiation. The developmental expression of murine *tbc1*, particularly in hematopoietic and germ cells, suggests that this gene plays a role in the terminal differentiation program of several tissues.

Consequently, an alteration in the expression of the *TBC-1* gene or in the amino acid sequence of the TBC-1 protein leading to an altered biological activity of the latter is likely to cause, directly or indirectly, cell proliferation disorders and thus diseases related to an abnormal cell proliferation such as cancer, particularly prostate cancer.

### Genomic Sequence Of *TBC-1*

The present invention concerns the genomic sequence of *TBC-1*. The present invention encompasses the *TBC-1* gene, or *TBC-1* genomic sequences consisting of, consisting essentially of, or comprising a sequence selected from the group consisting of SEQ ID Nos 1 and 2, a sequence  
5 complementary thereto, as well as fragments and variants thereof. These polynucleotides may be purified, isolated, or recombinant.

The inventors have sequenced two portions of the *TBC-1* genomic sequence. The first portion of the *TBC-1* gene sequence contains the three first exons of the *TBC-1* gene, designated as Exon 1, Exon 1bis and Exon 2, and the 5' regulatory sequence located upstream of the transcribed  
10 sequences. The sequence of the first portion of the genomic sequence is disclosed in SEQ ID No 1. The second portion contains the twelve last exons of the *TBC-1* gene, designated as exons A, B, C, D, E, F, G, H, I, J, K, and L, and the 3' regulatory sequence which is located downstream of the transcribed sequences.

The exon positions in SEQ ID Nos 1 and 2 are detailed below in Table A.

Table A

Exon	Position in SEQ ID No 1		Intron	Position in SEQ ID No 1	
	Beginning	End		Beginning	End
1	2001	2077	1	2078	12739
1bis	12292	12373	1bis	12374	12739
2	12740	13249	2	13250	at least 17590
Exon	Position in SEQ ID No 2		Intron	Position in SEQ ID No 2	
	Beginning	End		Beginning	End
A	4661	4789	A	4790	6115
B	6116	6202	B	6203	9918
C	9919	10199	C	10200	14520
D	14521	14660	D	14661	50256
E	50257	50442	E	50443	56255
F	56256	56417	F	56418	63325
G	63326	63484	G	63485	76035
H	76036	76280	H	76281	78363
I	78364	78523	I	78524	85294
J	85295	85464	J	85465	93416
K	93417	93590	K	93591	97475
L	97476	97960			

Intron 1 refers to the nucleotide sequence located between Exon 1 and Exon 2; Intron 1bis refers to the nucleotide sequence located between Exon 1bis and Exon 2; Intron A refers to the nucleotide sequence located between Exon A and Exon B; and so on. The position of the introns is  
20 detailed in Table A.

The *TBC-1* introns defined hereinafter for the purpose of the present invention are not exactly what is generally understood as "introns" by the one skilled in the art and will consequently be further defined below.

Generally, an intron is defined as a nucleotide sequence that is present both in the genomic DNA and in the unspliced mRNA molecule, and which is absent from the mRNA molecule which has already gone through splicing events. In the case of the *TBC-1* gene, the inventors have found that at least two different spliced mRNA molecules are produced when this gene is transcribed, as it will be described in detail in a further section of the specification. The first spliced mRNA molecule comprises Exons 1 and 2. Thus, the genomic nucleotide sequence comprised between Exon 1 and Exon 2 is an intronic sequence as regards to this first mRNA molecule, despite the fact that this intronic sequence contains Exon 1*bis*. In contrast, Exon 1*bis* is of course an exonic nucleotide sequence as regards to the second *TBC-1* mRNA molecule.

For the purpose of the present invention and in order to make a clear and unambiguous designation of the different nucleic acids encompassed, it has been postulated that the polynucleotides contained both in any of the nucleotide sequences of SEQ ID Nos 1 or 2 and in any of the nucleotide sequences of SEQ ID Nos 3 or 4 are considered as exonic sequences. Conversely, the polynucleotides contained in any of the nucleotide sequences of SEQ ID Nos 1 or 2 but which are absent both from the nucleotide sequence of SEQ ID No 3 and from the nucleotide sequence of SEQ ID No 4 are considered as intronic sequences.

The nucleic acids defining the *TBC-1* introns described above, as well as their fragments and variants, may be used as oligonucleotide primers or probes in order to detect the presence of a copy of the *TBC-1* gene in a test sample, or alternatively in order to amplify a target nucleotide sequence within the *TBC-1* intronic sequences.

Thus, the invention embodies purified, isolated, or recombinant polynucleotides comprising a nucleotide sequence selected from the group consisting of the 15 exons of the *TBC-1* gene which are described in the present invention, or a sequence complementary thereto. The invention also deals with purified, isolated, or recombinant nucleic acids comprising a combination of at least two exons of the *TBC-1* gene, wherein the polynucleotides are arranged within the nucleic acid, from the 5'-end to the 3'-end of said nucleic acid, in the same order as in SEQ ID Nos 1 and 2.

Thus, the invention embodies purified, isolated, or recombinant polynucleotides comprising a nucleotide sequence selected from the group consisting of the introns of the *TBC-1* gene, or a sequence complementary thereto.

The invention also encompasses a purified, isolated, or recombinant polynucleotide comprising a nucleotide sequence having at least 70, 75, 80, 85, 90, or 95% nucleotide identity with a sequence selected from the group consisting of SEQ ID Nos 1 and 2 or a complementary sequence thereto or a fragment thereof. The nucleotide differences as regards to the nucleotide sequence of SEQ ID Nos 1 or 2 may be generally randomly distributed throughout the entire nucleic acid.



Nevertheless, preferred nucleic acids are those wherein the nucleotide differences as regards to the nucleotide sequence of SEQ ID Nos 1 or 2 are predominantly located outside the coding sequences contained in the exons. These nucleic acids, as well as their fragments and variants, may be used as oligonucleotide primers or probes in order to detect the presence of a copy of the *TBC-1* gene in a test sample, or alternatively in order to amplify a target nucleotide sequence within the *TBC-1* sequences.

Another object of the invention consists of a purified, isolated, or recombinant nucleic acid that hybridizes with a sequence selected from the group consisting of SEQ ID Nos 1 and 2 or a complementary sequence thereto or a variant thereof, under the stringent hybridization conditions as defined above.

Particularly preferred nucleic acids of the invention include isolated, purified, or recombinant polynucleotides comprising a contiguous span of at least 12, 15, 18, 20, 25, 30, 35, 40, 50, 60, 70, 80, 90, 100, 150, 200, 500, or 1000 nucleotides of a nucleotide sequence selected from the group consisting of SEQ ID Nos 1 and 2, or the complements thereof. Additionally preferred nucleic acids of the invention include isolated, purified, or recombinant polynucleotides comprising a contiguous span of at least 12, 15, 18, 20, 25, 30, 35, 40, 50, 60, 70, 80, 90, 100, 150, 200, 500, or 1000 nucleotides of SEQ ID No 1 or the complements thereof, wherein said contiguous span comprises at least 1, 2, 3, 5, or 10 of the following nucleotide positions of SEQ ID No 1: 1-1000, 1001-2000, 2001-3000, 3001-4000, 4001-5000, 5001-6000, 6001-7000, 7001-8000, 8001-9000, 9001-10000, 10001-11000, 11001-12000, 12001-13000, 13001-14000, 14001-15000, 15001-16000, 16001-17000, and 17001-17590. Other preferred nucleic acids of the invention include isolated, purified, or recombinant polynucleotides comprising a contiguous span of at least 12, 15, 18, 20, 25, 30, 35, 40, 50, 60, 70, 80, 90, 100, 150, 200, 500, or 1000 nucleotides of SEQ ID No 2 or the complements thereof, wherein said contiguous span comprises at least 1, 2, 3, 5, or 10 of the following nucleotide positions of SEQ ID No 2: 1-5000, 5001-10000, 10001-15000, 15001-20000, 20001-25000, 25001-30000, 30001-35000, 35001-40000, 40001-45000, 45001-50000, 50001-55000, 55001-60000, 60001-65000, 65001-70000, 70001-75000, 75001-80000, 80001-85000, 85001-90000, 90001-95000, and 95001-99960.

While this section is entitled "Genomic Sequences of *TBC-1*," it should be noted that nucleic acid fragments of any size and sequence may also be comprised by the polynucleotides described in this section, flanking the genomic sequences of *TBC-1* on either side or between two or more such genomic sequences.

#### *TBC-1* cDNA Sequences

The inventors have discovered that the expression of the *TBC-1* gene leads to the production of at least two mRNA molecules, respectively a first and a second *TBC-1* transcription

product, as the results of alternative splicing events. They result from two distinct first exons, namely Exon 1 and Exon 1bis.

The first transcription product comprises Exons 1, 2, A, B, C, D, E, F, G, H, I, J, K, and L. This cDNA of SEQ ID No 3 includes a 5'-UTR region, spanning the whole Exon 1 and part of Exon 2. This 5'-UTR region starts from the nucleotide at position 1 and ends at the nucleotide at position 170 of the nucleotide sequence of SEQ ID No 3. The cDNA of SEQ ID No 3 includes a 3'-UTR region starting from the nucleotide at position 3726 and ending at the nucleotide at position 3983 of the nucleotide sequence of SEQ ID No 3. This first transcription product harbors a polyadenylation signal located between the nucleotide at position 3942 and the nucleotide at position 3947 of the nucleotide sequence of SEQ ID No 3.

The second *TBC-1* transcription product comprises Exons 1bis, 2, A, B, C, D, E, F, G, H, I, J, K, and L. This cDNA of SEQ ID No 4 includes a 5'-UTR region starting from the nucleotide at position 1 and ending at the nucleotide at position 175 of the nucleotide sequence of SEQ ID No 4. This second cDNA also includes a 3'-UTR region starting from the nucleotide at position 3731 and ending at the nucleotide at position 3988 of the nucleotide sequence of SEQ ID No 4. This second transcription product harbors a polyadenylation signal located between the nucleotide at position 3947 and the nucleotide at position 3952 of the nucleotide sequence of SEQ ID No 4.

The 5'-end sequence of this second *TBC-1* mRNA, more particularly the nucleotide sequence comprised between the nucleotide in position 1 and the nucleotide in position 458 of the nucleic acid of SEQ ID No 4 molecule corresponds to the nucleotide sequence of a 5'-EST that has been obtained from a human pancreas cDNA library and characterized following the teachings of the PCT Application No WO 96/34981. This 5'-EST is also part of the invention.

Another object of the invention consists of a purified or isolated nucleic acid comprising a polynucleotide selected from the group consisting of the nucleotide sequences of SEQ ID Nos 3 and 4 and to nucleic acid fragments thereof.

Preferred nucleic acid fragments of the nucleotide sequences of SEQ ID Nos 3 and 4 consist in polynucleotides comprising their respective Open Reading Frames encoding the TBC-1 protein.

Other preferred nucleic acid fragments of the nucleotide sequences of SEQ ID Nos 3 and 4 consist in polynucleotides comprising at least a part of their respective 5'-UTR or 3'-UTR regions.

The invention also pertains to a purified or isolated nucleic acid having at least a 95% of nucleotide identity with any one of the nucleotide sequences of SEQ ID Nos 3 and 4, or a fragment thereof.

Another object of the invention consists of purified, isolated or recombinant nucleic acids comprising a polynucleotide that hybridizes, under the stringent hybridization conditions defined herein, with any one of the nucleotide sequences of SEQ ID Nos 3 and 4, or a sequence complementary thereto or a fragment thereof.

T05290" TEE29250

The invention also relates to isolated, purified, or recombinant polynucleotides comprising a contiguous span of at least 12, 15, 18, 20, 25, 30, 35, 40, 50, 60, 70, 80, 90, 100, 150, 200, 500, or 1000 nucleotides of a nucleotide sequence selected from the group consisting of SEQ ID Nos 3 and 4, or the complements thereof. Particularly preferred nucleic acids of the invention include isolated, purified, or recombinant polynucleotides comprising a contiguous span of at least 12, 15, 18, 20, 25, 30, 35, 40, 50, 60, 70, 80, 90, 100, 150, 200, 500, or 1000 nucleotides of SEQ ID No 3 or the complements thereof, wherein said contiguous span comprises at least 1, 2, 3, 5, or 10 of the following nucleotide positions of SEQ ID No 3: 1-500, 501-1000, 1001-1500, 1501-2000, 2001-2500, 2501-3000, 3001-3500, and 3501-3983. Additionally preferred nucleic acids of the invention include isolated, purified, or recombinant polynucleotides comprising a contiguous span of at least 12, 15, 18, 20, 25, 30, 35, 40, 50, 60, 70, 80, 90, 100, 150, 200, 500, or 1000 nucleotides of SEQ ID No 4 or the complements thereof, wherein said contiguous span comprises at least 1, 2, 3, 5, or 10 of the following nucleotide positions of SEQ ID No 4: 1-500, 501-1000, 1001-1500, 1501-2000, 2001-2500, 2501-3000, 3001-3500, and 3501-3988. Such a nucleic acid is notably useful as polynucleotide probe or primer specific for the *TBC-1* gene or the *TBC-1* mRNAs and cDNAs.

While this section is entitled "*TBC-1* cDNA Sequences," it should be noted that nucleic acid fragments of any size and sequence may also be comprised by the polynucleotides described in this section, flanking the genomic sequences of *TBC-1* on either side or between two or more such genomic sequences.

### Coding Regions

The *TBC-1* open reading frame is contained in the two *TBC-1* mRNA molecules of about 4 kilobases isolated by the inventors.

More precisely, the effective *TBC-1* coding sequence is comprised between the nucleotide at position 171 and the nucleotide at position 3725 of SEQ ID No 3, and between the nucleotide at position 176 and the nucleotide at position 3730 of the nucleotide sequence of SEQ ID No 4.

The invention further provides a purified or isolated nucleic acid comprising a polynucleotide selected from the group consisting of a polynucleotide comprising a nucleic acid sequence located between the nucleotide at position 171 and the nucleotide at position 3725 of SEQ ID No 3, and a polynucleotide comprising a nucleic acid sequence located between the nucleotide at position 176 and the nucleotide at position 3730 of SEQ ID No 4 or a variant or fragment thereof or a sequence complementary thereto.

The present invention concerns a purified or isolated nucleic acid encoding a human TBC-1 protein, wherein said TBC-1 protein comprises an amino acid sequence of SEQ ID No 5, a nucleotide sequence complementary thereto, a fragment or a variant thereof. The present invention also embodies isolated, purified, and recombinant polynucleotides which encode a polypeptides comprising a contiguous span of at least 6 amino acids, preferably at least 8 or 10 amino acids,

more preferably at least 12, 15, 20, 25, 30, 40, 50, or 100 amino acids of SEQ ID No 5. In a preferred embodiment, the present invention embodies isolated, purified, and recombinant polynucleotides which encode a polypeptides comprising a contiguous span of at least 6 amino acids, preferably at least 8 or 10 amino acids, more preferably at least 12, 15, 20, 25, 30, 40, 50, or 100 amino acids of SEQ ID No 5 wherein said contiguous span includes at least 1, 2, 3, 5 or 10 of the following amino acid positions in SEQ ID No 5: 1-300, 301-600, 601-900, and 901-1168.

The above disclosed polynucleotide that contains only coding sequences derived from the *TBC-1* ORF may be expressed in a desired host cell or a desired host organism, when said polynucleotide is placed under the control of suitable expression signals. Such a polynucleotide, when placed under the suitable expression signals, may be inserted in a vector for its expression.

### Regulatory Sequences Of *TBC-1*

The invention further deals with a purified or isolated nucleic acid comprising the nucleotide sequence of a regulatory region which is located either upstream of the first exon of the *TBC-1* gene and which is contained in the *TBC-1* genomic sequence of SEQ ID No 1, or downstream of the last exon of the *TBC-1* gene and which is contained in the *TBC-1* genomic sequence of SEQ ID No 2.

The 5'-regulatory sequence of the *TBC-1* gene is localized between the nucleotide in position 1 and the nucleotide in position 2000 of the nucleotide sequence of SEQ ID No 1. The 3'-regulatory sequence of the *TBC-1* gene is localized between nucleotide position 97961 and nucleotide position 99960 of SEQ ID No 2.

Polynucleotides derived from the 5' and 3' regulatory regions are useful in order to detect the presence of at least a copy of a nucleotide sequence of SEQ ID Nos 1 or 2 or a fragment thereof in a test sample.

The promoter activity of the 5' regulatory regions contained in *TBC-1* can be assessed as described below.

Genomic sequences lying upstream of the *TBC-1* Exons are cloned into a suitable promoter reporter vector, such as the pSEAP-Basic, pSEAP-Enhancer, p $\beta$ gal-Basic, p $\beta$ gal-Enhancer, or pEGFP-1 Promoter Reporter vectors available from Clontech. Briefly, each of these promoter reporter vectors include multiple cloning sites positioned upstream of a reporter gene encoding a readily assayable protein such as secreted alkaline phosphatase, beta galactosidase, or green fluorescent protein. The sequences upstream of the *TBC-1* coding region are inserted into the cloning sites upstream of the reporter gene in both orientations and introduced into an appropriate host cell. The level of reporter protein is assayed and compared to the level obtained from a vector which lacks an insert in the cloning site. The presence of an elevated expression level in the vector containing the insert with respect to the control vector indicates the presence of a promoter in the insert. If necessary, the upstream sequences can be cloned into vectors which contain an enhancer

for increasing transcription levels from weak promoter sequences. A significant level of expression above that observed with the vector lacking an insert indicates that a promoter sequence is present in the inserted upstream sequence.

Promoter sequences within the upstream genomic DNA may be further defined by  
5 constructing nested deletions in the upstream DNA using conventional techniques such as Exonuclease III digestion. The resulting deletion fragments can be inserted into the promoter reporter vector to determine whether the deletion has reduced or obliterated promoter activity. In this way, the boundaries of the promoters may be defined. If desired, potential individual regulatory sites within the promoter may be identified using site directed mutagenesis or linker scanning to  
10 obliterate potential transcription factor binding sites within the promoter, individually or in combination. The effects of these mutations on transcription levels may be determined by inserting the mutations into the cloning sites in the promoter reporter vectors.

Thus, the minimal size of the promoter of the *TBC-1* gene can be determined through the measurement of *TBC-1* expression levels. For this assay, an expression vector comprising  
15 decreasing sizes from the promoter generally ranging from 2 kb to 100 bp, with a 3' end which is constant, operably linked to *TBC-1* coding sequence or to a reporter gene is used. Cells, which are preferably prostate cells and more preferably prostate cancer cells, are transfected with this vector and the expression level of the gene is assessed.

The strength and the specificity of the promoter of the *TBC-1* gene can be assessed through  
20 the expression levels of the gene operably linked to this promoter in different types of cells and tissues. In one embodiment, the efficacy of the promoter of the *TBC-1* gene is assessed in normal and cancer cells. In a preferred embodiment, the efficacy of the promoter of the *TBC-1* gene is assessed in normal prostate cells and in prostate cancer cells which can present different degrees of malignancy.

25 Polynucleotides carrying the regulatory elements located both at the 5' end and at the 3' end of the *TBC-1* cDNAs may be advantageously used to control the transcriptional and translational activity of an heterologous polynucleotide of interest.

Thus, the present invention also concerns a purified or isolated nucleic acid comprising a polynucleotide which is selected from the group consisting of the 5' and 3' regulatory regions, or a  
30 sequence complementary thereto or a biologically active fragment or variant thereof. "5' regulatory region" refers to the nucleotide sequence located between positions 1 and 2000 of SEQ ID No 1. "3' regulatory region" refers to the nucleotide sequence located between positions 97961 and 99960 of SEQ ID No 2.

The invention also pertains to a purified or isolated nucleic acid comprising a  
35 polynucleotide having at least 95% nucleotide identity with a polynucleotide selected from the group consisting of the 5' and 3' regulatory regions, advantageously 99 % nucleotide identity, preferably 99.5% nucleotide identity and most preferably 99.8% nucleotide identity with a

polynucleotide selected from the group consisting of the 5' and 3' regulatory regions, or a sequence complementary thereto or a variant thereof or a biologically active fragment thereof.

Another object of the invention consists of purified, isolated or recombinant nucleic acids comprising a polynucleotide that hybridizes, under the stringent hybridization conditions defined herein, with a polynucleotide selected from the group consisting of the nucleotide sequences of the 5'- and 3' regulatory regions, or a sequence complementary thereto or a variant thereof or a biologically active fragment thereof.

The 5'UTR and 3'UTR regions of a gene are of particular importance in that they often comprise regulatory elements which can play a role in providing appropriate expression levels, particularly through the control of mRNA stability.

A 5' regulatory polynucleotide of the invention may include the 5'-UTR located between the nucleotide at position 1 and the nucleotide at position 170 of SEQ ID No 3, or a biologically active fragment or variant thereof.

Alternatively, a 5'-regulatory polynucleotide of the invention may include the 5'-UTR located between the nucleotide at position 1 and the nucleotide at position 175 of SEQ ID No 4, or a biologically active fragment or variant thereof.

A 3' regulatory polynucleotide of the invention may include the 3'-UTR located between the nucleotide at position 3726 and the nucleotide at position 3983 of SEQ ID No 4, or a biologically active fragment or variant thereof.

Thus, the invention also pertains to a purified or isolated nucleic acid which is selected from the group consisting of :

- a) a nucleic acid comprising the nucleotide sequence of the 5' regulatory region;
- b) a nucleic acid comprising a biologically active fragment or variant of the nucleic acid of the 5' regulatory region.

Preferred fragments of the nucleic acid of the 5' regulatory region have a length of about 1000 nucleotides, more particularly of about 400 nucleotides, more preferably of about 200 nucleotides and most preferably about 100 nucleotides. More particularly, the invention further includes specific elements within this regulatory region, these elements preferably including the promoter region.

Preferred fragments of the 3' regulatory region are at least 50, 100, 150, 200, 300 or 400 bases in length.

By a "biologically active fragment or variant" of a *TBC-1* regulatory polynucleotide according to the present invention is intended a polynucleotide comprising or alternatively consisting in a fragment of said polynucleotide which is functional as a regulatory region for expressing a recombinant polypeptide or a recombinant polynucleotide in a recombinant cell host.

For the purpose of the invention, a nucleic acid or polynucleotide is "functional" as a regulatory region for expressing a recombinant polypeptide or a recombinant polynucleotide if said

regulatory polynucleotide contains nucleotide sequences which contain transcriptional and translational regulatory information, and if such sequences are "operatively linked" to nucleotide sequences which encode the desired polypeptide or the desired polynucleotide. An operable linkage is a linkage in which the regulatory nucleic acid and the DNA sequence sought to be expressed are

5 linked in such a way as to permit gene expression.

In order, to identify the relevant biologically active polynucleotide derivatives of the 5' or 3' regulatory region, the one skill in the art will refer to the book of Sambrook et al. (Sambrook, 1989) in order to use a recombinant vector carrying a marker gene (i.e. beta galactosidase, chloramphenicol acetyl transferase, etc.) the expression of which will be detected when placed

10 under the control of a biologically active derivative polynucleotide of the 5' or 3' regulatory region.

Regulatory polynucleotides of the invention may be prepared from any of the nucleotide sequences of SEQ ID Nos 1 or 2 by cleavage using the suitable restriction enzymes, the one skill in the art being guided by the book of Sambrook et al. (1989). Regulatory polynucleotides may also be prepared by digestion of any of the nucleotide sequences of SEQ ID Nos 1 or 2 by an exonuclease

15 enzyme, such as Bal31 (Wabiko et al., 1986). These regulatory polynucleotides can also be prepared by chemical synthesis, as described elsewhere in the specification, when the synthesis of oligonucleotide probes or primers is disclosed.

The regulatory polynucleotides according to the invention may be advantageously part of a recombinant expression vector that may be used to express a coding sequence in a desired host cell

20 or host organism. The recombinant expression vectors according to the invention are described elsewhere in the specification.

The invention also encompasses a polynucleotide comprising :

- a) a nucleic acid comprising a regulatory nucleotide sequence of the 5' regulatory region, or a biologically active fragment or variant thereof;
- 25 b) a polynucleotide encoding a desired polypeptide or nucleic acid, operably linked to the nucleic acid comprising a regulatory nucleotide sequence of the 5' regulatory region, or its biologically active fragment or variant.
- c) Optionally, a nucleic acid comprising a 3' regulatory polynucleotide, preferably a 3' regulatory polynucleotide of the invention.

30 The desired polypeptide encoded by the above described nucleic acid may be of various nature or origin, encompassing proteins of prokaryotic or eukaryotic origin. Among the polypeptides expressed under the control of a *TBC-1* regulatory region, it may be cited bacterial, fungal or viral antigens. Are also encompassed eukaryotic proteins such as intracellular proteins, such as "house keeping" proteins, membrane-bound proteins, like receptors, and secreted proteins

35 like the numerous endogenous mediators such as cytokines.

The desired nucleic acid encoded by the above described polynucleotide, usually a RNA molecule, may be complementary to a *TBC-1* coding sequence and thus useful as an antisense polynucleotide.

Such a polynucleotide may be included in a recombinant expression vector in order to  
5 express a desired polypeptide or a desired polynucleotide in host cell or in a host organism. Suitable recombinant vectors that contain a polynucleotide such as described hereinbefore are disclosed elsewhere in the specification.

### **TBC-1 Polypeptide And Peptide Fragments Thereof**

It is now easy to produce proteins in high amounts by genetic engineering techniques  
10 through expression vectors such as plasmids, phages or phagemids. The polynucleotide that code for one the polypeptides of the present invention is inserted in an appropriate expression vector in order to produce the polypeptide of interest *in vitro*.

Thus, the present invention also concerns a method for producing one of the polypeptides described herein, and especially a polypeptide of SEQ ID No 5 or a fragment or a variant thereof,  
15 wherein said method comprises the steps of :

a) culturing, in an appropriate culture medium, a cell host previously transformed or transfected with the recombinant vector comprising a nucleic acid encoding a TBC-1 polypeptide, or a fragment or a variant thereof;

b) harvesting the culture medium thus conditioned or lyse the cell host, for example by  
20 sonication or by an osmotic shock;

c) separating or purifying, from the said culture medium, or from the pellet of the resultant host cell lysate the thus produced polypeptide of interest.

d) Optionally characterizing the produced polypeptide of interest.

In a specific embodiment of the above method, step a) is preceded by a step wherein the  
25 nucleic acid coding for a TBC-1 polypeptide, or a fragment or a variant thereof, is inserted in an appropriate vector, optionally after an appropriate cleavage of this amplified nucleic acid with one or several restriction endonucleases. The nucleic acid coding for a TBC-1 polypeptide or a fragment or a variant thereof may be the resulting product of an amplification reaction using a pair of primers according to the invention (by SDA, TAS, 3SR NASBA, TMA etc.).

30 The polypeptides according to the invention may be characterized by binding onto an immunoaffinity chromatography column on which polyclonal or monoclonal antibodies directed to a polypeptide of SEQ ID No 5, or a fragment or a variant thereof, have previously been immobilized.

Purification of the recombinant proteins or peptides according to the present invention may  
35 be carried out by passage onto a Nickel or Copper affinity chromatography column. The Nickel chromatography column may contain the Ni-NTA resin (Porath et al., 1975).



The polypeptides or peptides thus obtained may be purified, for example by high performance liquid chromatography, such as reverse phase and/or cationic exchange HPLC, as described by Rougeot et al. (1994). The reason to prefer this kind of peptide or protein purification is the lack of byproducts found in the elution samples which renders the resultant purified protein or peptide more suitable for a therapeutic use.

Another object of the present invention consists in a purified or isolated TBC-1 polypeptide or a fragment or a variant thereof.

In a preferred embodiment, the TBC-1 polypeptide comprises an amino acid sequence of SEQ ID No 5 or a fragment or a variant thereof. The present invention also embodies isolated, purified, and recombinant polypeptides comprising a contiguous span of at least 6 amino acids, preferably at least 8 to 10 amino acids, more preferably at least 12, 15, 20, 25, 30, 40, 50, 100, 150 or 200 amino acids of SEQ ID No 5. The present invention also embodies isolated, purified, and recombinant polypeptides comprising a contiguous span of at least 6 amino acids, preferably at least 8 to 10 amino acids, more preferably at least 12, 15, 20, 25, 30, 40, 50, 100, 150 or 200 amino acids of SEQ ID No 5, wherein said contiguous span includes at least 1, 2, 3, 5 or 10 of the following amino acid positions: 1-200, 201-400, 401-600, 601-800, 801-1000, 1001-1168.

The invention also encompasses a purified, isolated, or recombinant polypeptides comprising an amino acid sequence having at least 90, 95, 98 or 99% amino acid identity with the amino acid sequence of SEQ ID No 5 or a fragment thereof.

The TBC-1 polypeptide of the invention possesses amino acid homologies as regards to the murine TBC-1 protein of 1141 amino acids in length which is described in US Patent No US 5,700,927. The TBC-1 protein of the invention also possesses some homologies with two other proteins : the Pollux drosophila protein (Zhang et al., 1996) and the CDC16 protein from *Caenorhabditis elegans* (Wilson et al., 1994). Figure 1 represents an amino acid alignment of a portion of the amino acid sequence of the TBC-1 protein of SEQ ID No 5 with other proteins sharing amino acid homology with TBC-1. The upper line shows the whole amino acid sequence of the murine tbc-1 protein described in US Patent No US 5,700,927; the second line represents part of the amino acid sequence of the TBC-1 protein of SEQ ID No 5; the third line (Genbank access No : *dmu50542*) depicts the amino acid sequence of the Pollux protein mentioned above; the fourth line (Genbank access No : *celf35h12*) shows the amino acid sequence of the *C. elegans* protein mentioned above; the fifth line presents positions in which consensus amino acids are identified, i.e. amino acids shared by the sequences presented in the four upper lines, when present.

The TBC-1 polypeptide of the amino acid sequence of SEQ ID No 5 has 1168 amino acids in length. The TBC-1 polypeptide includes a "TBC domain" which is spanning from the amino acid in position 786 to the amino acid in position 974 of the amino acid sequence of SEQ ID No 5. This TBC domain is represented in Figure 1 as a grey area spanning from the amino acid numbered 758 to the amino acid numbered 949. This TBC domain is likely to regulate protein-protein interactions.

Moreover, the TBC-1 TBC domain includes the amino acid sequence EVGYCQGL, spanning from the amino acid in position 886 to the amino acid in position 893 of the amino acid sequence of SEQ ID No 5. The EVGYCQGL amino acid sequence spans from the amino acid numbered 861 to the amino acid numbered 868 of Figure 1. This site may interact with a kinase. Based on the structural similarity to *cdc16*, a yeast regulator of mitosis, TBC-1 is likely to regulate mitosis and cytokinesis by interacting with other proteins which also participate with the regulation of mitosis, cytokinesis and septum formation.

Preferred polypeptides of the invention comprise the TBC domain of TBC-1, or alternatively at least the EVGYCQGL amino acid sequence motif.

10 A further object of the present invention concerns a purified or isolated polypeptide which is encoded by a nucleic acid comprising a nucleotide sequence selected from the group consisting of SEQ ID Nos 1, 2, 3, and 4 or fragments or variants thereof.

A single variant molecule of the TBC-1 protein is explicitly excluded from the scope of the present invention, which is a polypeptide having the same amino acid sequence than the murine *tbc1* protein described in the US Patent No 5,700,927.

Amino acid deletions, additions or substitutions in the TBC-1 protein are preferably located outside of the TBC domain as defined above. Most preferably, a mutated TBC-1 protein has an intact "EVGYCQGL" amino acid motif.

Such a mutated TBC-1 protein may be the target of diagnostic tools, such as specific monoclonal or polyclonal antibodies, useful for detecting the mutated TBC-1 protein in a sample.

The invention also encompasses a TBC-1 polypeptide or a fragment or a variant thereof in which at least one peptide bound has been modified as described in the "Definitions" section.

#### Antibodies That Bind *TBC-1* Polypeptides of the Invention

Any TBC-1 polypeptide or whole protein may be used to generate antibodies capable of specifically binding to an expressed TBC-1 protein or fragments thereof as described.

One antibody composition of the invention is capable of specifically binding or specifically bind to the variant of the TBC-1 protein of SEQ ID No 5. For an antibody composition to specifically bind to TBC-1, it must demonstrate at least a 5%, 10%, 15%, 20%, 25%, 50%, or 100% greater binding affinity for TBC-1 protein than for another protein in an ELISA, RIA, or other antibody-based binding assay.

In a preferred embodiment, the invention concerns antibody compositions, either polyclonal or monoclonal, capable of selectively binding, or selectively bind to an epitope-containing a polypeptide comprising a contiguous span of at least 6 amino acids, preferably at least 8 to 10 amino acids, more preferably at least 12, 15, 20, 25, 30, 40, 50, 100, 150 or 200 amino acids of SEQ ID No 5; Optionally said epitope comprises at least 1, 2, 3, 5 or 10 of the following amino acid positions : 1-200, 201-400, 401-600, 601-800, 801-1000, 1001-1168.

The invention also concerns a purified or isolated antibody capable of specifically binding to a mutated TBC-1 protein or to a fragment or variant thereof comprising an epitope of the mutated TBC-1 protein. In another preferred embodiment, the present invention concerns an antibody capable of binding to a polypeptide comprising at least 10 consecutive amino acids of a TBC-1 protein and including at least one of the amino acids which can be encoded by the trait causing mutations.

In a preferred embodiment, the invention concerns the use in the manufacture of antibodies of a polypeptide comprising a contiguous span of at least 6 amino acids, preferably at least 8 to 10 amino acids, more preferably at least 12, 15, 20, 25, 30, 40, 50, 100, 150 or 200 amino acids of SEQ ID No 5; Optionally said polypeptide comprises at least 1, 2, 3, 5 or 10 of the following amino acid positions : 1-200, 201-400, 401-600, 601-800, 801-1000, 1001-1168.

The antibodies of the invention may be labeled by any one of the radioactive, fluorescent or enzymatic labels known in the art.

The TBC-1 polypeptide of SEQ ID No 5 or a fragment thereof can be used for the preparation of polyclonal or monoclonal antibodies.

The TBC-1 polypeptide expressed from a DNA sequence comprising at least one of the nucleic acid sequences of SEQ ID Nos 1, 2, 3 and 4 may also be used to generate antibodies capable of specifically binding to the TBC-1 polypeptide of SEQ ID No 5 or a fragment thereof.

Preferred antibodies according to the invention are prepared using TBC-1 peptide fragments that do not comprise the EVGYCQGL amino acid motif.

Other preferred antibodies of the invention are prepared using TBC-1 peptide fragments that do not comprise the TBC domain defined elsewhere in the specification.

The antibodies may be prepared from hybridomas according to the technique described by Kohler and Milstein in 1975. The polyclonal antibodies may be prepared by immunization of a mammal, especially a mouse or a rabbit, with a polypeptide according to the invention that is combined with an adjuvant of immunity, and then by purifying of the specific antibodies contained in the serum of the immunized animal on an affinity chromatography column on which has previously been immobilized the polypeptide that has been used as the antigen.

The present invention also includes, chimeric single chain Fv antibody fragments (Martineau et al., 1998), antibody fragments obtained through phage display libraries (Ridder et al., 1995; Vaughan et al., 1995) and humanized antibodies (Reinmann et al., 1997; Leger et al., 1997).

Antibody preparations prepared according to either protocol are useful in quantitative immunoassays which determine concentrations of antigen-bearing substances in biological samples; they are also used semi-quantitatively or qualitatively to identify the presence of antigen in a biological sample. The antibodies may also be used in therapeutic compositions for killing cells expressing the protein or reducing the levels of the protein in the body.

Consequently, the invention is also directed to a method for detecting specifically the presence of a TBC-1 polypeptide according to the invention in a biological sample, said method comprising the following steps :

- a) bringing into contact the biological sample with a polyclonal or monoclonal antibody that specifically binds a TBC-1 polypeptide comprising an amino acid sequence of SEQ ID No 5, or to a peptide fragment or variant thereof; and
- b) detecting the antigen-antibody complex formed.

The invention also concerns a diagnostic kit for detecting *in vitro* the presence of a TBC-1 polypeptide according to the present invention in a biological sample, wherein said kit comprises:

- a) a polyclonal or monoclonal antibody that specifically binds a TBC-1 polypeptide comprising an amino acid sequence of SEQ ID No 5, or to a peptide fragment or variant thereof, optionally labeled;
- b) a reagent allowing the detection of the antigen-antibody complexes formed, said reagent carrying optionally a label, or being able to be recognized itself by a labeled reagent, more particularly in the case when the above-mentioned monoclonal or polyclonal antibody is not labeled by itself.

### ***TBC-1*-Related Biallelic Markers**

The inventors have discovered nucleotide polymorphisms located within the genomic DNA containing the *TBC-1* gene, and among them SNP that are also termed biallelic markers. The biallelic markers of the invention can be used for example for the generation of genetic map, the linkage analysis, the association studies.

#### **A- Identification Of *TBC-1*-related Biallelic Markers**

There are two preferred methods through which the biallelic markers of the present invention can be generated. In a first method, DNA samples from unrelated individuals are pooled together, following which the genomic DNA of interest is amplified and sequenced. The nucleotide sequences thus obtained are then analyzed to identify significant polymorphisms.

One of the major advantages of this method resides in the fact that the pooling of the DNA samples substantially reduces the number of DNA amplification reactions and sequencing which must be carried out. Moreover, this method is sufficiently sensitive so that a biallelic marker obtained therewith usually shows a sufficient degree of informativeness for conducting association studies.

In a second method for generating biallelic markers, the DNA samples are not pooled and are therefore amplified and sequenced individually. The resulting nucleotide sequences obtained are then also analyzed to identify significant polymorphisms.

It will readily be appreciated that when this second method is used, a substantially higher number of DNA amplification reactions must be carried out. It will further be appreciated that

09762311-06904  
T.05220 T.29260

including such potentially less informative biallelic markers in association studies to identify potential genetic associations with a trait may allow in some cases the direct identification of causal mutations, which may, depending on their penetrance, be rare mutations. This method is usually preferred when biallelic markers need to be identified in order to perform association studies within

5 candidate genes.

In both methods, the genomic DNA samples from which the biallelic markers of the present invention are generated are preferably obtained from unrelated individuals corresponding to a heterogeneous population of known ethnic background, or from familial cases.

The number of individuals from whom DNA samples are obtained can vary substantially, preferably from about 10 to about 1000, preferably from about 50 to about 200 individuals. It is usually preferred to collect DNA samples from at least about 100 individuals in order to have sufficient polymorphic diversity in a given population to generate as many markers as possible and to generate statistically significant results.

As for the source of the genomic DNA to be subjected to analysis, any test sample can be foreseen without any particular limitation. The preferred source of genomic DNA used in the context of the present invention is the peripheral venous blood of each donor.

The techniques of DNA extraction are well-known to the skilled technician. Details of a preferred embodiment are provided in Example 2.

DNA samples can be pooled or unpooled for the amplification step. DNA amplification techniques are well-known to those skilled in the art.

Amplification techniques that can be used in the context of the present invention include, but are not limited to, the ligase chain reaction (LCR) described in EP-A- 320 308, WO 9320227 and EP-A-439 182, the polymerase chain reaction (PCR, RT-PCR) and techniques such as the nucleic acid sequence based amplification (NASBA) described in Guatelli J.C., et al.(1990) and in Compton J.(1991), Q-beta amplification as described in European Patent Application No 4544610, strand displacement amplification as described in Walker et al.(1996) and EP A 684 315 and, target mediated amplification as described in PCT Publication WO 9322461.

LCR and Gap LCR are exponential amplification techniques, both depend on DNA ligase to join adjacent primers annealed to a DNA molecule. In Ligase Chain Reaction (LCR), probe pairs are used which include two primary (first and second) and two secondary (third and fourth) probes, all of which are employed in molar excess to target. The first probe hybridizes to a first segment of the target strand and the second probe hybridizes to a second segment of the target strand, the first and second segments being contiguous so that the primary probes abut one another in 5' phosphate-3'hydroxyl relationship, and so that a ligase can covalently fuse or ligate the two probes into a fused product. In addition, a third (secondary) probe can hybridize to a portion of the first probe and a fourth (secondary) probe can hybridize to a portion of the second probe in a similar abutting fashion. Of course, if the target is initially double stranded, the secondary probes also will

hybridize to the target complement in the first instance. Once the ligated strand of primary probes is separated from the target strand, it will hybridize with the third and fourth probes, which can be ligated to form a complementary, secondary ligated product. It is important to realize that the ligated products are functionally equivalent to either the target or its complement. By repeated  
5 cycles of hybridization and ligation, amplification of the target sequence is achieved. A method for multiplex LCR has also been described (WO 9320227). Gap LCR (GLCR) is a version of LCR where the probes are not adjacent but are separated by 2 to 3 bases.

For amplification of mRNAs, it is within the scope of the present invention to reverse transcribe mRNA into cDNA followed by polymerase chain reaction (RT-PCR); or, to use a single  
10 enzyme for both steps as described in U.S. Patent No. 5,322,770 or, to use Asymmetric Gap LCR (RT-AGLCR) as described by Marshall et al.(1994). AGLCR is a modification of GLCR that allows the amplification of RNA.

The PCR technology is the preferred amplification technique used in the present invention. A variety of PCR techniques are familiar to those skilled in the art. For a review of PCR  
15 technology, see White (1997) and the publication entitled "PCR Methods and Applications" (1991, Cold Spring Harbor Laboratory Press). In each of these PCR procedures, PCR primers on either side of the nucleic acid sequences to be amplified are added to a suitably prepared nucleic acid sample along with dNTPs and a thermostable polymerase such as Taq polymerase, Pfu polymerase, or Vent polymerase. The nucleic acid in the sample is denatured and the PCR primers are  
20 specifically hybridized to complementary nucleic acid sequences in the sample. The hybridized primers are extended. Thereafter, another cycle of denaturation, hybridization, and extension is initiated. The cycles are repeated multiple times to produce an amplified fragment containing the nucleic acid sequence between the primer sites. PCR has further been described in several patents including US Patents 4,683,195; 4,683,202; and 4,965,188.

25 The PCR technology is the preferred amplification technique used to identify new biallelic markers. A typical example of a PCR reaction suitable for the purposes of the present invention is provided in Example 3.

One of the aspects of the present invention is a method for the amplification of a *TBC-1* gene, particularly the genomic sequences of SEQ ID Nos 1 and 2 or of the cDNA sequence of SEQ  
30 ID Nos 3 or 4 or a fragment or variant thereof in a test sample, preferably using the PCR technology. The method comprises the steps of contacting a test sample suspected of containing the target *TBC-1* sequence or portion thereof with amplification reaction reagents comprising a pair of amplification primers.

Thus, the present invention also relates to a method for the amplification of a *TBC-1* gene  
35 sequence, particularly of a fragment of the genomic sequence of SEQ ID No 1 or of the cDNA sequence of SEQ ID No 2 or 3, or a fragment or a variant thereof in a test sample, said method comprising the steps of :

09762341.062504

a) contacting a test sample suspected of containing the targeted *TBC-1* gene sequence or portion thereof with amplification reaction reagents comprising a pair of amplification primers located on either side of the *TBC-1* region to be amplified, and

b) optionally, detecting the amplification products.

5 The invention also concerns a kit for the amplification of a *TBC-1* gene sequence, particularly of a portion of the genomic sequence of SEQ ID Nos 1 or 2, or of the cDNA sequence of SEQ ID Nos 3 or 4, or a variant thereof in a test sample, wherein said kit comprises:

a) a pair of oligonucleotide primers located on either side of the *TBC-1* region to be amplified;

10 b) optionally, the reagents necessary for performing the amplification reaction.

In one embodiment of the above amplification method and kit, the amplification product is detected by hybridization with a labeled probe having a sequence which is complementary to the amplified region. In another embodiment of the above amplification method and kit, primers comprise a sequence which is selected from the group consisting of B1 to B15, C1 to C15, D1 to

15 D19, and E1 to E19.

In a first embodiment of the present invention, biallelic markers are identified using genomic sequence information generated by the inventors. Sequenced genomic DNA fragments are used to design primers for the amplification of 500 bp fragments. These 500 bp fragments are amplified from genomic DNA and are scanned for biallelic markers. Primers may be designed using the OSP software (Hillier L. and Green P., 1991). All primers may contain, upstream of the specific target bases, a common oligonucleotide tail that serves as a sequencing primer. Those skilled in the art are familiar with primer extensions, which can be used for these purposes.

20 Preferred primers, useful for the amplification of genomic sequences encoding the candidate genes, focus on promoters, exons and splice sites of the genes. A biallelic marker presents a higher probability to be an eventual causal mutation if it is located in these functional regions of the gene. Preferred amplification primers of the invention include the nucleotide sequences of B1 to B15 and C1 to C15 further detailed in Example 3.

The amplification products generated as described above with the primers of the invention are then sequenced using methods known and available to the skilled technician. Preferably, the amplified DNA is subjected to automated dideoxy terminator sequencing reactions using a dye-primer cycle sequencing protocol. Following gel image analysis and DNA sequence extraction, sequence data are automatically processed with adequate software to assess sequence quality.

A polymorphism analysis software is used that detects the presence of biallelic sites among individual or pooled amplified fragment sequences. Polymorphism search is based on the presence of superimposed peaks in the electrophoresis pattern. These peaks which present distinct colors correspond to two different nucleotides at the same position on the sequence. The polymorphism has to be detected on both strands for validation.

097231-0690  
T09990-TE29460

19 biallelic markers were found in the *TBC-1* gene. They are detailed in the Table 2. They are located in intronic regions.

#### B- Genotyping Of *TBC-1*-Related Biallelic Markers

The polymorphisms identified above can be further confirmed and their respective frequencies can be determined through various methods using the previously described primers and probes. These methods can also be useful for genotyping either new populations in association studies or linkage analysis or individuals in the context of detection of alleles of biallelic markers which are known to be associated with a given trait. The genotyping of the biallelic markers is also important for the mapping. Those skilled in the art should note that the methods described below can be equally performed on individual or pooled DNA samples.

Once a given polymorphic site has been found and characterized as a biallelic marker as described above, several methods can be used in order to determine the specific allele carried by an individual at the given polymorphic base.

The identification of biallelic markers described previously allows the design of appropriate oligonucleotides, which can be used as probes and primers, to amplify a *TBC-1* gene containing the polymorphic site of interest and for the detection of such polymorphisms.

The biallelic markers according to the present invention may be used in methods for the identification and characterization of an association between alleles for one or several biallelic markers of the sequence of the *TBC-1* gene and a trait.

The identified polymorphisms, and consequently the biallelic markers of the invention, may be used in methods for the detection in an individual of *TBC-1* alleles associated with a trait, more particularly a trait related to a cell differentiation or abnormal cell proliferation disorders, and most particularly a trait related to cancer diseases, specifically prostate cancer.

In one embodiment the invention encompasses methods of genotyping comprising determining the identity of a nucleotide at a *TBC-1*-related biallelic marker or the complement thereof in a biological sample; optionally, wherein said *TBC-1*-related biallelic marker is selected from the group consisting of A1 to A19, and the complements thereof, or optionally the biallelic markers in linkage disequilibrium therewith; optionally, wherein said biological sample is derived from a single subject; optionally, wherein the identity of the nucleotides at said biallelic marker is determined for both copies of said biallelic marker present in said individual's genome; optionally, wherein said biological sample is derived from multiple subjects; Optionally, the genotyping methods of the invention encompass methods with any further limitation described in this disclosure, or those following, specified alone or in any combination; Optionally, said method is performed *in vitro*; optionally, further comprising amplifying a portion of said sequence comprising the biallelic marker prior to said determining step; Optionally, wherein said amplifying is performed by PCR, LCR, or replication of a recombinant vector comprising an origin of replication and said fragment in a host cell; optionally, wherein said determining is performed by a



hybridization assay, a sequencing assay, a microsequencing assay, or an enzyme-based mismatch detection assay.

#### Source of Nucleic Acids for genotyping

Any source of nucleic acids, in purified or non-purified form, can be utilized as the starting nucleic acid, provided it contains or is suspected of containing the specific nucleic acid sequence desired. DNA or RNA may be extracted from cells, tissues, body fluids and the like as described above. While nucleic acids for use in the genotyping methods of the invention can be derived from any mammalian source, the test subjects and individuals from which nucleic acid samples are taken are generally understood to be human.

#### 10 Amplification Of DNA Fragments Comprising Biallelic Markers

Methods and polynucleotides are provided to amplify a segment of nucleotides comprising one or more biallelic marker of the present invention. It will be appreciated that amplification of DNA fragments comprising biallelic markers may be used in various methods and for various purposes and is not restricted to genotyping. Nevertheless, many genotyping methods, although not all, require the previous amplification of the DNA region carrying the biallelic marker of interest. Such methods specifically increase the concentration or total number of sequences that span the biallelic marker or include that site and sequences located either distal or proximal to it. Diagnostic assays may also rely on amplification of DNA segments carrying a biallelic marker of the present invention. Amplification of DNA may be achieved by any method known in the art. Amplification techniques are described above in the section entitled, "Identification of *TBC-1*-related biallelic markers."

Some of these amplification methods are particularly suited for the detection of single nucleotide polymorphisms and allow the simultaneous amplification of a target sequence and the identification of the polymorphic nucleotide as it is further described below.

The identification of biallelic markers as described above allows the design of appropriate oligonucleotides, which can be used as primers to amplify DNA fragments comprising the biallelic markers of the present invention. Amplification can be performed using the primers initially used to discover new biallelic markers which are described herein or any set of primers allowing the amplification of a DNA fragment comprising a biallelic marker of the present invention.

In some embodiments the present invention provides primers for amplifying a DNA fragment containing one or more biallelic markers of the present invention. Preferred amplification primers are listed in Example 2. It will be appreciated that the primers listed are merely exemplary and that any other set of primers which produce amplification products containing one or more biallelic markers of the present invention are also of use.

The spacing of the primers determines the length of the segment to be amplified. In the context of the present invention, amplified segments carrying biallelic markers can range in size from at least about 25 bp to 35 kbp. Amplification fragments from 25-3000 bp are typical,

0976341-06501  
T05530-TE9260

fragments from 50-1000 bp are preferred and fragments from 100-600 bp are highly preferred. It will be appreciated that amplification primers for the biallelic markers may be any sequence which allow the specific amplification of any DNA fragment carrying the markers. Amplification primers may be labeled or immobilized on a solid support as described in "Oligonucleotide probes and  
5 primers".

#### Methods of Genotyping DNA samples for Biallelic Markers

Any method known in the art can be used to identify the nucleotide present at a biallelic marker site. Since the biallelic marker allele to be detected has been identified and specified in the present invention, detection will prove simple for one of ordinary skill in the art by employing any  
10 of a number of techniques. Many genotyping methods require the previous amplification of the DNA region carrying the biallelic marker of interest. While the amplification of target or signal is often preferred at present, ultrasensitive detection methods which do not require amplification are also encompassed by the present genotyping methods. Methods well-known to those skilled in the art that can be used to detect biallelic polymorphisms include methods such as, conventional dot  
15 blot analyzes, single strand conformational polymorphism analysis (SSCP) described by Orita et al.(1989), denaturing gradient gel electrophoresis (DGGE), heteroduplex analysis, mismatch cleavage detection, and other conventional techniques as described in Sheffield et al.(1991), White et al.(1992), Grompe et al.(1989 and 1993). Another method for determining the identity of the nucleotide present at a particular polymorphic site employs a specialized exonuclease-resistant  
20 nucleotide derivative as described in US patent 4,656,127.

Preferred methods involve directly determining the identity of the nucleotide present at a biallelic marker site by sequencing assay, enzyme-based mismatch detection assay, or hybridization assay. The following is a description of some preferred methods. A highly preferred method is the microsequencing technique. The term "sequencing" is generally used herein to refer to polymerase  
25 extension of duplex primer/template complexes and includes both traditional sequencing and microsequencing.

#### **1) Sequencing Assays**

The nucleotide present at a polymorphic site can be determined by sequencing methods. In a preferred embodiment, DNA samples are subjected to PCR amplification before sequencing as  
30 described above. DNA sequencing methods are described in "Sequencing Of Amplified Genomic DNA And Identification Of Single Nucleotide Polymorphisms".

Preferably, the amplified DNA is subjected to automated dideoxy terminator sequencing reactions using a dye-primer cycle sequencing protocol. Sequence analysis allows the identification of the base present at the biallelic marker site.

#### **2) Microsequencing Assays**

In microsequencing methods, the nucleotide at a polymorphic site in a target DNA is detected by a single nucleotide primer extension reaction. This method involves appropriate microsequencing primers which, hybridize just upstream of the polymorphic base of interest in the target nucleic acid. A polymerase is used to specifically extend the 3' end of the primer with one single ddNTP (chain terminator) complementary to the nucleotide at the polymorphic site. Next the identity of the incorporated nucleotide is determined in any suitable way.

Typically, microsequencing reactions are carried out using fluorescent ddNTPs and the extended microsequencing primers are analyzed by electrophoresis on ABI 377 sequencing machines to determine the identity of the incorporated nucleotide as described in EP 412 883, the disclosure of which is incorporated herein by reference in its entirety. Alternatively capillary electrophoresis can be used in order to process a higher number of assays simultaneously. An example of a typical microsequencing procedure that can be used in the context of the present invention is provided in Example 4.

Different approaches can be used for the labeling and detection of ddNTPs. A homogeneous phase detection method based on fluorescence resonance energy transfer has been described by Chen and Kwok (1997) and Chen et al.(1997). In this method, amplified genomic DNA fragments containing polymorphic sites are incubated with a 5'-fluorescein-labeled primer in the presence of allelic dye-labeled dideoxyribonucleoside triphosphates and a modified Taq polymerase. The dye-labeled primer is extended one base by the dye-terminator specific for the allele present on the template. At the end of the genotyping reaction, the fluorescence intensities of the two dyes in the reaction mixture are analyzed directly without separation or purification. All these steps can be performed in the same tube and the fluorescence changes can be monitored in real time. Alternatively, the extended primer may be analyzed by MALDI-TOF Mass Spectrometry. The base at the polymorphic site is identified by the mass added onto the microsequencing primer (see Haff and Smirnov, 1997).

Microsequencing may be achieved by the established microsequencing method or by developments or derivatives thereof. Alternative methods include several solid-phase microsequencing techniques. The basic microsequencing protocol is the same as described previously, except that the method is conducted as a heterogeneous phase assay, in which the primer or the target molecule is immobilized or captured onto a solid support. To simplify the primer separation and the terminal nucleotide addition analysis, oligonucleotides are attached to solid supports or are modified in such ways that permit affinity separation as well as polymerase extension. The 5' ends and internal nucleotides of synthetic oligonucleotides can be modified in a number of different ways to permit different affinity separation approaches, e.g., biotinylation. If a single affinity group is used on the oligonucleotides, the oligonucleotides can be separated from the incorporated terminator reagent. This eliminates the need of physical or size separation. More than one oligonucleotide can be separated from the terminator reagent and analyzed simultaneously if

09762311.062501

more than one affinity group is used. This permits the analysis of several nucleic acid species or more nucleic acid sequence information per extension reaction. The affinity group need not be on the priming oligonucleotide but could alternatively be present on the template. For example, immobilization can be carried out via an interaction between biotinylated DNA and streptavidin-coated microtitration wells or avidin-coated polystyrene particles. In the same manner, oligonucleotides or templates may be attached to a solid support in a high-density format. In such solid phase microsequencing reactions, incorporated ddNTPs can be radiolabeled (Syvänen, 1994) or linked to fluorescein (Livak and Hainer, 1994). The detection of radiolabeled ddNTPs can be achieved through scintillation-based techniques. The detection of fluorescein-linked ddNTPs can be based on the binding of anti fluorescein antibody conjugated with alkaline phosphatase, followed by incubation with a chromogenic substrate (such as *p*-nitrophenyl phosphate). Other possible reporter-detection pairs include: ddNTP linked to dinitrophenyl (DNP) and anti-DNP alkaline phosphatase conjugate (Harju et al., 1993) or biotinylated ddNTP and horseradish peroxidase-conjugated streptavidin with *o*-phenylenediamine as a substrate (WO 92/15712). As yet another alternative solid-phase microsequencing procedure, Nyren et al.(1993) described a method relying on the detection of DNA polymerase activity by an enzymatic luminometric inorganic pyrophosphate detection assay (ELIDA).

Pastinen et al.(1997) describe a method for multiplex detection of single nucleotide polymorphism in which the solid phase minisequencing principle is applied to an oligonucleotide array format. High-density arrays of DNA probes attached to a solid support (DNA chips) are further described below.

In one aspect the present invention provides polynucleotides and methods to genotype one or more biallelic markers of the present invention by performing a microsequencing assay. Preferred microsequencing primers include the nucleotide sequences D1 to D15 and E1 to E15. It will be appreciated that the microsequencing primers listed in Example 5 are merely exemplary and that, any primer having a 3' end immediately adjacent to the polymorphic nucleotide may be used. Similarly, it will be appreciated that microsequencing analysis may be performed for any biallelic marker or any combination of biallelic markers of the present invention. One aspect of the present invention is a solid support which includes one or more microsequencing primers listed in Example 5, or fragments comprising at least 8, 12, 15, 20, 25, 30, 40, or 50 consecutive nucleotides thereof, to the extent that such lengths are consistent with the primer described, and having a 3' terminus immediately upstream of the corresponding biallelic marker, for determining the identity of a nucleotide at a biallelic marker site.

### 3) Mismatch detection assays based on polymerases and ligases

In one aspect the present invention provides polynucleotides and methods to determine the allele of one or more biallelic markers of the present invention in a biological sample, by mismatch

detection assays based on polymerases and/or ligases. These assays are based on the specificity of polymerases and ligases. Polymerization reactions places particularly stringent requirements on correct base pairing of the 3' end of the amplification primer and the joining of two oligonucleotides hybridized to a target DNA sequence is quite sensitive to mismatches close to the ligation site, especially at the 3' end. Methods, primers and various parameters to amplify DNA fragments comprising biallelic markers of the present invention are further described above in "Amplification Of DNA Fragments Comprising Biallelic Markers".

### Allele Specific Amplification Primers

Discrimination between the two alleles of a biallelic marker can also be achieved by allele specific amplification, a selective strategy, whereby one of the alleles is amplified without amplification of the other allele. For allele specific amplification, at least one member of the pair of primers is sufficiently complementary with a region of a *TBC-1* gene comprising the polymorphic base of a biallelic marker of the present invention to hybridize therewith and to initiate the amplification. Such primers are able to discriminate between the two alleles of a biallelic marker.

This is accomplished by placing the polymorphic base at the 3' end of one of the amplification primers. Because the extension forms from the 3' end of the primer, a mismatch at or near this position has an inhibitory effect on amplification. Therefore, under appropriate amplification conditions, these primers only direct amplification on their complementary allele. Determining the precise location of the mismatch and the corresponding assay conditions are well within the ordinary skill in the art.

### Ligation/Amplification Based Methods

The "Oligonucleotide Ligation Assay" (OLA) uses two oligonucleotides which are designed to be capable of hybridizing to abutting sequences of a single strand of a target molecules. One of the oligonucleotides is biotinylated, and the other is detectably labeled. If the precise complementary sequence is found in a target molecule, the oligonucleotides will hybridize such that their termini abut, and create a ligation substrate that can be captured and detected. OLA is capable of detecting single nucleotide polymorphisms and may be advantageously combined with PCR as described by Nickerson et al.(1990). In this method, PCR is used to achieve the exponential amplification of target DNA, which is then detected using OLA.

Other amplification methods which are particularly suited for the detection of single nucleotide polymorphism include LCR (ligase chain reaction), Gap LCR (GLCR) which are described above in "DNA Amplification". LCR uses two pairs of probes to exponentially amplify a specific target. The sequences of each pair of oligonucleotides, is selected to permit the pair to hybridize to abutting sequences of the same strand of the target. Such hybridization forms a substrate for a template-dependant ligase. In accordance with the present invention, LCR can be performed with oligonucleotides having the proximal and distal sequences of the same strand of a biallelic marker site. In one embodiment, either oligonucleotide will be designed to include the

biallelic marker site. In such an embodiment, the reaction conditions are selected such that the oligonucleotides can be ligated together only if the target molecule either contains or lacks the specific nucleotide that is complementary to the biallelic marker on the oligonucleotide. In an alternative embodiment, the oligonucleotides will not include the biallelic marker, such that when they hybridize to the target molecule, a "gap" is created as described in WO 90/01069. This gap is then "filled" with complementary dNTPs (as mediated by DNA polymerase), or by an additional pair of oligonucleotides. Thus at the end of each cycle, each single strand has a complement capable of serving as a target during the next cycle and exponential allele-specific amplification of the desired sequence is obtained.

- 10 Ligase/Polymcrase-mediated Genetic Bit Analysis<sup>TM</sup> is another method for determining the identity of a nucleotide at a preselected site in a nucleic acid molecule (WO 95/21271). This method involves the incorporation of a nucleoside triphosphate that is complementary to the nucleotide present at the preselected site onto the terminus of a primer molecule, and their subsequent ligation to a second oligonucleotide. The reaction is monitored by detecting a specific label attached to the reaction's solid phase or by detection in solution.

#### 4) Hybridization Assay Methods

A preferred method of determining the identity of the nucleotide present at a biallelic marker site involves nucleic acid hybridization. The hybridization probes, which can be conveniently used in such reactions, preferably include the probes defined herein. Any hybridization assay may be used including Southern hybridization, Northern hybridization, dot blot hybridization and solid-phase hybridization (see Sambrook et al., 1989).

Hybridization refers to the formation of a duplex structure by two single stranded nucleic acids due to complementary base pairing. Hybridization can occur between exactly complementary nucleic acid strands or between nucleic acid strands that contain minor regions of mismatch.

- 25 Specific probes can be designed that hybridize to one form of a biallelic marker and not to the other and therefore are able to discriminate between different allelic forms. Allele-specific probes are often used in pairs, one member of a pair showing perfect match to a target sequence containing the original allele and the other showing a perfect match to the target sequence containing the alternative allele. Hybridization conditions should be sufficiently stringent that there is a significant difference in hybridization intensity between alleles, and preferably an essentially binary response, whereby a probe hybridizes to only one of the alleles. Stringent, sequence specific hybridization conditions, under which a probe will hybridize only to the exactly complementary target sequence are well known in the art (Sambrook et al., 1989). Stringent conditions are sequence dependent and will be different in different circumstances. Generally, stringent conditions are selected to be about 30 5°C lower than the thermal melting point (T<sub>m</sub>) for the specific sequence at a defined ionic strength and pH. Although such hybridization can be performed in solution, it is preferred to employ a

solid-phase hybridization assay. The target DNA comprising a biallelic marker of the present invention may be amplified prior to the hybridization reaction. The presence of a specific allele in the sample is determined by detecting the presence or the absence of stable hybrid duplexes formed between the probe and the target DNA. The detection of hybrid duplexes can be carried out by a number of methods. Various detection assay formats are well known which utilize detectable labels bound to either the target or the probe to enable detection of the hybrid duplexes. Typically, hybridization duplexes are separated from unhybridized nucleic acids and the labels bound to the duplexes are then detected. Those skilled in the art will recognize that wash steps may be employed to wash away excess target DNA or probe as well as unbound conjugate. Further, standard heterogeneous assay formats are suitable for detecting the hybrids using the labels present on the primers and probes.

Two recently developed assays allow hybridization-based allele discrimination with no need for separations or washes (see Landegren U. et al., 1998). The TaqMan assay takes advantage of the 5' nuclease activity of Taq DNA polymerase to digest a DNA probe annealed specifically to the accumulating amplification product. TaqMan probes are labeled with a donor-acceptor dye pair that interacts via fluorescence energy transfer. Cleavage of the TaqMan probe by the advancing polymerase during amplification dissociates the donor dye from the quenching acceptor dye, greatly increasing the donor fluorescence. All reagents necessary to detect two allelic variants can be assembled at the beginning of the reaction and the results are monitored in real time (see Livak et al., 1995). In an alternative homogeneous hybridization based procedure, molecular beacons are used for allele discriminations. Molecular beacons are hairpin-shaped oligonucleotide probes that report the presence of specific nucleic acids in homogeneous solutions. When they bind to their targets they undergo a conformational reorganization that restores the fluorescence of an internally quenched fluorophore (Tyagi et al., 1998).

The polynucleotides provided herein can be used to produce probes which can be used in hybridization assays for the detection of biallelic marker alleles in biological samples. These probes are characterized in that they preferably comprise between 8 and 50 nucleotides, and in that they are sufficiently complementary to a sequence comprising a biallelic marker of the present invention to hybridize thereto and preferably sufficiently specific to be able to discriminate the targeted sequence for only one nucleotide variation. A particularly preferred probe is 25 nucleotides in length. Preferably the biallelic marker is within 4 nucleotides of the center of the polynucleotide probe. In particularly preferred probes, the biallelic marker is at the center of said polynucleotide. Preferred probes comprise a nucleotide sequence selected from the group consisting of amplicons listed in Table 1 and the sequences complementary thereto, or a fragment thereof, said fragment comprising at least about 8 consecutive nucleotides, preferably 10, 15, 20, more preferably 25, 30, 40, 47, or 50 consecutive nucleotides and containing a polymorphic base. Preferred probes comprise a nucleotide sequence selected from the group consisting of P1 to P7, P9 to P13, P15 to

P19 and the sequences complementary thereto. In preferred embodiments the polymorphic base(s) are within 5, 4, 3, 2, 1, nucleotides of the center of the said polynucleotide, more preferably at the center of said polynucleotide.

Preferably the probes of the present invention are labeled or immobilized on a solid support.

- 5 Labels and solid supports are further described in "Oligonucleotide Probes and Primers". The probes can be non-extendable as described in "Oligonucleotide Probes and Primers".

By assaying the hybridization to an allele specific probe, one can detect the presence or absence of a biallelic marker allele in a given sample. High-Throughput parallel hybridization in array format is specifically encompassed within "hybridization assays" and are described below.

10 **5) Hybridization To Addressable Arrays Of Oligonucleotides**

- Hybridization assays based on oligonucleotide arrays rely on the differences in hybridization stability of short oligonucleotides to perfectly matched and mismatched target sequence variants. Efficient access to polymorphism information is obtained through a basic structure comprising high-density arrays of oligonucleotide probes attached to a solid support (e.g.,  
15 the chip) at selected positions. Each DNA chip can contain thousands to millions of individual synthetic DNA probes arranged in a grid-like pattern and miniaturized to the size of a dime.

- The chip technology has already been applied with success in numerous cases. For example, the screening of mutations has been undertaken in the BRCA1 gene, in *S. cerevisiae* mutant strains, and in the protease gene of HIV-1 virus (Hacia et al., 1996; Shoemaker et al., 1996;  
20 Kozal et al., 1996). Chips of various formats for use in detecting biallelic polymorphisms can be produced on a customized basis by Affymetrix (GeneChip™), Hyseq (HyChip and HyGnostics), and Protogene Laboratories.

- In general, these methods employ arrays of oligonucleotide probes that are complementary to target nucleic acid sequence segments from an individual which, target sequences include a  
25 polymorphic marker. EP 785280 describes a tiling strategy for the detection of single nucleotide polymorphisms. Briefly, arrays may generally be "tiled" for a large number of specific polymorphisms. By "tiling" is generally meant the synthesis of a defined set of oligonucleotide probes which is made up of a sequence complementary to the target sequence of interest, as well as preselected variations of that sequence, e.g., substitution of one or more given positions with one or  
30 more members of the basis set of nucleotides. Tiling strategies are further described in PCT application No. WO 95/11995. In a particular aspect, arrays are tiled for a number of specific, identified biallelic marker sequences. In particular, the array is tiled to include a number of detection blocks, each detection block being specific for a specific biallelic marker or a set of biallelic markers. For example, a detection block may be tiled to include a number of probes, which  
35 span the sequence segment that includes a specific polymorphism. To ensure probes that are complementary to each allele, the probes are synthesized in pairs differing at the biallelic marker.

09:62:11.062501  
T05290.T229.60



In addition to the probes differing at the polymorphic base, monosubstituted probes are also generally tiled within the detection block. These monosubstituted probes have bases at and up to a certain number of bases in either direction from the polymorphism, substituted with the remaining nucleotides (selected from A, T, G, C and U). Typically the probes in a tiled detection block will include substitutions of the sequence positions up to and including those that are 5 bases away from the biallelic marker. The monosubstituted probes provide internal controls for the tiled array, to distinguish actual hybridization from artefactual cross-hybridization. Upon completion of hybridization with the target sequence and washing of the array, the array is scanned to determine the position on the array to which the target sequence hybridizes. The hybridization data from the scanned array is then analyzed to identify which allele or alleles of the biallelic marker are present in the sample. Hybridization and scanning may be carried out as described in PCT application No. WO 92/10092 and WO 95/11995 and US patent No. 5,424,186.

Thus, in some embodiments, the chips may comprise an array of nucleic acid sequences of fragments of about 15 nucleotides in length. In further embodiments, the chip may comprise an array including at least one of the sequences selected from the group consisting of amplicons listed in table 1 and the sequences complementary thereto, or a fragment thereof, said fragment comprising at least about 8 consecutive nucleotides, preferably 10, 15, 20, more preferably 25, 30, 40, 47, or 50 consecutive nucleotides and containing a polymorphic base. In preferred embodiments the polymorphic base is within 5, 4, 3, 2, 1, nucleotides of the center of the said polynucleotide, more preferably at the center of said polynucleotide. In some embodiments, the chip may comprise an array of at least 2, 3, 4, 5, 6, 7, 8 or more of these polynucleotides of the invention. Solid supports and polynucleotides of the present invention attached to solid supports are further described in "Oligonucleotide Probes And Primers".

## 6) Integrated Systems

Another technique, which may be used to analyze polymorphisms, includes multicomponent integrated systems, which miniaturize and compartmentalize processes such as PCR and capillary electrophoresis reactions in a single functional device. An example of such technique is disclosed in US patent 5,589,136, which describes the integration of PCR amplification and capillary electrophoresis in chips.

Integrated systems can be envisaged mainly when microfluidic systems are used. These systems comprise a pattern of microchannels designed onto a glass, silicon, quartz, or plastic wafer included on a microchip. The movements of the samples are controlled by electric, electroosmotic or hydrostatic forces applied across different areas of the microchip to create functional microscopic valves and pumps with no moving parts.

For genotyping biallelic markers, the microfluidic system may integrate nucleic acid amplification, microsequencing, capillary electrophoresis and a detection method such as laser-induced fluorescence detection.

#### Association Studies With The Biallelic Markers Of The *TBC-1* Gene

5 The identification of genes involved in suspected heterogeneous, polygenic and multifactorial traits such as cancer can be carried out through two main strategies currently used for genetic mapping: linkage analysis and association studies. Association studies examine the frequency of marker alleles in unrelated trait positive (T+) individuals compared with trait negative (T-) controls, and are generally employed in the detection of polygenic inheritance. Association  
10 studies as a method of mapping genetic traits rely on the phenomenon of linkage disequilibrium.

If two genetic loci lie on the same chromosome, then sets of alleles of these loci on the same chromosomal segment (called haplotypes) tend to be transmitted as a block from generation to generation. When not broken up by recombination, haplotypes can be tracked not only through pedigrees but also through populations. The resulting phenomenon at the population level is that the  
15 occurrence of pairs of specific alleles at different loci on the same chromosome is not random, and the deviation from random is called linkage disequilibrium (LD).

If a specific allele in a given gene is directly involved in causing a particular trait T, its frequency will be statistically increased in a trait positive population when compared to the frequency in a trait negative population. As a consequence of the existence of linkage  
20 disequilibrium, the frequency of all other alleles present in the haplotype carrying the trait-causing allele (TCA) will also be increased in trait positive individuals compared to trait negative individuals. Therefore, association between the trait and any allele in linkage disequilibrium with the trait-causing allele will suffice to suggest the presence of a trait-related gene in that particular allele's region. Linkage disequilibrium allows the relative frequencies in trait positive and trait  
25 negative populations of a limited number of genetic polymorphisms (specifically biallelic markers) to be analyzed as an alternative to screening all possible functional polymorphisms in order to find trait-causing alleles.

The general strategy to perform association studies using biallelic markers derived from a candidate region is to scan two groups of individuals (trait positive and trait negative control  
30 individuals which are characterized by a well defined phenotype as described below) in order to measure and statistically compare the allele frequencies of such biallelic markers in both groups.

If a statistically significant association with a trait is identified for at least one or more of the analyzed biallelic markers, one can assume that : either the associated allele is directly responsible for causing the trait (associated allele is the trait-causing allele), or the associated allele  
35 is in linkage disequilibrium with the trait-causing allele. If the evidence indicates that the associated allele within the candidate region is most probably not the trait-causing allele but is in linkage

0976234 06504

**Collection of DNA samples from trait positive (trait +) and trait negative (trait – individuals (inclusion criteria)**

Nevertheless, even in the absence of such a bimodal distribution (as may in fact be the case for more complex genetic traits), any genetic trait may still be analyzed by the association method proposed here by carefully selecting the individuals to be included in the trait positive and trait negative phenotypic groups. The selection procedure involves to select individuals at opposite ends of the non-bimodal phenotype spectra of the trait under study, so as to include in these trait positive and trait negative populations individuals which clearly represent extreme, preferably non-overlapping phenotypes.

The definition of the inclusion criteria for the trait positive and trait negative populations is an important aspect of the present invention. The selection of drastically different but relatively uniform phenotypes enables efficient comparisons in association studies and the possible detection of marked differences at the genetic level, provided that the sample sizes of the populations under study are significant enough.

A first group of between 50 and 300 trait positive individuals, preferably about 100  
25 individuals, can be recruited according to clinical inclusion criteria.

In each case, a similar number of trait negative individuals, preferably more than 100 individuals, are included in such studies who are preferably both ethnically- and age-matched to the trait positive cases. They are checked for the absence of the clinical criteria defined above. Both trait positive and trait negative individuals should correspond to unrelated cases.

### 30 Genotyping of trait positive and trait negative individuals

Allelic frequencies of the biallelic markers in each of the above described population can be determined using one of the methods described above under the heading "Methods of Genotyping DNA samples for biallelic markers". Analyses are preferably performed on amplified fragments obtained by genomic PCR performed on the DNA samples from each individual in similar conditions as those described above for the generation of biallelic markers.

In a preferred embodiment, amplified DNA samples are subjected to automated microsequencing reactions using fluorescent ddNTPs (specific fluorescence for each ddNTP) and

the appropriate microsequencing oligonucleotides which hybridize just upstream of the polymorphic base.

Genotyping is further described in Example 5.

Associations studies can be carried out by the skilled technician using the biallelic markers of the invention defined above, with different trait positive and trait negative populations. Suitable examples of association studies using biallelic markers of the *TBC-1* gene, including the biallelic markers A1 to A19, involve studies on the following populations:

- a trait positive population suffering from a cancer, preferably prostate cancer and a healthy unaffected population; or

- 10 - a trait positive population suffering from prostate cancer treated with agents acting against prostate cancer and suffering from side-effects resulting from this treatment and an trait negative population suffering from prostate cancer treated with same agents without any substantial side-effects, or

- 15 - a trait positive population suffering from prostate cancer treated with agents acting against prostate cancer showing a beneficial response and a trait negative population suffering from prostate cancer treated with same agents without any beneficial response, or

- a trait positive population suffering from prostate cancer presenting highly aggressive prostate cancer tumors and a trait negative population suffering from prostate cancer with prostate cancer tumors devoid of aggressiveness.

20 It is another object of the present invention to provide a method for the identification and characterization of an association between an allele of one or more biallelic markers of a *TBC-1* gene and a trait. The method comprises the steps of :

- genotyping a marker or a group of biallelic markers according to the invention in trait positive;

- 25 - genotyping a marker or a group of biallelic markers according to the invention in and trait negative individuals; and

- establishing a statistically significant association between one allele of at least one marker and the trait.

Preferably, the trait positive and trait negative individuals are selected from non-  
30 overlapping phenotypes as regards to the trait under study. In one embodiment, the biallelic marker are selected from the group consisting of the biallelic markers A1 to A19.

In a preferred embodiment, the trait is cancer, prostate cancer, an early onset of prostate cancer, a susceptibility to prostate cancer, the level of aggressiveness of prostate cancer tumors, a modified expression of the *TBC-1* gene, a modified production of the TBC-1 protein, or the  
35 production of a modified TBC-1 protein.

In a further embodiment, the trait negative population can be replaced in the association studies by a random control population.

0976231-05201  
T05290-TE29260

The step of testing for and detecting the presence of DNA comprising specific alleles of a biallelic marker or a group of biallelic markers of the present invention can be carried out as described further below.

### Oligonucleotide Probes And Primers

5 The invention relates also to oligonucleotide molecules useful as probes or primers, wherein said oligonucleotide molecules hybridize specifically with a nucleotide sequence comprised in the *TBC-1* gene, particularly the *TBC-1* genomic sequence of SEQ ID Nos 1 and 2 or the *TBC-1* cDNAs sequences of SEQ ID Nos 3 and 4. More particularly, the present invention also concerns oligonucleotides for the detection of alleles of biallelic markers of the *TBC-1* gene. These  
10 oligonucleotides are useful either as primers for use in various processes such as DNA amplification and microsequencing or as probes for DNA recognition in hybridization analyses. Polynucleotides derived from the *TBC-1* gene are useful in order to detect the presence of at least a copy of a nucleotide sequence of SEQ ID Nos 1-4, or a fragment, complement, or variant thereof in a test sample.

15 Particularly preferred probes and primers of the invention include isolated, purified, or recombinant polynucleotides comprising a contiguous span of at least 12, 15, 18, 20, 25, 30, 35, 40, 50, 60, 70, 80, 90, 100, 150, 200, 500, or 1000 nucleotides of a nucleotide sequence selected from the group consisting of SEQ ID Nos 1 and 2, or the complements thereof. Additionally preferred probes and primers of the invention include isolated, purified, or recombinant polynucleotides  
20 comprising a contiguous span of at least 12, 15, 18, 20, 25, 30, 35, 40, 50, 60, 70, 80, 90, 100, 150, 200, 500, or 1000 nucleotides of SEQ ID No 1 or the complements thereof, wherein said contiguous span comprises at least 1, 2, 3, 5, or 10 of the following nucleotide positions of SEQ ID No 1: 1-1000, 1001-2000, 2001-3000, 3001-4000, 4001-5000, 5001-6000, 6001-7000, 7001-8000, 8001-9000, 9001-10000, 10001-11000, 11001-12000, 12001-13000, 13001-14000, 14001-15000,  
25 15001-16000, 16001-17000, and 17001-17590. Other preferred probes and primers of the invention include isolated, purified, or recombinant polynucleotides comprising a contiguous span of at least 12, 15, 18, 20, 25, 30, 35, 40, 50, 60, 70, 80, 90, 100, 150, 200, 500, or 1000 nucleotides of SEQ ID No 2 or the complements thereof, wherein said contiguous span comprises at least 1, 2, 3, 5, or 10 of the following nucleotide positions of SEQ ID No 2: 1-5000, 5001-10000, 10001-15000, 15001-  
30 20000, 20001-25000, 25001-30000, 30001-35000, 35001-40000, 40001-45000, 45001-50000, 50001-55000, 55001-60000, 60001-65000, 65001-70000, 70001-75000, 75001-80000, 80001-85000, 85001-90000, 90001-95000, and 95001-99960.

Moreover, preferred probes and primers of the invention include isolated, purified, or recombinant polynucleotides comprising a contiguous span of at least 12, 15, 18, 20, 25, 30, 35, 40,  
35 50, 60, 70, 80, 90, 100, 150, 200, 500, or 1000 nucleotides of a nucleotide sequence selected from the group consisting of SEQ ID Nos 3 and 4, or the complements thereof.. Particularly preferred

probes and primers of the invention include isolated, purified, or recombinant polynucleotides comprising a contiguous span of at least 12, 15, 18, 20, 25, 30, 35, 40, 50, 60, 70, 80, 90, 100, 150, 200, 500, or 1000 nucleotides of SEQ ID No 3 or the complements thereof, wherein said contiguous span comprises at least 1, 2, 3, 5, or 10 of the following nucleotide positions of SEQ ID  
5 No 3: 1-500, 501-1000, 1001-1500, 1501-2000, 2001-2500, 2501-3000, 3001-3500, and 3501-3983. Additional preferred probes and primers of the invention include isolated, purified, or recombinant polynucleotides comprising a contiguous span of at least 12, 15, 18, 20, 25, 30, 35, 40, 50, 60, 70, 80, 90, 100, 150, 200, 500, or 1000 nucleotides of SEQ ID No 4 or the complements thereof, wherein said contiguous span comprises at least 1, 2, 3, 5, or 10 of the following nucleotide  
10 positions of SEQ ID No 4: 1-500, 501-1000, 1001-1500, 1501-2000, 2001-2500, 2501-3000, 3001-3500, and 3501-3988.

Thus, the invention also relates to nucleic acid probes characterized in that they hybridize specifically, under the stringent hybridization conditions defined above, with a nucleic acid selected from the group consisting of the nucleotide sequences of SEQ ID Nos 1-4 or a variant thereof or a  
15 sequence complementary thereto.

In one embodiment the invention encompasses isolated, purified, and recombinant polynucleotides consisting of, or consisting essentially of a contiguous span of 8 to 50 nucleotides of any one of SEQ ID Nos 1 and 2 and the complement thereof, wherein said span includes a *TBC-I*-related biallelic marker in said sequence; optionally, wherein said *TBC-I*-related biallelic marker  
20 is selected from the group consisting of A1 to A19, and the complements thereof, or optionally the biallelic markers in linkage disequilibrium therewith; optionally, wherein said contiguous span is 18 to 35 nucleotides in length and said biallelic marker is within 4 nucleotides of the center of said polynucleotide; optionally, wherein said polynucleotide consists of said contiguous span and said contiguous span is 25 nucleotides in length and said biallelic marker is at the center of said  
25 polynucleotide; optionally, wherein the 3' end of said contiguous span is present at the 3' end of said polynucleotide; and optionally, wherein the 3' end of said contiguous span is located at the 3' end of said polynucleotide and said biallelic marker is present at the 3' end of said polynucleotide. In a preferred embodiment, said probes comprises, consists of, or consists essentially of a sequence selected from the following sequences: P1 to P7, P9 to P13, P15 to P19 and the complementary  
30 sequences thereto.

In another embodiment the invention encompasses isolated, purified and recombinant polynucleotides comprising, consisting of, or consisting essentially of a contiguous span of 8 to 50 nucleotides of SEQ ID Nos 1 and 2, or the complements thereof, wherein the 3' end of said contiguous span is located at the 3' end of said polynucleotide, and wherein the 3' end of said  
35 polynucleotide is located within 20 nucleotides upstream of a *TBC-I*-related biallelic marker in said sequence; optionally, wherein said *TBC-I*-related biallelic marker is selected from the group consisting of A1 to A19, and the complements thereof, or optionally the biallelic markers in linkage

T05290: T05290

disequilibrium therewith; optionally, wherein the 3' end of said polynucleotide is located 1 nucleotide upstream of said *TBC-1*-related biallelic marker in said sequence; and optionally, wherein said polynucleotide consists essentially of a sequence selected from the following sequences: D1 to D19 and E1 to E19.

5 In a further embodiment, the invention encompasses isolated, purified, or recombinant polynucleotides comprising, consisting of, or consisting essentially of a sequence selected from the following sequences: B1 to B15 and C1 to C15.

In an additional embodiment, the invention encompasses polynucleotides for use in hybridization assays, sequencing assays, and enzyme-based mismatch detection assays for  
10 determining the identity of the nucleotide at a *TBC-1*-related biallelic marker in SEQ ID Nos 1 and 2, or the complements thereof, as well as polynucleotides for use in amplifying segments of nucleotides comprising a *TBC-1*-related biallelic marker in SEQ ID Nos 1 and 2, or the complements thereof; optionally, wherein said *TBC-1*-related biallelic marker is selected from the group consisting of A1 to A19, and the complements thereof, or optionally the biallelic markers in  
15 linkage disequilibrium therewith.

A probe or a primer according to the invention has between 8 and 1000 nucleotides in length, or is specified to be at least 12, 15, 18, 20, 25, 35, 40, 50, 60, 70, 80, 100, 250, 500 or 1000 nucleotides in length. More particularly, the length of these probes and primers can range from 8, 10, 15, 20, or 30 to 100 nucleotides, preferably from 10 to 50, more preferably from 15 to 30  
20 nucleotides. Shorter probes and primers tend to lack specificity for a target nucleic acid sequence and generally require cooler temperatures to form sufficiently stable hybrid complexes with the template. Longer probes and primers are expensive to produce and can sometimes self-hybridize to form hairpin structures. The appropriate length for primers and probes under a particular set of assay conditions may be empirically determined by one of skill in the art. A preferred probe or  
25 primer consists of a nucleic acid comprising a polynucleotide selected from the group of the nucleotide sequences of P1 to P7, P9 to P13, P15 to P19 and the complementary sequence thereto, B1 to B15, C1 to C15, D1 to D19, E1 to E19, for which the respective locations in the sequence listing are provided in Tables 2, 3 and 4.

The formation of stable hybrids depends on the melting temperature ( $T_m$ ) of the DNA. The  
30  $T_m$  depends on the length of the primer or probe, the ionic strength of the solution and the G+C content. The higher the G+C content of the primer or probe, the higher is the melting temperature because G:C pairs are held by three H bonds whereas A:T pairs have only two. The GC content in the probes of the invention usually ranges between 10 and 75 %, preferably between 35 and 60 %, and more preferably between 40 and 55 %.

35 The primers and probes can be prepared by any suitable method, including, for example, cloning and restriction of appropriate sequences and direct chemical synthesis by a method such as the phosphodiester method of Narang et al.(1979), the phosphodiester method of Brown et

09624-00001

al.(1979), the diethylphosphoramidite method of Beaucage et al.(1981) and the solid support method described in EP 0 707 592.

Detection probes are generally nucleic acid sequences or uncharged nucleic acid analogs such as, for example peptide nucleic acids which are disclosed in International Patent Application WO 92/20702, morpholino analogs which are described in U.S. Patents Numbered 5,185,444; 5,034,506 and 5,142,047. The probe may have to be rendered "non-extendable" in that additional dNTPs cannot be added to the probe. In and of themselves analogs usually are non-extendable and nucleic acid probes can be rendered non-extendable by modifying the 3' end of the probe such that the hydroxyl group is no longer capable of participating in elongation. For example, the 3' end of the probe can be functionalized with the capture or detection label to thereby consume or otherwise block the hydroxyl group. Alternatively, the 3' hydroxyl group simply can be cleaved, replaced or modified, U.S. Patent Application Serial No. 07/049,061 filed April 19, 1993 describes modifications, which can be used to render a probe non-extendable.

Any of the polynucleotides of the present invention can be labeled, if desired, by incorporating any label known in the art to be detectable by spectroscopic, photochemical, biochemical, immunochemical, or chemical means. For example, useful labels include radioactive substances (including,  $^{32}\text{P}$ ,  $^{35}\text{S}$ ,  $^3\text{H}$ ,  $^{125}\text{I}$ ), fluorescent dyes (including, 5-bromodesoxyuridin, fluorescein, acetylaminofluorene, digoxigenin) or biotin. Preferably, polynucleotides are labeled at their 3' and 5' ends. Examples of non-radioactive labeling of nucleic acid fragments are described in the French patent No. FR-7810975 or by Urdea et al (1988) or Sanchez-Pescador et al (1988). In addition, the probes according to the present invention may have structural characteristics such that they allow the signal amplification, such structural characteristics being, for example, branched DNA probes as those described by Urdea et al. in 1991 or in the European patent No. EP 0 225 807 (Chiron).

A label can also be used to capture the primer, so as to facilitate the immobilization of either the primer or a primer extension product, such as amplified DNA, on a solid support. A capture label is attached to the primers or probes and can be a specific binding member which forms a binding pair with the solid's phase reagent's specific binding member (e.g. biotin and streptavidin). Therefore depending upon the type of label carried by a polynucleotide or a probe, it may be employed to capture or to detect the target DNA. Further, it will be understood that the polynucleotides, primers or probes provided herein, may, themselves, serve as the capture label. For example, in the case where a solid phase reagent's binding member is a nucleic acid sequence, it may be selected such that it binds a complementary portion of a primer or probe to thereby immobilize the primer or probe to the solid phase. In cases where a polynucleotide probe itself serves as the binding member, those skilled in the art will recognize that the probe will contain a sequence or "tail" that is not complementary to the target. In the case where a polynucleotide



primer itself serves as the capture label, at least a portion of the primer will be free to hybridize with a nucleic acid on a solid phase. DNA Labeling techniques are well known to the skilled technician.

The probes of the present invention are useful for a number of purposes. They can be notably used in Southern hybridization to genomic DNA. The probes can also be used to detect

5 PCR amplification products. They may also be used to detect mismatches in the *TBC-1* gene or mRNA using other techniques.

Any of the polynucleotides, primers and probes of the present invention can be conveniently immobilized on a solid support. Solid supports are known to those skilled in the art and include the walls of wells of a reaction tray, test tubes, polystyrene beads, magnetic beads,

10 nitrocellulose strips, membranes, microparticles such as latex particles, sheep (or other animal) red blood cells, duracytes and others. The solid support is not critical and can be selected by one skilled in the art. Thus, latex particles, microparticles, magnetic or non-magnetic beads, membranes, plastic tubes, walls of microtiter wells, glass or silicon chips, sheep (or other suitable animal's) red blood cells and duracytes are all suitable examples. Suitable methods for immobilizing nucleic

15 acids on solid phases include ionic, hydrophobic, covalent interactions and the like. A solid support, as used herein, refers to any material which is insoluble, or can be made insoluble by a subsequent reaction. The solid support can be chosen for its intrinsic ability to attract and immobilize the capture reagent. Alternatively, the solid phase can retain an additional receptor which has the ability to attract and immobilize the capture reagent. The additional receptor can

20 include a charged substance that is oppositely charged with respect to the capture reagent itself or to a charged substance conjugated to the capture reagent. As yet another alternative, the receptor molecule can be any specific binding member which is immobilized upon (attached to) the solid support and which has the ability to immobilize the capture reagent through a specific binding reaction. The receptor molecule enables the indirect binding of the capture reagent to a solid

25 support material before the performance of the assay or during the performance of the assay. The solid phase thus can be a plastic, derivatized plastic, magnetic or non-magnetic metal, glass or silicon surface of a test tube, microtiter well, sheet, bead, microparticle, chip, sheep (or other suitable animal's) red blood cells, duracytes® and other configurations known to those of ordinary skill in the art. The polynucleotides of the invention can be attached to or immobilized on a solid

30 support individually or in groups of at least 2, 5, 8, 10, 12, 15, 20, or 25 distinct polynucleotides of the invention to a single solid support. In addition, polynucleotides other than those of the invention may be attached to the same solid support as one or more polynucleotides of the invention.

Consequently, the invention also deals with a method for detecting the presence of a nucleic

35 acid comprising a nucleotide sequence selected from a group consisting of SEQ ID Nos 1-4, a fragment or a variant thereof and a complementary sequence thereto in a sample, said method comprising the following steps of:

097634.050  
T0320:TE29250

a) bringing into contact a nucleic acid probe or a plurality of nucleic acid probes which can hybridize with a nucleotide sequence included in a nucleic acid selected from the group consisting of the nucleotide sequences of SEQ ID Nos 1-4, a fragment or a variant thereof and a complementary sequence thereto and the sample to be assayed; and

5 b) detecting the hybrid complex formed between the probe and a nucleic acid in the sample.

The invention further concerns a kit for detecting the presence of a nucleic acid comprising a nucleotide sequence selected from a group consisting of SEQ ID Nos 1-4, a fragment or a variant thereof and a complementary sequence thereto in a sample, said kit comprising:

a) a nucleic acid probe or a plurality of nucleic acid probes which can hybridize with a  
10 nucleotide sequence included in a nucleic acid selected from the group consisting of the nucleotide sequences of SEQ ID Nos 1-4, a fragment or a variant thereof and a complementary sequence thereto; and

b) optionally, the reagents necessary for performing the hybridization reaction.

In a first preferred embodiment of this detection method and kit, said nucleic acid probe or  
15 the plurality of nucleic acid probes are labeled with a detectable molecule. In a second preferred embodiment of said method and kit, said nucleic acid probe or the plurality of nucleic acid probes has been immobilized on a substrate. In a third preferred embodiment, the nucleic acid probe or the plurality of nucleic acid probes comprise either a sequence which is selected from the group consisting of the nucleotide sequences of P1 to P7, P9 to P13, P15 to P19 and the complementary  
20 sequence thereto, B1 to B15, C1 to C15, D1 to D19, E1 to E19 or a biallelic marker selected from the group consisting of A1 to A19 and the complements thereto.

### Oligonucleotide Arrays

A substrate comprising a plurality of oligonucleotide primers or probes of the invention may be used either for detecting or amplifying targeted sequences in the *TBC-1* gene and may also  
25 be used for detecting mutations in the coding or in the non-coding sequences of the *TBC-1* gene.

Any polynucleotide provided herein may be attached in overlapping areas or at random locations on the solid support. Alternatively the polynucleotides of the invention may be attached in an ordered array wherein each polynucleotide is attached to a distinct region of the solid support which does not overlap with the attachment site of any other polynucleotide. Preferably, such an  
30 ordered array of polynucleotides is designed to be "addressable" where the distinct locations are recorded and can be accessed as part of an assay procedure. Addressable polynucleotide arrays typically comprise a plurality of different oligonucleotide probes that are coupled to a surface of a substrate in different known locations. The knowledge of the precise location of each polynucleotides location makes these "addressable" arrays particularly useful in hybridization  
35 assays. Any addressable array technology known in the art can be employed with the polynucleotides of the invention. One particular embodiment of these polynucleotide arrays is known as the Genechips™, and has been generally described in US Patent 5,143,854; PCT

09634-03501  
T0530-TE2260

publications WO 90/15070 and 92/10092. These arrays may generally be produced using mechanical synthesis methods or light directed synthesis methods which incorporate a combination of photolithographic methods and solid phase oligonucleotide synthesis (Fodor et al., 1991). The immobilization of arrays of oligonucleotides on solid supports has been rendered possible by the development of a technology generally identified as "Very Large Scale Immobilized Polymer Synthesis" (VLSIPS™) in which, typically, probes are immobilized in a high density array on a solid surface of a chip. Examples of VLSIPS™ technologies are provided in US Patents 5,143,854; and 5,412,087 and in PCT Publications WO 90/15070, WO 92/10092 and WO 95/11995, which describe methods for forming oligonucleotide arrays through techniques such as light-directed synthesis techniques. In designing strategies aimed at providing arrays of nucleotides immobilized on solid supports, further presentation strategies were developed to order and display the oligonucleotide arrays on the chips in an attempt to maximize hybridization patterns and sequence information. Examples of such presentation strategies are disclosed in PCT Publications WO 94/12305, WO 94/11530, WO 97/29212 and WO 97/31256.

In another embodiment of the oligonucleotide arrays of the invention, an oligonucleotide probe matrix may advantageously be used to detect mutations occurring in the *TBC-1* gene and preferably in its regulatory region. For this particular purpose, probes are specifically designed to have a nucleotide sequence allowing their hybridization to the genes that carry known mutations (either by deletion, insertion or substitution of one or several nucleotides). By known mutations, it is meant, mutations on the *TBC-1* gene that have been identified according, for example to the technique used by Huang et al.(1996) or Samson et al.(1996).

Another technique that is used to detect mutations in the *TBC-1* gene is the use of a high-density DNA array. Each oligonucleotide probe constituting a unit element of the high density DNA array is designed to match a specific subsequence of the *TBC-1* genomic DNA or cDNA.

Thus, an array consisting of oligonucleotides complementary to subsequences of the target gene sequence is used to determine the identity of the target sequence with the wild gene sequence, measure its amount, and detect differences between the target sequence and the reference wild gene sequence of the *TBC-1* gene. In one such design, termed 4L tiled array, is implemented a set of four probes (A, C, G, T), preferably 15-nucleotide oligomers. In each set of four probes, the perfect complement will hybridize more strongly than mismatched probes. Consequently, a nucleic acid target of length L is scanned for mutations with a tiled array containing 4L probes, the whole probe set containing all the possible mutations in the known wild reference sequence. The hybridization signals of the 15-mer probe set tiled array are perturbed by a single base change in the target sequence. As a consequence, there is a characteristic loss of signal or a "footprint" for the probes flanking a mutation position. This technique was described by Chee et al. in 1996.

Consequently, the invention concerns an array of nucleic acid molecules comprising at least one polynucleotide described above as probes and primers. Preferably, the invention concerns an

array of nucleic acid comprising at least two polynucleotides described above as probes and primers.

A further object of the invention consists of an array of nucleic acid sequences comprising either at least one of the sequences selected from the group consisting of P1 to P7, P9 to P13, P15 to P19, B1 to B15, C1 to C15, D1 to D19, E1 to E19, the sequences complementary thereto, a fragment thereof of at least 8, 10, 12, 15, 18, 20, 25, 30, or 40 consecutive nucleotides thereof, and at least one sequence comprising a biallelic marker selected from the group consisting of A1 to A19 and the complements thereto.

The invention also pertains to an array of nucleic acid sequences comprising either at least two of the sequences selected from the group consisting of P1 to P7, P9 to P13, P15 to P19, B1 to B15, C1 to C15, D1 to D19, E1 to E19, the sequences complementary thereto, a fragment thereof of at least 8 consecutive nucleotides thereof, and at least two sequences comprising a biallelic marker selected from the group consisting of A1 to A19 and the complements thereof.

#### Vectors For The Expression Of A Regulatory Or A Coding Polynucleotide Of *TBC-1*.

Any of the regulatory polynucleotides or the coding polynucleotides of the invention may be inserted into recombinant vectors for expression in a recombinant host cell or a recombinant host organism.

Thus, the present invention also encompasses a family of recombinant vectors that contains either a regulatory polynucleotide selected from the group consisting of any one of the regulatory polynucleotides derived from the *TBC-1* genomic sequences of SEQ ID Nos 1 and 2, or a polynucleotide comprising the *TBC-1* coding sequence, or both.

In a first preferred embodiment, a recombinant vector of the invention is used as an expression vector : (a) the *TBC-1* regulatory sequence comprised therein drives the expression of a coding polynucleotide operably linked thereto; (b) the *TBC-1* coding sequence is operably linked to regulation sequences allowing its expression in a suitable cell host and/or host organism.

In a second preferred embodiment, a recombinant vector of the invention is used to amplify the inserted polynucleotide derived from the *TBC-1* genomic sequences of SEQ ID Nos 1 and 2 or *TBC-1* cDNAs in a suitable cell host , this polynucleotide being amplified at every time that the recombinant vector replicates.

More particularly, the present invention relates to expression vectors which include nucleic acids encoding a TBC-1 protein, preferably the TBC-1 protein of the amino acid sequence of SEQ ID No 5 described therein, under the control of a regulatory sequence selected among the *TBC-1* regulatory polynucleotides, or alternatively under the control of an exogenous regulatory sequence.

A recombinant expression vector comprising a nucleic acid selected from the group consisting of 5' and 3' regulatory regions, or biologically active fragments or variants thereof, is also part of the present invention.

The invention also encompasses a recombinant expression vector comprising :

a) a nucleic acid comprising the 5' regulatory polynucleotide of the nucleotide sequence  
SEQ ID No 1, or a biologically active fragment or variant thereof;

b) a polynucleotide encoding a polypeptide or a polynucleotide of interest operably linked  
5 with said nucleic acid.

c) optionally, a nucleic acid comprising a 3'-regulatory polynucleotide, preferably a 3'-  
regulatory polynucleotide of the invention, or a biologically active fragment or variant thereof.

The nucleic acid comprising the 5' regulatory polynucleotide or a biologically active  
fragment or variant thereof may also comprises the 5'-UTR sequence from any of the two cDNA of  
10 the invention or a biologically active fragment or variant thereof.

The invention also pertains to a recombinant expression vector useful for the expression of  
the *TBC-1* coding sequence, wherein said vector comprises a nucleic acid selected from the group  
consisting of SEQ ID Nos 3 and 4 or a nucleic acid having at least 95% nucleotide identity with a  
polynucleotide selected from the group consisting of the nucleotide sequences of SEQ ID Nos 3 and

15 4.

Another recombinant expression vector of the invention consists in a recombinant vector  
comprising a nucleic acid comprising the nucleotide sequence beginning at the nucleotide in  
position 176 and ending in position 3730 of the polynucleotide of SEQ ID No 4.

Generally, a recombinant vector of the invention may comprise any of the polynucleotides  
20 described herein, including regulatory sequences, and coding sequences, as well as any *TBC-1*  
primer or probe as defined above. More particularly, the recombinant vectors of the present  
invention can comprise any of the polynucleotides described in the "*TBC-1* cDNA Sequences"  
section, the "Coding Regions" section, "Genomic sequence of *TBC-1*" section and the  
"Oligonucleotide Probes And Primers" section.

25 Some of the elements which can be found in the vectors of the present invention are  
described in further detail in the following sections.

#### a) Vectors

A recombinant vector according to the invention comprises, but is not limited to, a YAC  
(Yeast Artificial Chromosome), a BAC (Bacterial Artificial Chromosome), a phage, a phagemid, a  
30 cosmid, a plasmid or even a linear DNA molecule which may consist of a chromosomal, non-  
chromosomal and synthetic DNA. Such a recombinant vector can comprise a transcriptional unit  
comprising an assembly of :

(1) a genetic element or elements having a regulatory role in gene expression, for example  
promoters or enhancers. Enhancers are cis-acting elements of DNA, usually from about 10 to 300  
35 bp in length that act on the promoter to increase the transcription.

(2) a structural or coding sequence which is transcribed into mRNA and eventually  
translated into a polypeptide, and

T 0 5 3 9 9 0 " T 2 2 9 1 6 0

(3) appropriate transcription initiation and termination sequences. Structural units intended for use in yeast or eukaryotic expression systems preferably include a leader sequence enabling extracellular secretion of translated protein by a host cell. Alternatively, where a recombinant protein is expressed without a leader or transport sequence, it may include an N-terminal residue.

- 5 This residue may or may not be subsequently cleaved from the expressed recombinant protein to provide a final product.

Generally, recombinant expression vectors will include origins of replication, selectable markers permitting transformation of the host cell, and a promoter derived from a highly expressed gene to direct transcription of a downstream structural sequence. The heterologous structural

- 10 sequence is assembled in appropriate phase with translation initiation and termination sequences, and preferably a leader sequence capable of directing secretion of the translated protein into the periplasmic space or the extracellular medium.

The selectable marker genes for selection of transformed host cells are preferably dihydrofolate reductase or neomycin resistance for eukaryotic cell culture, TRP1 for *S. cerevisiae* or

- 15 tetracycline, rifampicin or ampicillin resistance in *E. coli*, or levan saccharase for mycobacteria.

As a representative but non-limiting example, useful expression vectors for bacterial use can comprise a selectable marker and a bacterial origin of replication derived from commercially available plasmids comprising genetic elements of pBR322 (ATCC 37017). Such commercial vectors include, for example, pKK223-3 (Pharmacia, Uppsala, Sweden), and GEM1 (Promega

- 20 Biotec, Madison, WI, USA).

Large numbers of suitable vectors and promoters are known to those of skill in the art, and commercially available, such as bacterial vectors : pQE70, pQE60, pQE-9 (Qiagen), pbs, pD10, phagescript, psiX174, pbluescript SK, pbsks, pNH8A, pNH16A, pNH18A, pNH46A (Stratagene); ptrc99a, pKK223-3, pKK233-3, pDR540, pRIT5 (Pharmacia); or eukaryotic vectors : pWLNEO,

- 25 pSV2CAT, pOG44, pXT1, pSG (Stratagene); pSVK3, pBPV, pMSG, pSVL (Pharmacia); baculovirus transfer vector pVL1392/1393 (Pharminogen); pQE-30 (QIAexpress).

A suitable vector for the expression of the TBC-1 polypeptide of SEQ ID No 5 is a baculovirus vector that can be propagated in insect cells and in insect cell lines. A specific suitable host vector system is the pVL1392/1393 baculovirus transfer vector (Pharminogen) that is used to

- 30 transfect the SF9 cell line (ATCC N<sup>o</sup>CRL 1711) which is derived from *Spodoptera frugiperda*.

Other suitable vectors for the expression of the TBC-1 polypeptide of SEQ ID No 5 in a baculovirus expression system include those described by Chai et al. (1993), Vlasak et al. (1983) and Lenhard et al. (1996).

Mammalian expression vectors will comprise an origin of replication, a suitable promoter

- 35 and enhancer, and also any necessary ribosome binding sites, polyadenylation site, splice donor and acceptor sites, transcriptional termination sequences, and 5' flanking nontranscribed sequences. DNA sequences derived from the SV40 viral genome, for example SV40 origin, early promoter,

enhancer, splice and polyadenylation sites may be used to provide the required nontranscribed genetic elements.

#### b) Promoters

The suitable promoter regions used in the expression vectors according to the present invention are chosen taking into account the cell host in which the heterologous gene has to be expressed.

A suitable promoter may be heterologous with respect to the nucleic acid for which it controls the expression or alternatively can be endogenous to the native polynucleotide containing the coding sequence to be expressed. Additionally, the promoter is generally heterologous with respect to the recombinant vector sequences within which the construct promoter/coding sequence has been inserted.

Preferred bacterial promoters are the LacI, LacZ, the T3 or T7 bacteriophage RNA polymerase promoters, the polyhedrin promoter, or the p10 protein promoter from baculovirus (Kit Novagen) (Smith et al., 1983; O'Reilly et al., 1992), the lambda P<sub>R</sub> promoter or also the trc promoter.

Promoter regions can be selected from any desired gene using, for example, CAT (chloramphenicol transferase) vectors and more preferably pKK232-8 and pCM7 vectors. Particularly preferred bacterial promoters include lacI, lacZ, T3, T7, gpt, lambda PR, PL and trp. Eukaryotic promoters include CMV immediate early, HSV thymidine kinase, early and late SV40, LTRs from retrovirus, and mouse metallothionein-L. Selection of a convenient vector and promoter is well within the level of ordinary skill in the art.

The choice of a promoter is well within the ability of a person skilled in the field of genetic engineering. For example, one may refer to the book of Sambrook et al. (1989) or also to the procedures described by Fullér et al. (1996).

The vector containing the appropriate DNA sequence as described above, more preferably a TBC-1 gene regulatory polynucleotide, a polynucleotide encoding the TBC-1 polypeptide of SEQ ID No 5 or both of them, can be utilized to transform an appropriate host to allow the expression of the desired polypeptide or polynucleotide.

#### c) Other types of vectors

The *in vivo* expression of a TBC-1 polypeptide of SEQ ID No 5 may be useful in order to correct a genetic defect related to the expression of the native gene in a host organism or to the production of a biologically inactive TBC-1 protein.

Consequently, the present invention also deals with recombinant expression vectors mainly designed for the *in vivo* production of the TBC-1 polypeptide of SEQ ID No 5 by the introduction of the appropriate genetic material in the organism of the patient to be treated. This genetic material may be introduced *in vitro* in a cell that has been previously extracted from the organism, the

modified cell being subsequently reintroduced in the said organism, directly *in vivo* into the appropriate tissue.

By « vector » according to this specific embodiment of the invention is intended either a circular or a linear DNA molecule.

5 One specific embodiment for a method for delivering a protein or peptide to the interior of a cell of a vertebrate *in vivo* comprises the step of introducing a preparation comprising a physiologically acceptable carrier and a naked polynucleotide operatively coding for the polypeptide of interest into the interstitial space of a tissue comprising the cell, whereby the naked polynucleotide is taken up into the interior of the cell and has a physiological effect.

10 In a specific embodiment, the invention provides a composition for the *in vivo* production of the TBC-1 protein or polypeptide described herein. It comprises a naked polynucleotide operatively coding for this polypeptide, in solution in a physiologically acceptable carrier, and suitable for introduction into a tissue to cause cells of the tissue to express the said protein or polypeptide.

15 Compositions comprising a polynucleotide are described in PCT application N° WO 90/11092 (Vical Inc.) and also in PCT application N° WO 95/11307 (Institut Pasteur, INSERM, Université d'Ottawa) as well as in the articles of Tacson et al. (1996) and of Huygen et al. (1996).

The amount of vector to be injected to the desired host organism varies according to the site of injection. As an indicative dose, it will be injected between 0,1 and 100 µg of the vector in an animal body, preferably a mammal body, for example a mouse body.

20 In another embodiment of the vector according to the invention, it may be introduced *in vitro* in a host cell, preferably in a host cell previously harvested from the animal to be treated and more preferably a somatic cell such as a muscle cell. In a subsequent step, the cell that has been transformed with the vector coding for the desired TBC-1 polypeptide or the desired fragment thereof is reintroduced into the animal body in order to deliver the recombinant protein within the  
25 body either locally or systemically.

In one specific embodiment, the vector is derived from an adenovirus. Preferred adenovirus vectors according to the invention are those described by Feldman and Steg (1996) or Ohno et al. (1994). Another preferred recombinant adenovirus according to this specific embodiment of the present invention is the human adenovirus type 2 or 5 (Ad 2 or Ad 5) or an adenovirus of animal  
30 origin ( French patent application N° FR-93.05954).

Retrovirus vectors and adeno-associated virus vectors are generally understood to be the recombinant gene delivery systems of choice for the transfer of exogenous polynucleotides *in vivo*, particularly to mammals, including humans. These vectors provide efficient delivery of genes into cells, and the transferred nucleic acids are stably integrated into the chromosomal DNA of the host

35 Particularly preferred retroviruses for the preparation or construction of retroviral *in vitro* or *in vitro* gene delivery vehicles of the present invention include retroviruses selected from the group consisting of Mink-Cell Focus Inducing Virus, Murine Sarcoma Virus, Reticuloendotheliosis virus

09762314-062501  
T05290-T1229260



and Rous Sarcoma virus. Particularly preferred Murine Leukemia Viruses include the 4070A and the 1504A viruses, Abelson (ATCC No VR-999), Friend (ATCC No VR-245), Gross (ATCC No VR-590), Rauscher (ATCC No VR-998) and Moloney Murine Leukemia Virus (ATCC No VR-190; PCT Application No WO 94/24298). Particularly preferred Rous Sarcoma Viruses include

5 Bryan high titer (ATCC Nos VR-334, VR-657, VR-726, VR-659 and VR-728). Other preferred retroviral vectors are those described in Roth et al. (Roth J.A. et al., 1996), PCT Application No WO 93/25234, PCT Application No WO 94/ 06920, Roux et al., 1989, Julan et al., 1992 and Neda et al., 1991.

Yet another viral vector system that is contemplated by the invention consists in the adeno-

10 associated virus (AAV). The adeno-associated virus is a naturally occurring defective virus that requires another virus, such as an adenovirus or a herpes virus, as a helper virus for efficient replication and a productive life cycle (Muzyczka et al., 1992). It is also one of the few viruses that may integrate its DNA into non-dividing cells, and exhibits a high frequency of stable integration (Flotte et al., 1992; Samulski et al., 1989; McLaughlin et al., 1989). One advantageous feature of

15 AAV derives from its reduced efficacy for transducing primary cells relative to transformed cells.

Other compositions containing a vector of the invention advantageously comprise an oligonucleotide fragment of a nucleic sequence selected from the group consisting of SEQ ID Nos 3 or 4 as an antisense tool that inhibits the expression of the corresponding *TBC-1* gene. Preferred methods using antisense polynucleotide according to the present invention are the procedures

20 described by Sczakiel et al. (1995) or those described in PCT Application No WO 95/24223.

### Host cells

Another object of the invention consists in host cell that have been transformed or transfected with one of the polynucleotides described therein, and more precisely a polynucleotide either comprising a *TBC-1* regulatory polynucleotide or the coding sequence of the *TBC-1*

25 polypeptide having the amino acid sequence of SEQ ID No 5. Are included host cells that are transformed (prokaryotic cells) or that are transfected (eukaryotic cells) with a recombinant vector such as one of those described above.

A recombinant host cell of the invention comprises any one of the polynucleotides or the recombinant vectors described therein. More particularly, the cell hosts of the present invention can

30 comprise any of the polynucleotides described in "*TBC-1* cDNA Sequences" section, the "Coding Regions" section, "Genomic sequence of *TBC-1*" section and the "Oligonucleotide Probes And Primers" section.

Another preferred recombinant cell host according to the present invention is characterized in that its genome or genetic background (including chromosome, plasmids) is modified by the

35 nucleic acid coding for the *TBC-1* polypeptide of SEQ ID No 5.

Preferred host cells used as recipients for the expression vectors of the invention are the following :

- a) Prokaryotic host cells : *Escherichia coli* strains (I.E. DH5- $\alpha$  strain) or *Bacillus subtilis*.
- b) Eukaryotic host cells : HeLa cells (ATCC N<sup>o</sup>CCL2; N<sup>o</sup>CCL2.1; N<sup>o</sup>CCL2.2), Cv 1 cells
- 5 (ATCC N<sup>o</sup>CCL70), COS cells (ATCC N<sup>o</sup>CRL1650; N<sup>o</sup>CRL1651), Sf-9 cells (ATCC N<sup>o</sup>CRL1711).

The constructs in the host cells can be used in a conventional manner to produce the gene product encoded by the recombinant sequence.

Following transformation of a suitable host and growth of the host to an appropriate cell density, the selected promoter is induced by appropriate means, such as temperature shift or

10 chemical induction, and cells are cultivated for an additional period.

Cells are typically harvested by centrifugation, disrupted by physical or chemical means, and the resulting crude extract retained for further purification.

Microbial cells employed in the expression of proteins can be disrupted by any convenient method, including freeze-thaw cycling, sonication, mechanical disruption, or use of cell lysing

15 agents. Such methods are well known by the skill artisan.

#### Transgenic animals

The terms "transgenic animals" or "host animals" are used herein to designate animals that have their genome genetically and artificially manipulated so as to include one of the nucleic acids according to the invention. Preferred animals are non-human mammals and include those belonging

20 to a genus selected from *Mus* (e.g. mice), *Rattus* (e.g. rats) and *Oryctogalus* (e.g. rabbits) which have their genome artificially and genetically altered by the insertion of a nucleic acid according to the invention.

The transgenic animals of the invention all include within a plurality of their cells a cloned recombinant or synthetic DNA sequence, more specifically one of the purified or isolated nucleic

25 acids comprising a *TBC-1* coding sequence, a *TBC-1* regulatory polynucleotide or a DNA sequence encoding an antisense polynucleotide such as described in the present specification.

More particularly, transgenic animals according to the invention contain in their somatic cells and/or in their germ line cells any of the polynucleotides described in "TBC-1 cDNA Sequences" section, the "Coding Regions" section, "Genomic sequence of *TBC-1*" section, the

30 "Oligonucleotide Probes And Primers" section and the "Vectors for the expression of a regulatory or coding polynucleotide of *TBC-1*" section.

The transgenic animals of the invention thus contain specific sequences of exogenous genetic material such as the nucleotide sequences described above in detail.

In a first preferred embodiment, these transgenic animals may be good experimental models

35 in order to study the diverse pathologies related to cell differentiation, in particular concerning the

T 0 9 2 9 0 1 1 0 6 2 5 0 1

transgenic animals within the genome of which has been inserted one or several copies of a polynucleotide encoding a native TBC-1 protein, or alternatively a mutant TBC-1 protein.

In a second preferred embodiment, these transgenic animals may express a desired polypeptide of interest under the control of the regulatory polynucleotides of the *TBC-1* gene, leading to good yields in the synthesis of this protein of interest, and eventually a tissue specific expression of this protein of interest.

Since it is possible to produce transgenic animals of the invention using a variety of different sequences, a general description will be given of the production of transgenic animals by referring generally to exogenous genetic material. This general description can be adapted by those skilled in the art in order to incorporate the DNA sequences into animals. For more details regarding the production of transgenic animals, and specifically transgenic mice, it may be referred to Sandou et al. (1994) and also to US Patents Nos 4,873,191, issued Oct.10, 1989, 5,968,766, issued Dec. 16, 1997 and 5,387,742, issued Feb. 28, 1995, these documents being herein incorporated by reference to disclose methods for producing transgenic mice.

Transgenic animals of the present invention are produced by the application of procedures which result in an animal with a genome that incorporates exogenous genetic material which is integrated into the genome. The procedure involves obtaining the genetic material, or a portion thereof, which encodes either a *TBC-1* coding sequence, a *TBC-1* regulatory polynucleotide or a DNA sequence encoding an antisense polynucleotide such as described in the present specification.

A recombinant polynucleotide of the invention is inserted into an embryonic or ES stem cell line. The insertion is made using electroporation. The cells subjected to electroporation are screened (e.g. Southern blot analysis) to find positive cells which have integrated the exogenous recombinant polynucleotide into their genome. An illustrative positive-negative selection procedure that may be used according to the invention is described by Mansour et al. (1988). Then, the positive cells are isolated, cloned and injected into 3.5 days old blastocysts from mice. The blastocysts are then inserted into a female host animal and allowed to grow to term. The offsprings of the female host are tested to determine which animals are transgenic e.g. include the inserted exogenous DNA sequence and which are wild-type.

#### Screening Of Agents Interacting With TBC-1

In a further embodiment, the present invention also concerns a method for the screening of new agents, or candidate substances interacting with TBC-1. These new agents could be useful against cancer.

In a preferred embodiment, the invention relates to a method for the screening of candidate substances comprising the following steps:

- providing a cell line, an organ, or a mammal expressing a *TBC-1* gene or a fragment thereof, preferably the regulatory region or the promoter region of the *TBC-1* gene.

- obtaining a candidate substance preferably a candidate substance capable of inhibiting the binding of a transcription factor to the *TBC-1* regulatory region,
- testing the ability of the candidate substance to decrease the symptoms of prostate cancer and/or to modulate the expression levels of *TBC-1*.

5 In some embodiments, the cell line, organ or mammal expresses a heterologous protein, the coding sequence of which is operably linked to the *TBC-1* regulatory or promoter sequence. In other embodiments, they express a *TBC-1* gene comprising alleles of one or more *TBC-1*-related biallelic markers.

10 A candidate substance is a substance which can interact with or modulate, by binding or other intramolecular interactions, expression, stability, and function of *TBC-1*. Such substances may be potentially interesting for patients who are not responsive to existing drugs or develop side effects to them. Screening may be effected using either *in vitro* methods or *in vivo* methods.

Such methods can be carried out in numerous ways such as on transformed cells which express the considered alleles of the *TBC-1* gene, on tumors induced by said transformed cells, for  
15 example in mice, or on a *TBC-1* protein encoded by the considered allelic variant of *TBC-1*.

Screening assays of the present invention generally involve determining the ability of a candidate substance to present a cytotoxic effect, to change the characteristics of transformed cells such as proliferative and invasive capacity, to affect the tumor growth, or to modify the expression level of *TBC-1*.

20 Typically, this method includes preparing transformed cells with different forms of *TBC-1* sequences containing particular alleles of one or more biallelic markers and/or trait causing mutations described above. This is followed by testing the cells expressing the *TBC-1* with a candidate substance to determine the ability of the substance to present cytotoxic effect, to affect the characteristics of transformed cells, the tumor growth, or to modify the expression level of *TBC-1*.

25 Typical examples of such drug screening assays are provided below. It is to be understood that the parameters set forth in these examples can be modified by the skilled person without undue experimentation.

#### Methods for screening substances interacting with a *TBC-1* polypeptide

30 A method for the screening of a candidate substance according to the invention comprises the following steps :

- a) providing a polypeptide comprising the amino acid sequence SEQ ID No 5, or a peptide fragment or a variant thereof;
- b) obtaining a candidate substance;
- c) bringing into contact said polypeptide with said candidate substance;
- 35 d) detecting the complexes formed between said polypeptide and said candidate substance.

For the purpose of the present invention, a ligand means a molecule, such as a protein, a peptide, an antibody or any synthetic chemical compound capable of binding to the *TBC-1* protein

or one of its fragments or variants or to modulate the expression of the polynucleotide coding for TBC-1 or a fragment or variant thereof.

In the ligand screening method according to the present invention, a biological sample or a defined molecule to be tested as a putative ligand of the TBC-1 protein is brought into contact with a purified TBC-1 protein, for example a purified recombinant TBC-1 protein produced by a recombinant cell host as described hereinbefore, in order to form a complex between the TBC-1 protein and the putative ligand molecule to be tested.

#### A. Candidate ligands obtained from random peptide libraries

In a particular embodiment of the screening method, the putative ligand is the expression product of a DNA insert contained in a phage vector (Parmley and Smith, 1988). Specifically, random peptide phages libraries are used. The random DNA inserts encode peptides of 8 to 20 aminoacids in length (Oldenburg K.R. et al., 1992.; Valadon P., et al., 1996; Lucas A.H., 1994; Westerink M.A.J., 1995; Castagnoli L. et al., 1991). According to this particular embodiment, the recombinant phages expressing a protein that binds to the immobilized TBC-1 protein are retained and the complex formed between the TBC-1 protein and the recombinant phage may be subsequently immunoprecipitated by a polyclonal or a monoclonal antibody directed against the TBC-1 protein.

Once the ligand library in recombinant phages has been constructed, the phage population is brought into contact with the immobilized TBC-1 protein. Then the preparation of complexes is washed in order to remove the non-specifically bound recombinant phages. The phages that bind specifically to the TBC-1 protein are then eluted by a buffer (acid pH) or immunoprecipitated by the anti-TBC-1 monoclonal antibody produced by a hybridoma, and this phage population is subsequently amplified by an over-infection of bacteria (for example E. coli). The selection step may be repeated several times, preferably 2-4 times, in order to select the more specific recombinant phage clones. The last step consists in characterizing the peptide produced by the selected recombinant phage clones either by expression in infected bacteria and isolation, expressing the phage insert in another host-vector system, or sequencing the insert contained in the selected recombinant phages.

#### B. Candidate ligands obtained through a two-hybrid screening assay.

The yeast two-hybrid system is designed to study protein-protein interactions *in vivo* (Fields and Song, 1989), and relies upon the fusion of a bait protein to the DNA binding domain of the yeast Gal4 protein. This technique is also described in US Patent N° US 5,667,973 and US Patent N° 5,283,173 (Fields et al.) the technical teachings of both patents being herein incorporated by reference.

The general procedure of library screening by the two-hybrid assay may be performed as described by Harper et al. (Harper JW et al., 1993) or as described by Cho et al. (1998) or also Fromont-Racine et al. (1997).

The bait protein or polypeptide consists of a TBC-1 polypeptide or a fragment or variant thereof.

More precisely, the nucleotide sequence encoding the TBC-1 polypeptide or a fragment or variant thereof is fused to a polynucleotide encoding the DNA binding domain of the GAL4 protein, the fused nucleotide sequence being inserted in a suitable expression vector, for example pAS2 or pM3.

Then, a human cDNA library is constructed in a specially designed vector, such that the human cDNA insert is fused to a nucleotide sequence in the vector that encodes the transcriptional domain of the GAL4 protein. Preferably, the vector used is the pACT vector. The polypeptides encoded by the nucleotide inserts of the human cDNA library are termed "pray" polypeptides.

A third vector contains a detectable marker gene, such as beta galactosidase gene or CAT gene that is placed under the control of a regulation sequence that is responsive to the binding of a complete Gal4 protein containing both the transcriptional activation domain and the DNA binding domain. For example, the vector pG5EC may be used.

Two different yeast strains are also used. As an illustrative but non limiting example the two different yeast strains may be the following :

- Y190, the phenotype of which is (*MATa, Leu2-3, 112 ura3-12, trp1-901, his3-D200, ade2-101, gal4Dgal180D URA3 GAL-LacZ, LYS GAL-HIS3, cyh'*);
- Y187, the phenotype of which is (*MATa gal4 gal80 his3 trp1-901 ade2-101 ura3-52 leu2-3, - 112 URA3 GAL-lacZmet'*), which is the opposite mating type of Y190.

Briefly, 20 µg of pAS2/TBC-1 and 20 µg of pACT-cDNA library are co-transformed into yeast strain Y190. The transformants are selected for growth on minimal media lacking histidine, leucine and tryptophan, but containing the histidine synthesis inhibitor 3-AT (50 mM). Positive colonies are screened for beta galactosidase by filter lift assay. The double positive colonies (*His<sup>+</sup>, beta-gal<sup>+</sup>*) are then grown on plates lacking histidine, leucine, but containing tryptophan and cycloheximide (10 mg/ml) to select for loss of pAS2/TBC-1 plasmids but retention of pACT-cDNA library plasmids. The resulting Y190 strains are mated with Y187 strains expressing TBC-1 or non-related control proteins; such as cyclophilin B, lamin, or SNF1, as *Gal4* fusions as described by Harper et al. (1993) and by Bram et al. (1993), and screened for beta galactosidase by filter lift assay. Yeast clones that are *beta gal-* after mating with the control *Gal4* fusions are considered false positives.

In another embodiment of the two-hybrid method according to the invention, the interaction between TBC-1 or a fragment or variant thereof with cellular proteins may be assessed using the Matchmaker Two Hybrid System 2 (Catalog No. K1604-1, Clontech). ). As described in the manual accompanying the Matchmaker Two Hybrid System 2 (Catalog No. K1604-1, Clontech), the disclosure of which is incorporated herein by reference, nucleic acids encoding the TBC-1 protein or a portion thereof, are inserted into an expression vector such that they are in frame with DNA encoding the DNA

binding domain of the yeast transcriptional activator GAL4. A desired cDNA, preferably human cDNA, is inserted into a second expression vector such that they are in frame with DNA encoding the activation domain of GAL4. The two expression plasmids are transformed into the yeast cells and the yeast cells are plated on selection medium which selects for expression of selectable markers on each of  
5 the expression vectors as well as GAL4 dependent expression of the HIS3 gene. Transformants capable of growing on medium lacking histidine are screened for GAL4 dependent lacZ expression. Those cells which are positive in both the histidine selection and the lacZ assay are those in which an interaction between TBC-1 and the protein or peptide encoded by the initially selected cDNA insert has taken place.

#### 10 Method for screening ligands that modulate the expression of the *TBC-1* gene.

Another subject of the present invention is a method for screening molecules that modulate the expression of the TBC-1 protein. Such a screening method comprises the steps of :

- a) cultivating a prokaryotic or an eukaryotic cell that has been transfected with a nucleotide sequence encoding the TBC-1 protein, operably linked to a *TBC-1* 5'-regulatory sequence;
- 15 b) bringing into contact the cultivated cell with a molecule to be tested;
- c) quantifying the expression of the TBC-1 protein.

Using DNA recombination techniques well known by the one skill in the art, the TBC-1 protein encoding DNA sequence is inserted into an expression vector, downstream from a *TBC-1* 5'-regulatory sequence that contains a *TBC-1* promoter sequence.

- 20 The quantification of the expression of the TBC-1 protein may be realized either at the mRNA level or at the protein level. In the latter case, polyclonal or monoclonal antibodies may be used to quantify the amounts of the TBC-1 protein that have been produced, for example in an ELISA or a RIA assay.

- In a preferred embodiment, the quantification of the *TBC-1* mRNAs is realized by a  
25 quantitative PCR amplification of the cDNAs obtained by a reverse transcription of the total mRNA of the cultivated *TBC-1*-transfected host cell, using a pair of primers specific for *TBC-1*.

- Expression levels and patterns of *TBC-1* may be analyzed by solution hybridization with long probes as described in International Patent Application No. WO 97/05277, the entire contents of which are incorporated herein by reference. Briefly, the *TBC-1* cDNA or the *TBC-1* genomic DNA described  
30 above, or fragments thereof, is inserted at a cloning site immediately downstream of a bacteriophage (T3, T7 or SP6) RNA polymerase promoter to produce antisense RNA. Preferably, the *TBC-1* insert comprises at least 100 or more consecutive nucleotides of the genomic DNA sequence or the cDNA sequences, particularly those comprising one of the nucleotide sequences of SEQ ID Nos 3, 4 and 6-8 or those encoding a mutated TBC-1. The plasmid is linearized and transcribed in the presence of  
35 ribonucleotides comprising modified ribonucleotides (i.e. biotin-UTP and DIG-UTP). An excess of this doubly labeled RNA is hybridized in solution with mRNA isolated from cells or tissues of interest. The hybridizations are performed under standard stringent conditions (40-50°C for 16 hours in an 80%

- formamide, 0.4 M NaCl buffer, pH 7-8). The unhybridized probe is removed by digestion with ribonucleases specific for single-stranded RNA (i.e. RNases CL3, T1, Phy M, U2 or A). The presence of the biotin-UTP modification enables capture of the hybrid on a microtitration plate coated with streptavidin. The presence of the DIG modification enables the hybrid to be detected and quantified by
- 5 ELISA using an anti-DIG antibody coupled to alkaline phosphatase.

- Quantitative analysis of *TBC-1* gene expression may also be performed using arrays. As used herein, the term array means a one dimensional, two dimensional, or multidimensional arrangement of a plurality of nucleic acids of sufficient length to permit specific detection of expression of mRNAs capable of hybridizing thereto. For example, the arrays may contain a
- 10 plurality of nucleic acids derived from genes whose expression levels are to be assessed. The arrays may include the *TBC-1* genomic DNA, the *TBC-1* cDNA sequences or the sequences complementary thereto or fragments thereof, particularly those comprising at least one of the biallelic markers according to the present invention. Preferably, the fragments are at least 15 nucleotides in length. In other embodiments, the fragments are at least 25 nucleotides in length. In
- 15 some embodiments, the fragments are at least 50 nucleotides in length. More preferably, the fragments are at least 100 nucleotides in length. In another preferred embodiment, the fragments are more than 100 nucleotides in length. In some embodiments the fragments may be more than 500 nucleotides in length.

- For example, quantitative analysis of *TBC-1* gene expression may be performed with a
- 20 complementary DNA microarray as described by Schena et al. (1995). Full length *TBC-1* cDNAs or fragments thereof are amplified by PCR and arrayed from a 96-well microtiter plate onto silylated microscope slides using high-speed robotics. Printed arrays are incubated in a humid chamber to allow rehydration of the array elements and rinsed, once in 0.2% SDS for 1 min, twice in water for 1 min and once for 5 min in sodium borohydride solution. The arrays are submerged in water for 2
- 25 min at 95°C, transferred into 0.2% SDS for 1 min, rinsed twice with water, air dried and stored in the dark at 25°C.

- Cell or tissue mRNA is isolated or commercially obtained and probes are prepared by a single round of reverse transcription. Probes are hybridized to 1 cm<sup>2</sup> microarrays under a 14 x 14 mm glass coverslip for 6-12 hours at 60°C. Arrays are washed for 5 min at 25°C in low stringency
- 30 wash buffer (1 x SSC/0.2% SDS), then for 10 min at room temperature in high stringency wash buffer (0.1 x SSC/0.2% SDS). Arrays are scanned in 0.1 x SSC using a fluorescence laser scanning device fitted with a custom filter set. Accurate differential expression measurements are obtained by taking the average of the ratios of two independent hybridizations.

- Quantitative analysis of *TBC-1* gene expression may also be performed with full length
- 35 *TBC-1* cDNAs or fragments thereof in complementary DNA arrays as described by Pietu et al. (1996). The full length *TBC-1* cDNA or fragments thereof is PCR amplified and spotted on membranes. Then, mRNAs originating from various tissues or cells are labeled with radioactive



nucleotides. After hybridization and washing in controlled conditions, the hybridized mRNAs are detected by phospho-imaging or autoradiography. Duplicate experiments are performed and a quantitative analysis of differentially expressed mRNAs is then performed.

Alternatively, expression analysis using the *TBC-1* genomic DNA, the *TBC-1* cDNAs, or  
5 fragments thereof can be done through high density nucleotide arrays or chips as described by Lockhart et al. (1996) and Sosnowsky et al. (1997). Oligonucleotides of 15-50 nucleotides from the sequences of the *TBC-1* genomic DNA, the *TBC-1* cDNA sequences particularly those comprising at least one of biallelic markers according the present invention, preferably at least one of SEQ ID No 7-8 or those comprising the trait causing mutation, or the sequences complementary thereto, are  
10 synthesized directly on the chip (Lockhart et al., supra) or synthesized and then addressed to the chip (Sosnowski et al., supra). Preferably, the oligonucleotides are about 20 nucleotides in length.

*TBC-1* cDNA probes labeled with an appropriate compound, such as biotin, digoxigenin or fluorescent dye, are synthesized from the appropriate mRNA population and then randomly  
15 fragmented to an average size of 50 to 100 nucleotides. The said probes are then hybridized to the chip. After washing as described in Lockhart et al., supra and application of different electric fields (Sosnowsky et al., 1997), the dyes or labeling compounds are detected and quantified. Duplicate hybridizations are performed. Comparative analysis of the intensity of the signal originating from cDNA probes on the same target oligonucleotide in different cDNA samples indicates a differential expression of *TBC-1* mRNAs.

20 Thus, is also part of the present invention a method for screening of a candidate substance or molecule that modulates the expression of the *TBC-1* gene according to the invention, wherein this method comprises the following steps :

a) providing a recombinant cell host containing a nucleic acid, wherein said nucleic acid comprises the 5' regulatory region sequence or a biologically active fragment or variant thereof, the  
25 5' regulatory region or its biologically active fragment or variant being operably linked to a polynucleotide encoding a detectable protein;

b) obtaining a candidate substance, and

c) determining the ability of the candidate substance to modulate the expression levels of the polynucleotide encoding the detectable protein.

30 In a preferred embodiment of the above screening method, the nucleic acid comprising the 5' regulatory region sequence or a biologically active fragment or variant thereof also includes a 5'UTR region of one of the *TBC-1* cDNAs of SEQ ID Nos 3 and 4, or one of their biologically active fragments or variants thereof.

A second method for the screening of a candidate substance or molecule that modulates the  
35 expression of the *TBC-1* gene comprises the following steps :

a) providing a recombinant cell host containing a nucleic acid, wherein said nucleic acid comprises a 5'UTR sequence of one of the *TBC-1* cDNAs of SEQ ID Nos 3 and 4, or one of their

09762311-06501  
T0660-TE29Z60

biologically active fragments or variants, the 5'UTR sequence or its biologically active fragment or variant being operably linked to a polynucleotide encoding a detectable protein;

b) obtaining a candidate substance, and

c) determining the ability of the candidate substance to modulate the expression levels of the polynucleotide encoding the detectable protein.

In a preferred embodiment of the screening method described above, the nucleic acid that comprises a nucleotide sequence selected from the group consisting of the 5'UTR sequence of one of the *TBC-1* cDNAs of SEQ ID Nos 3 and 4 or one of their biologically active fragments or variants, includes a promoter sequence, wherein said promoter sequence can be either endogenous, or in contrast exogenous with respect to the *TBC-1* 5'UTR sequences defined therein.

Among the preferred polynucleotides encoding a detectable protein, there may be cited polynucleotides encoding beta galactosidase, green fluorescent protein (GFP) and chloramphenicol acetyl transferase (CAT).

For the design of suitable recombinant vectors useful for performing the screening methods described above, it will be referred to the section of the present specification wherein the preferred recombinant vectors of the invention are detailed.

#### Screening using transgenic animals

*In vivo* methods can utilize transgenic animals for drug screening. Nucleic acids including at least one of the biallelic polymorphisms of interest can be used to generate genetically modified non-human animals or to generate site specific gene modifications in cell lines. The term "transgenic" is intended to encompass genetically modified animals having a deletion or other knock-out of *TBC-1* gene activity, having an exogenous *TBC-1* gene that is stably transmitted in the host cells, or having an exogenous *TBC-1* promoter operably linked to a reporter gene. Transgenic animals may be made through homologous recombination, where the *TBC-1* locus is altered.

Alternatively, a nucleic acid construct is randomly integrated into the genome. Vectors for stable integration include for example plasmids, retroviruses and other animal viruses, and YACs. Of interest are transgenic mammals e.g. cows, pigs, goats, horses, and particularly rodents such as rats and mice. Transgenic animals allow to study both efficacy and toxicity of the candidate drug.

#### Methods for inhibiting the expression of a *TBC-1* gene

Other therapeutic compositions according to the present invention comprise advantageously an oligonucleotide fragment of the nucleic sequence of *TBC-1* as an antisense tool that inhibits the expression of the corresponding *TBC-1* gene. Preferred methods using antisense polynucleotide according to the present invention are the procedures described by Sczakiel et al. (1995).

Preferably, the antisense tools are chosen among the polynucleotides (15-200 bp long) that are complementary to the 5' end of the *TBC-1* mRNA. In another embodiment, a combination of different antisense polynucleotides complementary to different parts of the desired targetted gene are used.

Preferred antisense polynucleotides according to the present invention are complementary to a sequence of the mRNAs of *TBC-1* that contains the translation initiation codon ATG.

The antisense nucleic acid molecules to be used in gene therapy may be either DNA or RNA sequences. They comprise a nucleotide sequence complementary to the targeted sequence of the PTCA-1 genomic DNA, the sequence of which can be determined using one of the detection methods of the present invention. The targeted DNA or RNA sequence preferably comprises at least one of the biallelic markers according to the present invention. The antisense nucleic acids should have a length and melting temperature sufficient to permit formation of an intracellular duplex having sufficient stability to inhibit the expression of the *TBC-1* mRNA in the duplex. Strategies for designing antisense nucleic acids suitable for use in gene therapy are disclosed in Green et al., (1986) and Izant and Weintraub, (1984), the disclosures of which are incorporated herein by reference.

In some strategies, antisense molecules are obtained by reversing the orientation of the *TBC-1* coding region with respect to a promoter so as to transcribe the opposite strand from that which is normally transcribed in the cell. The antisense molecules may be transcribed using in vitro transcription systems such as those which employ T7 or SP6 polymerase to generate the transcript. Another approach involves transcription of *TBC-1* antisense nucleic acids in vivo by operably linking DNA containing the antisense sequence to a promoter in a suitable expression vector.

Alternatively, suitable antisense strategies are those described by Rossi et al. (1991), in the International Applications Nos. WO 94/23026, WO 95/04141, WO 92/18522 and in the European Patent Application No. EP 0 572 287 A2

An alternative to the antisense technology that is used according to the present invention consists in using ribozymes that will bind to a target sequence via their complementary polynucleotide tail and that will cleave the corresponding RNA by hydrolyzing its target site (namely « hammerhead ribozymes »). Briefly, the simplified cycle of a hammerhead ribozyme consists of (1) sequence specific binding to the target RNA via complementary antisense sequences; (2) site-specific hydrolysis of the cleavable motif of the target strand; and (3) release of cleavage products, which gives rise to another catalytic cycle. Indeed, the use of long-chain antisense polynucleotide (at least 30 bases long) or ribozymes with long antisense arms are advantageous. A preferred delivery system for antisense ribozyme is achieved by covalently linking these antisense ribozymes to lipophilic groups or to use liposomes as a convenient vector. Preferred antisense ribozymes according to the present invention are prepared as described by Sczakiel et al. (1995), the specific preparation procedures being referred to in said article being herein incorporated by reference.

Throughout this application, various publications, patents and published patent applications are cited. The disclosures of these publications, patents and published patent specification

referenced in this application are hereby incorporated by reference into the present disclosure to more fully describe the state of the art to which this invention pertains.

## EXAMPLES

### EXAMPLE 1 :

#### 5      **Analysis of the first mRNA encoding a TBC-1 polypeptide synthesized by the cells.**

*TBC-1* cDNA was obtained as follows : 4 µl of ethanol suspension containing 1 mg of human prostate total RNA (Clontech laboratories, Inc., Palo Alto, USA; Catalogue N. 64038-1) was centrifuged, and the resulting pellet was air dried for 30 minutes at room temperature.

First strand cDNA synthesis was performed using the Advantage™ RT-for-PCR kit  
10 (Clontech laboratories Inc., catalogue N. K1402-1). 1 µl of 20 mM solution of a specific oligo dT primer was added to 12.5 µl of RNA solution in water, heated at 74°C for 2.5 min and rapidly quenched in an ice bath. 10 µl of 5 x RT buffer (50 mM Tris-HCl, pH 8.3, 75 mM KCl, 3 mM MgCl<sub>2</sub>), 2.5 µl of dNTP mix (10 mM each), 1.25 µl of human recombinant placental RNA inhibitor were mixed with 1 ml of MMLV reverse transcriptase (200 units). 6.5 µl of this solution were  
15 added to RNA-primer mix and incubated at 42°C for one hour. 80 µl of water were added and the solution was incubated at 94°C for 5 minutes.

5 µl of the resulting solution were used in a Long Range PCR reaction with hot start, in 50 µl final volume, using 2 units of rT<sub>HXL</sub>, 20 pmol/µl of each of 5'-TGACCACCATGCCCCATGCT-3' (271-289 in SEQ ID No 3) and 5'-  
20 GCATTATTCACGTCCACGCC-3' (3929-3949 in SEQ ID No 3) primers with 35 cycles of elongation for 6 minutes at 67°C in thermocycler.

The amplification products corresponding to both cDNA strands were partially sequenced in order to ensure the specificity of the amplification reaction.

Results of Northern blot analysis of prostate mRNAs supported the existence of the first  
25 *TBC-1* cDNA having about 4 kb in length, which is the nucleotide sequence of SEQ ID No 3.

### Example 2 :

#### **Detection of *TBC-1* biallelic markers: DNA extraction**

Donors were unrelated and healthy. They presented a sufficient diversity for being representative of a French heterogeneous population. The DNA from 100 individuals was extracted  
30 and tested for the detection of the biallelic markers.

30 ml of peripheral venous blood were taken from each donor in the presence of EDTA. Cells (pellet) were collected after centrifugation for 10 minutes at 2000 rpm. Red cells were lysed by a lysis solution (50 ml final volume : 10 mM Tris pH7.6; 5 mM MgCl<sub>2</sub>; 10 mM NaCl). The

solution was centrifuged (10 minutes, 2000 rpm) as many times as necessary to eliminate the residual red cells present in the supernatant, after resuspension of the pellet in the lysis solution.

The pellet of white cells was lysed overnight at 42°C with 3.7 ml of lysis solution composed of:

- 5                   - 3 ml TE 10-2 (Tris-HCl 10 mM, EDTA 2 mM) / NaCl 0.4 M
- 200 µl SDS 10%
- 500 µl K-proteinase (2 mg K-proteinase in TE 10-2 / NaCl 0.4 M).

For the extraction of proteins, 1 ml saturated NaCl (6M) (1/3.5 v/v) was added. After vigorous agitation, the solution was centrifuged for 20 minutes at 10000 rpm.

- 10           For the precipitation of DNA, 2 to 3 volumes of 100% ethanol were added to the previous supernatant, and the solution was centrifuged for 30 minutes at 2000 rpm. The DNA solution was rinsed three times with 70% ethanol to eliminate salts, and centrifuged for 20 minutes at 2000 rpm. The pellet was dried at 37°C, and resuspended in 1 ml TE 10-1 or 1 ml water. The DNA concentration was evaluated by measuring the OD at 260 nm (1 unit OD = 50 µg/ml DNA).

- 15           To determine the presence of proteins in the DNA solution, the OD 260 / OD 280 ratio was determined. Only DNA preparations having a OD 260 / OD 280 ratio between 1.8 and 2 were used in the subsequent examples described below.

The pool was constituted by mixing equivalent quantities of DNA from each individual.

### Example 3 :

#### 20           **Detection of the biallelic markers: amplification of genomic DNA by PCR**

The amplification of specific genomic sequences of the DNA samples of example 2 was carried out on the pool of DNA obtained previously. In addition, 50 individual samples were similarly amplified.

PCR assays were performed using the following protocol:

25	Final volume	25 µl
	DNA	2 ng/µl
	MgCl <sub>2</sub>	2 mM
	dNTP (each)	200 µM
	primer (each)	2.9 ng/µl
30	Ampli Taq Gold DNA polymerase	0.05 unit/µl
	PCR buffer (10x = 0.1 M TrisHCl pH8.3 0.5M KCl	1x

Each pair of first primers was designed using the sequence information of the *TBC-1* gene disclosed herein and the OSP software (Hillier & Green, 1991). This first pair of primers was about 20 nucleotides in length and had the sequences disclosed in Table 1 in the columns labeled PU and RP.

Table 1

Amplicon	Position range of the amplicon in SEQ ID 1		Primer name	Position range of amplification primer in SEQ ID No 1		Primer name	Complementary position range of amplification primer in SEQ ID No 1	
99-430	9391	9845	B1	9391	9408	C1	9828	9845
Amplicon	Position range of the amplicon in SEQ ID 2		Primer name	Position range of amplification primer in SEQ ID No 2		Primer name	Complementary position range of amplification primer in SEQ ID No 2	
99-20508	988	1529	B2	988	1006	C2	1509	1529
99-20469	5039	5554	B3	5039	5056	C3	5534	5554
5-254	5997	6350	B4	5997	6015	C4	6332	6350
5-257	14371	14817	B5	14371	14390	C5	14798	14817
99-20511	18751	19217	B6	18751	18771	C6	19198	19217
99-20510	19605	20005	B7	19605	19625	C7	19986	20005
99-20504	29529	30061	B8	29529	29547	C8	30041	30061
99-20493	42268	42752	B9	42268	42287	C9	42732	42752
99-20499	69026	69543	B10	69026	69046	C10	69525	69543
99-20473	76323	76790	B11	76323	76343	C11	76771	76790
5-249	78292	78721	B12	78292	78309	C12	78704	78721
99-20485	81893	82372	B13	81893	81912	C13	82353	82372
99-20481	84392	84929	B14	84392	84412	C14	84909	84929
99-20480	89746	90198	B15	89746	89765	C15	90179	90198

Preferably, the primers contained a common oligonucleotide tail upstream of the specific bases targeted for amplification which was useful for sequencing.

5 Primers PU contain the following additional PU 5' sequence :

TGTAACGACGGCCAGT (SEQ ID No 6); primers RP contain the following RP 5' sequence : CAGGAAACAGCTATGACC (SEQ ID No 7).

The synthesis of these primers was performed following the phosphoramidite method, on a GENSET UFPS 24.1 synthesizer.

10 DNA amplification was performed on a Genius II thermocycler. After heating at 95°C for 10 min, 40 cycles were performed. Each cycle comprised: 30 sec at 95°C, 54°C for 1 min, and 30 sec at 72°C. For final elongation, 10 min at 72°C ended the amplification. The quantities of the amplification products obtained were determined on 96-well microtiter plates, using a fluorometer and Picogreen as intercalant agent (Molecular Probes).

15 **Example 4 :**

**Detection of the biallelic markers: sequencing of amplified genomic DNA and identification of polymorphisms.**

The sequencing of the amplified DNA obtained in example 3 was carried out on ABI 377 sequencers. The sequences of the amplification products were determined using automated dideoxy  
20 terminator sequencing reactions with a dye terminator cycle sequencing protocol. The products of

the sequencing reactions were run on sequencing gels and the sequences were determined using gel image analysis [ABI Prism DNA Sequencing Analysis software (2.1.2 version)].

The sequence data were further evaluated to detect the presence of biallelic markers among the pooled amplified fragments. The polymorphism search was based on the presence of

- 5 superimposed peaks in the electrophoresis pattern resulting from different bases occurring at the same position as described previously.

15 fragments of amplification was analyzed. In this segment, 19 biallelic markers were detected. The localization of the biallelic marker is as shown in Table 2.

Table 2

Amplicon	BM	Marker Name	Localization in <i>TBC-1</i> gene	Polymorphism		BM position in SEQ ID No 1
				Allele 1	allele 2	
99-430	A1	99-430-352	Intron 1	A	G	9494
Amplicon	BM	Marker Name	Localization in <i>TBC-1</i> gene	Polymorphism		BM position in SEQ ID No 2
				allele 1	allele 2	
99-20508	A2	99-20508-456	Intron upstream to Exon A	C	T	1443
99-20469	A3	99-20469-213	Intron A	C	T	5247
5-254	A4	5-254-227	Intron B	A	G	6223
5-257	A5	5-257-353	Intron D	C	T	14723
99-20511	A6	99-20511-32	Intron D	C	T	19186
99-20511	A7	99-20511-221	Intron D	A	G	18997
99-20510	A8	99-20510-115	Intron D	deletion of TCT		19891
99-20504	A9	99-20504-90	Intron D	A	G	29617
99-20493	A10	99-20493-238	Intron D	A	C	42519
99-20499	A11	99-20499-221	Intron G	A	G	69324
99-20499	A12	99-20499-364	Intron G	A	T	69181
99-20499	A13	99-20499-399	Intron G	A	G	69146
99-20473	A14	99-20473-138	Intron H	deletion of TAACA		76458
5-249	A15	5-249-304	Intron I	A	G	78595
99-20485	A16	99-20485-269	Intron I	A	G	82159
99-20481	A17	99-20481-131	Intron I	G	C	84522
99-20481	A18	99-20481-419	Intron I	A	T	84810
99-20480	A19	99-20480-233	Intron J	A	G	89967

- 10 BM refers to "biallelic marker". All1 and all2 refer respectively to allele 1 and allele 2 of the biallelic marker.

Table 3

BM	Marker Name	Position range of probes in SEQ ID No 1		Probes
A1	99-430-352	9482	9506	P1

BM	Marker Name	Position range of probes in SEQ ID No 2		Probes
A2	99-20508-456	1431	1455	P2
A3	99-20469-213	5235	5259	P3
A4	5-254-227	6211	6235	P4
A5	5-257-353	14711	14735	P5
A6	99-20511-32	19174	19198	P6
A7	99-20511-221	18985	19009	P7
A9	99-20504-90	29605	29629	P9
A10	99-20493-238	42507	42531	P10
A11	99-20499-221	69312	69336	P11
A12	99-20499-364	69169	69193	P12
A13	99-20499-399	69134	69158	P13
A15	5-249-304	78583	78607	P15
A16	99-20485-269	82147	82171	P16
A17	99-20481-131	84510	84534	P17
A18	99-20481-419	84798	84822	P18
A19	99-20480-233	89955	89979	P19

#### Example 5 :

##### Validation of the polymorphisms through microsequencing

The biallelic markers identified in example 4 were further confirmed and their respective frequencies were determined through microsequencing. Microsequencing was carried out for each individual DNA sample described in Example 2.

Amplification from genomic DNA of individuals was performed by PCR as described above for the detection of the biallelic markers with the same set of PCR primers (Table 1).

The preferred primers used in microsequencing were about 19 nucleotides in length and hybridized just upstream of the considered polymorphic base. According to the invention, the primers used in microsequencing are detailed in Table 4.

Table 4

Marker Name	Biallelic Marker	Mis. 1	Position range of microsequencing primer mis 1 in SEQ ID No 1		Mis. 2	Complementary position range of microsequencing primer mis. 2 in SEQ ID No 1	
99-430-352	A1	D1	9475	9493	E1	9495	9513
Marker Name	Biallelic Marker	Mis. 1	Position range of microsequencing primer mis 1 in SEQ ID No 2		Mis. 2	Complementary position range of microsequencing primer mis. 2 in SEQ ID No 2	
99-20508-456	A2	D2	1424	1442	E2	1444	1462
99-20469-213	A3	D3	5228	5246	E3	5248	5266
5-254-227	A4	D4	6204	6222	E4	6224	6242
5-257-353	A5	D5	14704	14722	E5	14724	14742
99-20511-32	A6	D6	19167	19185	E6	19187	19205
99-20511-221	A7	D7	18978	18996	E7	18998	19016



99-20510-115	A8	D8	19872	19890	E8	19892	19910
99-20504-90	A9	D9	29598	29616	E9	29618	29636
99-20493-238	A10	D10	42500	42518	E10	42520	42538
99-20499-221	A11	D11	69305	69323	E11	69325	69343
99-20499-364	A12	D12	69162	69180	E12	69182	69200
99-20499-399	A13	D13	69127	69145	E13	69147	69165
99-20473-138	A14	D14	76439	76457	E14	76459	76477
5-249-304	A15	D15	78576	78594	E15	78596	78614
99-20485-269	A16	D16	82140	82158	E16	82160	82178
99-20481-131	A17	D17	84503	84521	E17	84523	84541
99-20481-419	A18	D18	84791	84809	E18	84811	84829
99-20480-233	A19	D19	89948	89966	E19	89968	89986

The microsequencing reaction was performed as follows :

After purification of the amplification products, the microsequencing reaction mixture was prepared by adding, in a 20µl final volume: 10 pmol microsequencing oligonucleotide, 1 U

- 5 Thermosequase (Amersham E79000G), 1.25 µl Thermosequase buffer (260 mM Tris HCl pH 9.5, 65 mM MgCl<sub>2</sub>), and the two appropriate fluorescent ddNTPs (Perkin Elmer, Dye Terminator Set 401095) complementary to the nucleotides at the polymorphic site of each biallelic marker tested, following the manufacturer's recommendations. After 4 minutes at 94°C, 20 PCR cycles of 15 sec at 55°C, 5 sec at 72°C, and 10 sec at 94°C were carried out in a Tetrad PTC-225
- 10 thermocycler (MJ Research). The unincorporated dye terminators were then removed by ethanol precipitation. Samples were finally resuspended in formamide-EDTA loading buffer and heated for 2 min at 95°C before being loaded on a polyacrylamide sequencing gel. The data were collected by an ABI PRISM 377 DNA sequencer and processed using the GENESCAN software (Perkin Elmer).

Following gel analysis, data were automatically processed with software that allows the

- 15 determination of the alleles of biallelic markers present in each amplified fragment.

The software evaluates such factors as whether the intensities of the signals resulting from the above microsequencing procedures are weak, normal, or saturated, or whether the signals are ambiguous. In addition, the software identifies significant peaks (according to shape and height criteria). Among the significant peaks, peaks corresponding to the targeted site are identified based

- 20 on their position. When two significant peaks are detected for the same position, each sample is categorized classification as homozygous or heterozygous type based on the height ratio.

## References

- Altschul et al., 1990, J. Mol. Biol. 215(3):403-410 / Altschul et al., 1993, Nature Genetics 3:266-272 / Altschul et al., 1997, Nuc. Acids Res. 25:3389-3402 / Ausubel et al. (1989)Current Protocols in Molecular Biology, Green Publishing Associates and Wiley Interscience, N.Y. / Beaucage et al., *Tetrahedron Lett* 1981, 22: 1859-1862 / Bram RJ et al., 1993, Mol. Cell Biol., 13 : 4760-4769. / Brown EL, Belagaje R, Ryan MJ, Khorana HG, *Methods Enzymol* 1979;68:109-151 / Castagnoli L. et al. (Felici F.), 1991, J. Mol. Biol., 222:301-310. /
- 25

- Chai H. et al., 1993, *Biotechnol. Appl. Biochem.*, **18**:259-273 / Chee et al. (1996) *Science*. 274:610-614. / Chen and Kwok *Nucleic Acids Research* 25:347-353 1997 / Chen et al. *Proc. Natl. Acad. Sci. USA* 94/20 10756-10761, 1997 / Cho RJ et al., 1998, *Proc. Natl. Acad. Sci. USA*, **95**(7) : 3752-3757. / Chumakov I. et al., 1995, *Nature*, **377**(6547 Suppl): 175-297. / Compton J.
- 5 (1991) *Nature*. 350(6313):91-92. / Dib et al., 1996, *Nature*, **380**: III-V. / Ellis NA, 1997 *Curr. Op. Genet. Dev.*, **7** : 354-363 / Feldman and Steg, 1996, *Medecine/Sciences, synthese*, **12**:47-55 / Fields and Song, 1989, *Nature*, Vol. 340 : 245-246. / Fishel R & Wilson T. 1997, *Curr. Op. Genet. Dev.* **7**: 105-113 / Flotte et al., 1992, *Am. J. Respir. Cell Mol. Biol.*, **7** : 349-356. / Fodor et al. (1991) *Science* 251:767-777. / Fromont-Racine M. et al., 1997, *Nature Genetics*,
- 10 **16**(3) : 277-282. / Fuller S.A. et al., 1996, *Immunology in Current Protocols in Molecular Biology*, Ausubel et al. Eds, John Wiley & Sons, Inc., USA / Geysen H. Mario et al. 1984. *Proc. Natl. Acad. Sci. U.S.A.* **81**:3998-4002 / Gonnet et al., 1992, *Science* 256:1443-1445 / Green et al., *Ann. Rev. Biochem.* **55**:569-597 (1986) / Grompe, M. et al., *Proc. Natl. Acad. Sci. U.S.A* 1989; **86**:5855-5892 / Grompe, M. *Nature Genetics* 1993; **5**:111-117 / Guatelli J C et al. *Proc. Natl.*
- 15 *Acad. Sci. USA*. **35**:273-286. / Haber D & Harlow E, 1997, *Nature Genet.* **16**:320-322. / Hacia JG, Brody LC, Chee MS, Fodor SP, Collins FS, *Nat Genet* 1996;**14**(4):441-447 / Haff L. A. and Smirnov I. P. (1997) *Genome Research*, **7**:378-388. / Hames B.D. and Higgins S.J. (1985) *Nucleic Acid Hybridization: A Practical Approach*. Hames and Higgins Ed., IRL Press, Oxford. / Harju L, et al., *Clin Chem* 1993;**39**(11Pt 1):2282-2287 / Harper JW et al., 1993, *Cell*, Vol. **75** :
- 20 805-816. / Harris H et al., 1969, *Nature* 223:363-368. / Henikoff and Henikoff, 1993, *Proteins* **17**:49-61 / Higgins et al., 1996, *Methods Enzymol.* **266**:383-402 / Hillier L. and Green P. *Methods Appl.*, 1991, **1**: 124-8. / Huang L. et al. (1996) *Cancer Res* **56**(5):1137-1141. / Huygen et al., 1996, *Nature Medicine*, **2**(8):893-898 / Izant and Weintraub, *Cell* **36**:1007-1015 (1984) / Julan et al., 1992, *J. Gen. Virol.*, **73** : 3251 - 3255. / Karlin and Altschul, 1990, *Proc. Natl. Acad. Sci. USA* **87**:2267-2268 / Koch Y., 1977, *Biochem. Biophys. Res. Commun.*, **74**:488-491 /
- 25 Kohler G. and Milstein C., 1975, *Nature*, **256** : 495. / Kozal MJ, et al., *Nat Med* 1996;**2**(7):753-759 / Landegren U. et al. (1998) *Genome Research*, **8**:769-776. / Leger OJ, et al., 1997, *Hum Antibodies*, **8**(1): 3-16 / Lenhard T. et al., 1996, *Gene*, **169**:187-190 / Livak et al., *Nature Genetics*, **9**:341-342, 1995 / Livak KJ, and Hainer JW., 1994, *Hum Mutat.*, **3**(4): 379-385. /
- 30 Lockhart et al. *Nature Biotechnology* **14**: 1675-1680, 1996 / Lucas A.H., 1994, In : *Development and Clinical Uses of Haemophilus b Conjugate*. / Mansour SL et al., 1988, *Nature*, **336** : 348-352. / Marshall R. L. et al. (1994) *PCR Methods and Applications*. **4**:80-84. / Martineau P, Jones P, Winter G, 1998, *J Mol Biol*, **280**(1):117-127 / Mc Whorter W.P., et al. A screening study of prostate cancer in high risk families. *J Urol* 1992;**148**:826-828. / McLaughlin et al., 1989, *J. Virol.*,
- 35 **62** : 1963 - 1973. / Muzyczka et al., 1992, *Curr. Topics in Micro. and Immunol.*, **158** : 97-129. / Narang SA, Hsiung HM, Brousseau R, *Methods Enzymol* 1979;**68**:90-98 / Neda et al., 1991, *J. Biol. Chem.*, **266** : 14143 - 14146. / Nickerson D.A. et al. (1990) *Proc. Natl. Acad. Sci. U.S.A.*

- 87:8923-8927. / Nyren P, Pettersson B, Uhlen M, *Anal Biochem* 1993;208(1):171-175 / O'Reilly et al., 1992, Baculovirus expression vectors : a Laboratory Manual. W.H. Freeman and Co., New York / Ohno et al., 1994, *Sciences*, 265:781-784 / Oldenburg K.R. et al., 1992, *Proc. Natl. Acad. Sci.*, 89:5393-5397. / Orita et al., *Proc. Natl. Acad. Sci. U.S.A.* 1989;86: 2776-2770 /
- 5 Parmley and Smith, *Gene*, 1988, 73:305-318. / Pastinen et al., *Genome Research* 1997; 7:606-614 / PCR Methods and Applications", 1991, Cold Spring Harbor Laboratory Press. / Pearson and Lipman, 1988, *Proc. Natl. Acad. Sci. USA* 85(8):2444-2448 / Pietu et al. *Genome Research* 6:492-503, 1996 / Porath J et al., 1975, *Nature*, 258(5536) : 598-599. / Reimann KA, et al., 1997, *AIDS Res Hum Retroviruses*. 13(11): 933-943 / Ridder R, et al., 1995, *Biotechnology (N Y)*,
- 10 13(3):255-260 / Rossi et al., *Pharmacol. Ther.* 50:245-254, (1991) / Roth J.A. et al., 1996, *Nature Medicine*, 2(9):985-991 / Rougeot, C. et al., *Eur. J. Biochem.* 219 (3): 765-773, 1994 / Roux et al., 1989, *Proc. Natl Acad. Sci. USA*, 86 : 9079 – 9083. / Sambrook, et al. 1989. *Molecular cloning: a laboratory manual*. 2ed. Cold Spring Harbor Laboratory, Cold spring Harbor, New York. / Samson M, et al. (1996) *Nature*, 382(6593):722-725. / Samulski et al., 1989, *J. Virol.*, 63 :
- 15 3822-3828. / Sanchez-Pescador R., 1988, *J. Clin. Microbiol.*, 26(10):1934-1938 / Sandou et al., 1994, *Science*, 265 : 1875-1878. / Schena et al. *Science* 270:467-470, 1995 / Schwartz and Dayhoff, eds., 1978, *Matrices for Detecting Distance Relationships: Atlas of Protein Sequence and Structure*, Washington: National Biomedical Research Foundation / Sczakiel G. et al., 1995, *Trends Microbiol.*, 1995, 3(6):213-217 / Sheffield, V.C. et al, *Proc. Natl. Acad. Sci. U.S.A* 1991;
- 20 49:699-706 / Shoemaker DD, et al., *Nat Genet* 1996;14(4):450-456 / Smith et al., 1983, *Mol. Cell. Biol.*, 3:2156-2165. / Sosnowski RG, et al., *Proc Natl Acad Sci U S A* 1997;94:1119-1123 / Steinberg G.D., et al. Family history and the risk of prostate cancer, *The prostate* 1990;17,337-347. / Stryer, L., *Biochemistry*, 4th edition, 1995 / Syvanen AC, et al., 1994, *Hum Mutat.*, 3(3): 172-179. / Tacson et al., 1996, *Nature Medicine*, 2(8):888-892. / Thompson et al., 1994, *Nucleic*
- 25 *Acids Res.* 22(2):4673-4680 / Tyagi et al. (1998) *Nature Biotechnology*. 16:49-53. / Urdea M.S., 1988, *Nucleic Acids Research*, 11: 4937-4957 / Urdea MS et al., 1991, *Nucleic Acids Symp Ser.*, 24: 197-200. / Valadon P., et al., 1996, *J. Mol. Biol.*, Vol. 261:11-22. / Vaughan TJ, et al., 1996, *Nat Biotechnol.* 14(3): 309-314 / Vlasak R. et al., 1983, *Eur. J. Biochem.*, 135:123-126 / Wabiko et al., 1986, *DNA*, 5(4):305-314. / Walker et al. (1996) *Clin. Chem.* 42:9-13. /
- 30 Westerink M.A.J., 1995, *Proc. Natl. Acad. Sci.*, 92:4021-4025. / White, M.B. et al. (1992) *Genomics*. 12:301-306. / White, M.B. et al. (1997) *Genomics*. 12:301-306. / Wilson R. et al., 1994, *Nature*, 368(6466) : 32-38. / Zhang SD et al., 1996, *Genes and development*, 10 : 1108-1119.

## SEQUENCE LISTING FREE TEXT

- 35 The following free text appears in the accompanying Sequence Listing :  
5' regulatory region

5

polymorphic base

complement

3' regulatory region

deletion of

or

probe

homology with Genset 5' EST in ref

sequencing oligonucleotide PrimerPU

sequencing oligonucleotide PrimerRP

T05290-T05290

<110> Genset SA

<120> Nucleic acids encoding human TBC-1 protein and polymorphic markers thereof.

<130> D.18363

<150> US 60/095,653

<151> 1998-08-07

<160> 7

<170> Patent.pm

<210> 1

<211> 17590

<212> DNA

<213> Homo sapiens

<220>

<221> misc\_feature

<222> 1..2000

<223> 5' regulatory region

<220>

<221> exon

<222> 2001..2077

<223> exon 1

<220>

<221> exon

<222> 12292..12373

<223> exon 1b

<220>

<221> exon

<222> 12740..13249

<223> exon 2

<220>

<221> allele

<222> 9494

<223> 99-430-352 : polymorphic base A or G

0976241-062501  
T05290="T022460

```
<221> primer_bind
```

<223> 99-430.rp

```
<221> primer_bind
```

<223> 99-430.pu complement

```
<221> primer_bind
```

<223> 99-430-352.mis

```
<221> primer_bind
```

<223> 99-430-352.mis complement

```
<221> primer_bind
```

<223> 99-430-352.probe

```
<221> misc_feature
```

<223> n=a, g, c or t

aggacagtat	ctagcacaat	accccaaattc	gactaactcc	tccgtaaaga	atagctacca	60
ctattgtgag	agtttttaagt	caagctgtga	ataaaactct	tgggtccact	taaaaatacc	120
tcccctggat	gtaagcatcc	agggaaatca	gggaatgcca	taagacagcc	ctaattctaaa	180
agcctacaag	aagctcagtg	ggcttcaagg	aagacactgc	tcttgggtacg	atgaggaaac	240
ctggccctct	atttgccctcc	tgggccacag	taatattgat	aatagctgct	gcttttagtt	300
gaggaccatg	tacgtctgtg	tactgcact	ggccacttta	cttacacttt	cctgctttgt	360
cctcaciaaag	atcctgtaag	gtgtgtattg	gtcccattta	gcaggtaaga	caatgaagac	420
cagaggtcca	gcaccttgcc	taaaccacac	ctgctgggat	ttggattcaa	gtccaaccgt	480

acagctcaaa cgctcagcca cttccctaaa gtccaccccc agctacatta agtaaaaaaa 540  
tccagaaaga tgccacctgg gggctctggaa ctgcctcctc cgagcacccg gctctcccct 600  
ccctgcggac tcttctctgg agaggatgtg atgcttctta cttttctcag atccctctcc 660  
ccaccctgcg agtgacgttg cgcctctgtg cctgggtggga tagggatctg ggagcttcgc 720  
ctgttttttg cacactgcca tcccctagtc ttagggagcg agctctgtcc cgcttttcac 780  
atctccgcgt ctttccctgc actctacatc accgctggga atgtccccag acctgatcgg 840  
ggcatgcaca ctgggggtgtg cgtgtgcgtg tgggtgtgtgt tcttgccgct gtgccgggct 900  
cgccggggcag gaaaaagcgc ctaatccagg ctctgcgtca ctcccgcaat tgggttagaaa 960  
tggagtttcc tgggtgtttaa tcccgggagg gcaacttcgcc ttcgttgttt cccagagtcc 1020  
ctgattttcc tgcctcgcgt gccagcgccc catagggcat ccgtgcctca gtacacctct 1080  
tgccatcctc caaggacggg gagaaggggt aaggcggggg agagcaagggt ggcttggtcg 1140  
cccccgggcc ccgcccccca tgttgtgtgc agtttccacc acgtctgttt cggagggaga 1200  
agaggagggt gcagatgagg cgaggcgctc tcgggagcgc ggagagcggg caggcagtcg 1260  
cacctgctga gagccactca ggccgagcaa gcggcgggca gtgccacctg ctataaatag 1320  
gccgccaagg acaggggtgtg cgactgtaca tcccgccacg agggcctgca tcacgcgcgg 1380  
ggccccgcgc ccccggtctc ccagggaaac gctgtgcccc gatcctgcgc aggggtctgg 1440  
atggggcggc ggcccagta cttccccctc attcccccca cagacactgg ctgaggatgg 1500  
cccgcgggct tgggggcggg ggggtggcaag gaggggaggg aggccgcggc ggaccgcgag 1560  
tgcagcagct gttgctcgcg tgtgactcgc ccgtccgggc cgtgctgccc aggcacagtc 1620  
acacggcgca gtggggagga ggaggacacc gagtccccct cccagctccc cggggaccga 1680  
gtggggagat cccggctcct gtcttcccc cgcctccagc gcgctcgcgc aggtctggag 1740  
gaggaaacca gagccgcgcg cagacacctc ctcttctctc tctcttctt cctcctctc 1800  
ctcctcctcc tctcctctt cggtgctgc tcttgggtgc gccaccgtcc gccggtgct 1860  
gttctgctccg ccgcccggg acctgctgtg tctcagctg ggtggagaag aggcgggcgc 1920  
cgagccgagg ggagccccct ccccgctccc ccgcgggggg aagagcgag ccagccgggt 1980  
gcgatggact ccccgccccg ccaggccgtc cccaggatgc ccccaagcac ctgcgcgtcc 2040  
cggccccggc ccgggctctg agcgcgcgcg ggcacaggta aggcgcttcc tggggcttcg 2100  
tcttggccac cctgctggct cctctcgggg cgtcgcggcc gccccctccc gcagcacgcc 2160  
cctgccccgc ctggccgcgg aggggaaggc atctggccgc ccacggagcg gaggccaggg 2220  
tctctcgggg gaggaagttc attgcctct cgttgcccc cttaccccc ccccccgcc 2280  
gcccttgagc gaaagcgaac ccttaatgtt gctagcgacc cgagagctcc gccggcttct 2340  
cccccaacct ccgccagctc actggctcgc gcctctctcc cctccccct cccgccaatt 2400  
atcctagcgt gtttgcaagg cgaccagatt ggaaagagt tggtcagagt gaccccaagc 2460  
cacgctttaa aagttcaggg tactttgcag tagtaacttt ggcagctcca ccagtgcgcg 2520  
caacatttct ttctatgggt acatcctgta ccagtcattt tgaaacctg cttcattgtt 2580  
tctagccgct tctgatggc tctgtgatta tgagaccccc ctcaaacttc accaggcatt 2640  
aagggtttgt ttttgctttt ttttcagaga ggtatcattt cgtttgaaat ccacctagat 2700  
gtggcttttc ctgttttgat tttacttaac atagcttatt ctctggaagt tgctttaaaa 2760  
agaaattgaa agtgatgggt gtctcttcca ccaaacagtt taattttcag ggtgcctcat 2820  
attaatggat atgttttccc ttcatagatt tctcattgtt tcccttatga tgggatgatt 2880  
tcatttatta ataaaatcag actttgaaag agcattttaa aatgacctgg tttaaatagg 2940  
tcacacccaa gaaactcagc tatctgtaca agttcaaact tctaaacttt ttcaatgagc 3000

09631-0601  
T0330-T0340

taggggtggt ggcacccacc tgtagtccca gctacttggg aggctgaggc aggaggatca 3060  
 cttgagccca ggagttcgag gccatagtg gctatgactg tgccacctca ctggagcctg 3120  
 ggtgacaaag tgagatccca tctcttaaaa aaaaagagtt taggggacat tttctgaagt 3180  
 gaacacaagt agagcattct aacactattg agtgcaagga gacctggaag ggactaagtg 3240  
 gttcaaagca ggaaataaaa tcatcaggtg ataattaaaa taatttcttt cctgtggatt 3300  
 tgtccagcca tttgcaaacc aggagaatag gaaaaaaaat cactagtgtg gttataaatt 3360  
 attacattac gttttcaaag gaaaattttg caaatgcgtc tccttgtcat agtctattgt 3420  
 tatctacccc actgagagtg ctggggcttc cccttttcac cacgacagca tttctggttg 3480  
 ggtggcagtc atgcagtgtt gacctggtgt ccataaggc acagtgtgtc aaaacactag 3540  
 tgggtattag gaggaacgt gcaactctga agcaacagag cttgccctt cttctcatt 3600  
 atccagctgg tgataatccc tgtccccac ttccctagaa gacagctttg accaggaagg 3660  
 ctgcaatgac aatgagatgt accctatgc agagccagat gtgggcgggt ggcttttttg 3720  
 tgggtccagat cttctaggat cttctaggat gtaaccctgg caagcagtgg ggagcctgaa 3780  
 tcaagcagca tggctgttac ctcttctgtg ttcacagcag catcttcagt tgtcttgggtg 3840  
 cctggagcag gcaccacagc tgcctgctct gttggccacc agctttctag agtagatggt 3900  
 agggaggaga gcaaggggct caagaggatt ctgtctttga acatgctttt aantttgatc 3960  
 tgacagaatg gcagctccct gaagtccttc ctactcttc cacagcattt ctctgtagg 4020  
 cccagtttt tgctcttttc agattcccag aggacntgaa aatgtatcac ggcccatttg 4080  
 gggacttct gtatatgtgt gggtgctca ggatcatttg ttttgccctt ttccagtcta 4140  
 ccgtgctgcc cttctcaagt ttaatgnacc acgttagttt caatatttta tatatttctc 4200  
 agcagtttt atctcttgggt cattaaactt gagaagtaaa atctgctcat taaaatgact 4260  
 gagtccatgg ccaggcatgg tggctcatgc ctgtaatccc agcacttttg gagtccaagg 4320  
 cgggtggatc acttgaggtc aggagtctga gaccagcctg gccagcatgg caaaaccctg 4380  
 tctctacaaa aatatagatc tacaaaaact agccaggcat ggtggcatgt gctgtagtc 4440  
 ccagctattt gggaggctga gacaggagaa tcgcttgaag ccaggaggcg gaggttgag 4500  
 tgaaacatga tcgtgccact gagtccattc agcagcagag tagtggtggg gtttgtatcc 4560  
 ctgtagtgat gacgaaggat ttaggttttc agtcagaact gttaccttac aatttccttc 4620  
 actgactttt cttcctttcc aacaccacat tccaataaaa aatatcttta gaccagattc 4680  
 ttcacgaaag acatgaagg 4740  
 aaggagtctt gctgtgtcac ccaggctgga gtgcagtggc gtgatctcag ctactgcaa 4800  
 cctccgctc ctgggttcaa gtgattctcc tgcctcagcc tcccaagtag ctgggactac 4860  
 aggctgctc taccacggcc ggctaaattt tgtgttttta gtagaggcga ggtttcacca 4920  
 tcttgccag gctggctctg aactcccgac cttgtgatcc acccgcttg gctcccaaa 4980  
 gtgctgggat tacagggtgtg agccacggcg ccgaccagt ttttgacatt tctaagccaa 5040  
 aagttccatt tgatgaggtc ttagatgcag gggcaatgtg tcccttttca gatttcagat 5100  
 gtttagaaaa agatgtgtca tatttgggcc aactgaaaa ctcttgatat gtaggttttt 5160  
 atgaagctgt gcagaatgta ggaaatacat tttagaacca acaaagaggc atttaatttt 5220  
 gagtgtgcct gtctcctttg agatgagcaa cagctatttt tctcttcaa agacaatgcg 5280  
 tgtatttatc agcacatttt atataatcag caaatctaaa cctctgaatt aggtaagccc 5340  
 tataggtttg ttgccagaat tagtgaattt atacatgcaa agtgcttaga acagtgcctg 5400  
 gtacacagtg agcactcaat attatttatt gctattatta tgtttattta ttttatactt 5460  
 ttagagtata attttgatgt taggtttgga ttgctgaggc caagcaaat ttagatagac 5520



caaccagct aatccactag aaagatatatt gaggggttatt cccatctaaa gatctatggg 5580  
 atctttggat atcatctgtg agaaacaaca gaagtttgta gataagacag atatagattc 5640  
 aaatgccacc ttcacaattt attagtggag gaaccctttg gtaaaatgag catgacaaaa 5700  
 cctacttctc agctttgtga acgtacagaa gatcatgaat gtaaaatgtc catgaagtgg 5760  
 taggtgggtca acagatatgt ctctagcaaa gtgggttaaga gcaagcaaac tctggagcca 5820  
 aagggcctgg gttcaaattc cgattctgcc acttcttata gtatggcctt gggcaggtga 5880  
 cctcactttt ctgtgccact attcaatgat aatattcctt tattgtccaa cgttttgtaa 5940  
 gttaaatcag ttaataaaca cactatgata atgtgttggg aactattctt tttactttta 6000  
 gcagaataac ttaaaggaac actgcaggta ggaggggttat acataatctc tgagggccag 6060  
 ctaggacact cgccatttc ccaccctttt tctgtgcaa tgaagagtat aagaagtgc 6120  
 agtgccattc taaaggacta gccttgagtt ggctctaatt tatatgactc gtgcctgtaa 6180  
 tcccagcact ttgggaggcc gaggtggca gatcacnttg aggtcaggag ttggagacca 6240  
 gcctnggcaa acatgacgaa atctcatctc taccaaaaat acaaagatca gccgggcatg 6300  
 gtagcacaca cctgtaatcc cagcttctgg ggaggtgag gcagaagaat tgtttgaacc 6360  
 cagggggcgg aggatgcagt gagctgagat cagccactg cactacagcc tgggtgacag 6420  
 agtgagattc tgtctcaaat aataataata atttatatga gaaagaagtc attcaaaagc 6480  
 atcattactt tacatgtcaa attagaaagg cacaccccag tactaaagca tcttgatga 6540  
 tgaaaacatt tagaaccact ggtttcagga gctccatgca atgggtgaaac agcctctact 6600  
 ccaaggggtg ttgtccctg tgggattcct ggggtaggaa cactgctc ccgttgggggt 6660  
 ggaatcctgt ggaggaagtg gatgaagagt gtagccaagt cagtgagcct actgcatggg 6720  
 attagagtac ttcaggttgc agtataattc tgttcaggtg catgctcact ccactctggc 6780  
 taaagaacag agaaattaaa ccattgattc acagagcaat atgagtagct gcctggggac 6840  
 ttctgtcca ttctggggtt cccaacagcc aatcaatatt ggccgggttc taatctgacc 6900  
 tagagctaga ggccactagc acccctccat tcatccttct tctctccct acccactccc 6960  
 accagcattc tgaggaggtg agggctgaag ctgcagaggc tgttgactg tcagttactg 7020  
 taaagtcccc atcctgctct ctgtagtttg ctgtgaagga gtggaggggg ctgggaacaa 7080  
 aggggggttc caataagag cttactaata cctacccttg cttctctcac ttcctgatca 7140  
 ataattccca tctttcttc atgtacctcc cccacatttt tgttctttaa ggaagaagg 7200  
 gagcagcttt ccatccccac acaatgttgg ggacatttgg tattatacat tatgaaaata 7260  
 agatttacca gatttttagag atggaagaaa acttggggat gatcttggtc cattctctta 7320  
 taagaacaaa caatatgtg agaagctgag taacttgctt tttcaacttc acacttgaaa 7380  
 gtgttttcat gaggaagttg gggttctctg cagcacttgg atgggagtca gggacttgga 7440  
 ttgtcccaat tctgtcacta aatttggaag agccacttaa cttttccaga atctagttgc 7500  
 ctcatctcaa atattaggga aatttctaaa tggcttaaaa ggagcttggt agctttaaaa 7560  
 ttttatgatt ctaagtgtat gctgccagag atatgtagca tagcaggaca cattaacaag 7620  
 attattgaaa ctgttctaata aaaggacatc tttgtgtctt gggtagctac tatgtttaaa 7680  
 gactgtgcta ggtgggagtt gtgcagaata cacaggttgg ctgtagagg atagggcgtg 7740  
 tacacagaca actctactaa caagaacgtt actagaagct tattggaatc acagtatttc 7800  
 ttgctgaggg tatgaaacat aagagttctc cttggaatat gaggttctat ttggggctta 7860  
 aagaatggtc aaaggttgag tgcaataaac atggattgag atggctttaa aaaataatca 7920  
 aatgggttgt tagtattaaa ctggtgcaga aataattgca gtttttgcca ttccttttaa 7980  
 tggcaaaaat tgcagttact tttaaaccaa atccctaata ttatttgcac agtttatctc 8040

tgttatggaa gtttttattg acaagtaatg tagatattca cctgatctaa gttaccctga 8100  
 atcttatatt agcagaatct gaattgctta taaataatta tggctatgtt ggatgtagaa 8160  
 cttattattt gatagtttat gaacagtgtt aagggtcta ctacttttta cagagaagct 8220  
 aagaacatgc tacagctggt tgaaaaacaa aaacttcagg cattgaaatg ttttgtcaat 8280  
 gaaatggcag gactcatttg atgactgatt attatcaact gatttaaattg actgaatttt 8340  
 tgggtactgtg tacatctata ctctaagaag gaaattgaaa gtaattctgc tatgcttggt 8400  
 gccactatat taataactgc atcatctaaa ataattgata gagctcagat ttatcctttg 8460  
 taataattct agtacttctt taaacatgtt ttgggattag cagctgtcaa cagttagaac 8520  
 atgaaacaga ttctgttaca ggagtagaag tcgatccaga catttaattgt ctttttcacc 8580  
 tgtgagagag agaataaaga gaaagagaga tcattattta tgggattatg tgaacttcaa 8640  
 gtccgttttc attattagga gaagctgtgc tttaaaggac agtcagggac tttactttca 8700  
 tgaaatgcct gagctgtaaa taaagtattg ctttattttt tatttcttga acatttgaaa 8760  
 taaaaaatta gctatgagtt atgttcaaat tatattataa aaatttgctc ttagcattgt 8820  
 gcatatatat tatacagaaa aacacagagt aaaaagaata gacttcagtt cctgttcaga 8880  
 aaagggttaa aatttgaata ctgatttttg aaaccccaa ccttaagaat tcaagaagct 8940  
 tacggtcttc ttgagggaca cctattcaaa ctcttaata tggtgattgg gtagaaagtg 9000  
 cagaaaagcc tgctgataca tgccctaaaa caccttgga aaaagaggtg gtagttgctt 9060  
 gaggtaggac ttaagtacta gttggaaata gaagacaagg atggagactg ttggtagatg 9120  
 actctccatg ggtccttctt gtttctacac accttgtaag cagggcattg agtgcctgtg 9180  
 ttccaaacta ccttttccat catgtttcta cagcaaacag tcatggaaga tagaaataga 9240  
 gtcttctctt ggagcaaagg gcagacacgc ttgcttctctg tacttccac tataagatat 9300  
 tccggctccc taaactcagc tgcttttctt gtaaccacc atgatacaga tgtcacctga 9360  
 cctgtgggaa ttgggggtca gggaaccaag agaaatgctg actgtctggc tactgtgact 9420  
 gccctgagta ataaattgtc cttcgtctcc aaccaggag tctcatgttt tctaccagca 9480  
 ggataactgt ggcrggctaa cgtgttagtt tgcaagtaag gtaaaatctc agaccctttg 9540  
 cagtttgttg cagggattat attctgagga gagaggaacc gtatgcacca tggtcagag 9600  
 gcatgagaaa cggggaacaa taactagttc tctatcttca gagcctttaa aaggtgcacc 9660  
 aaggagggca ttttagggga gaatataaag ttggagatat agacacagcc agattcctga 9720  
 gagaccttat atgccaggta gaagacttca gattgtatgg gggaattatt agagaatttt 9780  
 tagcaggggt gtgatatgat aaattttgtg ttgattaagt tactccagga aatatgcgat 9840  
 ggggtgattg aaggatgggg caccttttct ctaggacgaa aaagaaagag tagttggtga 9900  
 agtcagttag aggtagtaat aggatgaaga agggatctga atgaccctt ggccattcag 9960  
 tgagtagtga tgctattcac ctagatacag cacatagnng ggaaangaaa tinctgggaag 10020  
 gagggagatg agaccgagtt agcttttaaa taactaaatt caggcctagg agcctatagg 10080  
 ctatccagat agaaatattt aatngcctat atggatctgg aactcaggaa ggaggcttcc 10140  
 gtgggagcag aacacttggg caccattagg gtgtatgtgg tagatgcatt cttgtgcagc 10200  
 agtcaagggg atgggattta gactcaagtg caaattgccc cccatctcct gtgataagtg 10260  
 actgaagctc tccgggcttc agtttcttag ttcatcatag tgggctctag cggataaatg 10320  
 ttacaaaggt taaatgagac aacataggca aagtgcgtgg tactcaatag aagtcagctg 10380  
 ctgtcatcag cagcaggatc accagaatgt ggtgcttgac accaaaagat taggtgagat 10440  
 tgcccaaac agcaggtgaa atgaggggag aggatgnaag tcaaacacag gaagaaaagc 10500  
 ctttgaagta tgtggaaaga aacaaccaga aaggtaagat aagaaccaga agagattcaa 10560

gaaggaaggt gtggccgggc gcggtggctc aagcctgtaa tcccagcact tugggaggcc 10620  
 gaggcgggcg gaacacgagg tcaggagatc gagaccatcc tggctaacac ggtgaaaccc 10680  
 cgtctgtact aaagatacaa aagaattagc cgggcgcggt ggcaggcgcc tgtagtccca 10740  
 gctactcggg aggctgacgc gggagaatgg cgcgaacccg ggaggcggag cttgcagtga 10800  
 gccgagatcg cgccactgca cntcnagcct gggcgacaga gcgaggagcc gtctcaaaaa 10860  
 aaaaagaaaa aaaaaaaaaa gtaaggaagg tgtggccaag attgagaaat tcgtcagagc 10920  
 aaacaaggca gtcaggggct aaatagcctc ctttaaattt tacaaccttg aggacctcg 10980  
 caactttaac agaatttcag tggatcccta gggcaaacca ggccttatac accaggaatg 11040  
 gatggtcaat aggaagtggg gacagtaagt gtagacctta ccttgaggag aaggnaagag 11100  
 aaagagccat ggccaaggga agtttgaaat caaaggaaat atcttttttt ttttttttcg 11160  
 attggagaga cctcagttat tcttttaaaa tacttattga gccctcagt tattctttta 11220  
 aaatacgtat tgagtcccta ctttgagtca ggcacnatgg cagacacgag ggngatagca 11280  
 gtgaatcaga cagatgcaac gcctgccttc atggagtttc accttagcat ctgtccatat 11340  
 gctaggggag tggggcaggg gcaggagct ggatacagga gagactgaag atccagggag 11400  
 caagtgagta aagaataggg cttgagatcc cacagacaac tcagctttga acaaaagggt 11460  
 tttgtcatcc aataggacaa gaaggcgta ggatacatca aacgtggttg ttgaaaacag 11520  
 aaaagggtcg ggcactgtgg ctcatgccta taatcccagc actttgggag gccaagggtg 11580  
 gcagatcact tgaggccagg agttcgagac cagcctggcc aacatggtga aaccccatct 11640  
 ctactaaaaa taaaaaatt agccagggtg ggtggtgcat gcctgtaatc ccagctactt 11700  
 ggaaggctga ggcaggagaa ttgcttgaac ccagggggtg gaggttgag tgagccacga 11760  
 tcgtgccact gcactccagc ccgggcaaca ggcgagact ctgtctcaa aaaaaaaaaa 11820  
 ggaagaaaga acatagacag ggaaatgtag ttaagggnag tttgggttg ggtttggtag 11880  
 aagcgttttc tgttnnttgt ttgtttgttt tcagaaagag tctactctg ttgtccagac 11940  
 tggagtgcag tggcacaatc ttggcttgct gcagcctctg cctcctggat tcaagcaatt 12000  
 ctctgcctc agcctcctga gtagctggga ttacagacac ctaccaccac accaggctaa 12060  
 tttttgtatt tttagtagag acgggggttc accatgttgg ccaggctggt ctcaaactcc 12120  
 tgacctcagg tgatccacct atcttggcct ctcaaagtgc tgggattaca ggtgtgagcc 12180  
 actgcacctg gcctaacatt gatatctgtt gatgagaaga agccagggtg tggagtgata 12240  
 gcttatagca catgaactga ataaaacagt gtttaagaca atgtttgcaa cataataggc 12300  
 actgaagaca tgttaatgga aggtggattt gtgattcaga acctctagac tacctgggcg 12360  
 agtcttttaa aatgtaagta atatcttaag tgatattact tgtccagat cagttgttta 12420  
 aaactgaggt ttaatgctgt cagagtagca ctgtatcgtc ttctatcatg ggggcctttg 12480  
 ttggctttag gaggtttgtg tttcatagta gtttccaggt gggctctttg ttacctgtaa 12540  
 tgagtgtgac agttatgcca taaccagggt ttatatggaa tacaattttg agaaagttct 12600  
 ttctaggcag agaagcttat ttgaacctct tattatattt gggtttcagg cttttgagtt 12660  
 cttctgaaat aatagccctt tgaaggtagc tattgctatg acttcattaa attctaattg 12720  
 ctctggtttt ctccccagg tttctgcata tgaagtgtgt aaaatagatt gcttgatcca 12780  
 aaacagaaaa acagtgataa ctgttttgct gagttccag acccttccca agatggaacc 12840  
 aataacattc acagcaagga aacatctgct ttctaacgag gtctcgggtg attttggcct 12900  
 gcagctgggtg ggctccctgc ctgtgcattc cctgaccacc atgcccatgc tgccctgggt 12960  
 tgtggctgag gtgcgaagac tcagcaggca gtccaccaga aaggaaacct taaccaagca 13020  
 agtccggctt tgcgtttcac cctctggact gagatgtgaa cctgagccag ggagaagtca 13080

acagtgggat cccctgatct attccagcat ctttgagtgc aagcctcagc gtgttcacaa 13140  
actgattcac aacagtcattg acccaagtta ctttgcttgt ctgattaagg aagacgctgt 13200  
ccaccggcag agtatctgct atgtgttcaa agccgatgat caaacaaaag taagtggat 13260  
ggagatccaa aagactaagg tgtggctggc tggtttttat tgtatggggg tcaggatatt 13320  
tattttaagt atactgaaat gaataaggaa ttaatgctgc agttataaat tgattactta 13380  
gctgaatttt tgttttatgg tgatagttaa tagtttttaa gcacatttga aaacagatac 13440  
gagaaattat cagtttttga gttcaaaaat tcaagagaaa tcagtctaaa actactaatt 13500  
aagagcagaa gtgttaagat gtacattatt tcagatgaat gttctaaagc catgcctctc 13560  
aaactgaaat gagcttgtga gtcacctggg gatcttgtta aaatgtgaat cttgattcag 13620  
taggtctggg gtggaccca agactgcatt tgtaacaagc tgccaagaaa tgctgatgct 13680  
gcccttttgc aggttgcaat ttgagtggca aagttctaaa tctccacatt tgtaatccta 13740  
ttaagaaaaa tatagtcatt cgtaaactgt gtaaaaatgc tactggccag tttcccaagg 13800  
cataatgttc acttaggcaa aggtcattga taagaacgct ggatatgcat ctaagttttg 13860  
atgcgatcag gggttctttg tgttttttct tttcgcaaac ctcaggctag atctgattag 13920  
cttgttatta tcacatgata tggctgaaaa aaaatgtgag acatggtaaa agttctgctc 13980  
tttctcgtt catttgtgct tgctttgtta ttagcattcg ttgtagctct gggcaggact 14040  
catttgaaga tgcttgnccc attttatgag gattagctta gataaaattg aaaatataat 14100  
gcaaatagca actttctcag ttgggctcag ggctccacag ctaaccccat ggactgtgga 14160  
gtcttgccgt tgttttgggt gccaaagcaag ccaagtcaca tgtgattcaa gctgtctgcc 14220  
acatgtacag ggcgaggatg cgagtgtcaa tccacctgtt aactgtcagt gaagccttga 14280  
aagcttctca tattttcaag gttaaaatct ggatagaaat gctaaagttt tctctctgca 14340  
ctccattagg ttattttatg tactctctag ggtgtaagga ccttatttag aaattaatat 14400  
tcttggtatc aagtagatgc ctttttgctt gttcatttgt tggttcttct agtcattcag 14460  
aattgctgtt gcaggctactg ttggagatga tattagcaga ggcttgtagg aaggcaggag 14520  
catcagtggg gaataggacc aggtgatcta tgtataggac ataatggaag gactgagaag 14580  
ggagcctaac acacacccaa agggtagaga aggctttgtg aaataaaggc taatatggag 14640  
ctcaaaaacca ccatttcaat cacagaatca aactctcata ttataaatca tttcatgtta 14700  
ttgtccacac atctcaagtg ggcacggcag catcaggctt ggagattcag agggactaac 14760  
ttcctgtact ctaatcctac ttctgcaccc ataaactggg tggcctcagg caattgagtc 14820  
tgttttctta tctgttaaatt ggggataatt acagtattta tccaatagag ttgctggaaa 14880  
gactaaatga ggtagcactc gacctgaaac ttagtaagca tttatagcca taaaaacatt 14940  
ttcattcaag aaaattttac tagaggcaga ttatatgcta atttcatttc acgtcttagg 15000  
taaaaagaaa catgatacct agatgagtgc cttcagcttt caaagatgag attctgggtca 15060  
tatttgagga acatttttaa aactacacgt ataacttaat ggctcctatt atttgacaa 15120  
attccagaat gaaaatgaga ggactgaaca gcctgtacct cagtccagct ctatatagta 15180  
tttgactga atttccttgg ggagagtttg tgcgtggaat cggtgttcag cattttacac 15240  
atttgactct tccccaaat cttttacggc catctgagaa taggcttctg gccagtcatt 15300  
cggatgcctg acaagagaaa gagatttata accaaattct gtaattggga cttccagtct 15360  
ttccccaggt agagaattgg acttactcta tatgctaaaa acccatgggt gaaatatgaa 15420  
ttagttctta agtgattttt ggcttgcata ccatttttgc aaacacaaat tgctattact 15480  
ctgctcattt aataaaagaa taattttag tataggtata tacctcaatc agtgattttg 15540  
ttggtggaaa cagaacagta aatcacactg gccatgatgc taacagcgtg atagattttc 15600

tgttcttggg acaccaatgt cactgtatct catagcgaag gattatctgc tgtaggagca 15660  
 ttctcttgac tacttataac atttgctggg tgaaataatt ctccagggtta aggcctcttc 15720  
 taaacagatg aggtcagcac taactgcatt tgccagagaa gacatatgca tttactgccca 15780  
 gcatcataaa cacaaaacta cagtttgca ggaaccctt tgaccagcat ctaattaatt 15840  
 cactgagtaa tgtcttggga gaagaggcat gtaaaggaac aattttataa gcatgccatg 15900  
 agattgtttt cnnattgtat gttccataga atatgaggaa acttcaaac attttgtgga 15960  
 aaaattgaat taaaaagtaa aaaacacata tatacataag ctttatttct caagataaac 16020  
 tttatcaagt tcaagacact tttgtaagca atgttaacag ccattgagtc ggtctctaaa 16080  
 gaactgaggg tcctgggaat ttaaccatgt ttatacagtc ttttatacat tattaactgg 16140  
 agaaaaattg gcgctcttta aagattttttt aaaattgaga agcaaaagga cgtcagaagg 16200  
 agccaaatta ggctgnnaa gtggatgctt aatgatttcc catggaaact cttgcaaat 16260  
 tgctctgtt tgatgagagg aatgagcagg aacattgtca tgggtggaaa ggactctggt 16320  
 gaagctttnn caggcgattt tctgctaaag ctttggttaa ctttctcaaa acactctcat 16380  
 gataaacaga tgttatcatt ctttggcctt ccagaaagtc aacaaacaaa atgccttggg 16440  
 catcccaaaa aactattgca accatttgc cttgaccagt ccactttcgc tttgactgga 16500  
 ccacttctgc tctcagtagc cattgcttaa atttgtcttg atcttttagga ttgcgtggt 16560  
 aaaactatgt ttcatacct gttacaattc tttgaagaaa tgcttcagga tcttgatccc 16620  
 acccgtttaa aatttccatt agaaactctg ctcttgtctg cagctgatct gagggcaatg 16680  
 gttttggcac ccatctagta aaacgtttgc tcagtgttaa tttttcatcc aggattgtgt 16740  
 aagctgaacc agcagagatg tctatgatat tggctagtgt gtcctcttca atgagggcat 16800  
 gaacaagatg aatattttcc tcaacaatt atctggatgg tctgctgctg caggcttcat 16860  
 cttcaatatt gtctcgcccc ttctttttct tttccccccc gcttgagaca cagtctgtt 16920  
 ctgttgccca ggttgagtg cagtggcccg atttcggctc actgcaacct ctgcctcccg 16980  
 ggctcaagcg attctcctgc ctcagccac caagtagctg ggattacagg tacacatgat 17040  
 cgtgcctggc taatttttgt attttttagta gagacagggt ttcaccgtgt tggccaggct 17100  
 ggtctcgaac tcctgacctc aagtaatcca cctgccttgg cctcccaaag tgctgggatt 17160  
 ataaacatga gccaccaaac ctggcctcat ctttcttaa aatgagttat acatttgtaa 17220  
 gctgctgatt tctttggaca ttgtgcctat aaactttttg taaagcatca gtgatttcac 17280  
 cattcttcca cccaaacttc accataagtt tgatgtttct tcttgctttg attttagcag 17340  
 gattcatggt tctctgatag ggggtctttt caaactgatg tcttatectt cttagagcct 17400  
 catccagat cctgttcaga catgctacaa gttaatacaa gtttatttgg tgccaaaaaa 17460  
 tggaaatcca tgcatagttt ttaaataata tgcatttttc atgnactttt tgaagacccc 17520  
 ttgtatactt aaactgctcc acatggaaaa gcttccatga tcaaatgcag taaggcagca 17580  
 tctcaaacat 17590

&lt;210&gt; 2

&lt;211&gt; 99960

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; exon

T05290.F00060

<222> 4661..4789

<223> exon A

<220>

<221> exon

<222> 6116..6202

<223> exon B

<220>

<221> exon

<222> 9919..10199

<223> exon C

<220>

<221> exon

<222> 14521..14660

<223> exon D

<220>

<221> exon

<222> 50257..50442

<223> exon E

<220>

<221> exon

<222> 56256..56417

<223> exon F

<220>

<221> exon

<222> 63326..63484

<223> exon G

<220>

<221> exon

<222> 76036..76280

<223> exon H

<220>

<221> exon

<222> 78364..78523

<223> exon I

096231.06504  
T05250.TE29460

<220>

<221> exon

<222> 85295..85464

<223> exon J

<220>

<221> exon

<222> 93417..93590

<223> exon K

<220>

<221> exon

<222> 97476..97960

<223> exon L

<220>

<221> misc\_feature

<222> 97961..99960

<223> 3' regulatory region

<220>

<221> allele

<222> 1443

<223> 99-20508-456 : polymorphic base C or T

<220>

<221> allele

<222> 5247

<223> 99-20469-213 : polymorphic base C or T

<220>

<221> allele

<222> 6223

<223> 5-254-227 : polymorphic base A or G

<220>

<221> allele

<222> 14723

<223> 5-257-353 : polymorphic base C or T

<220>

092634-062501  
T0520-TE220

<221> allele

<222> 19186

<223> 99-20511-32 : polymorphic base C or T

<220>

<221> allele

<222> 18997

<223> 99-20511-221 : polymorphic base A or G

<220>

<221> allele

<222> 19891

<223> 99-20510-115 : deletion of TCT

<220>

<221> allele

<222> 29617

<223> 99-20504-90 : polymorphic base A or G

<220>

<221> allele

<222> 42519

<223> 99-20493-238 : polymorphic base A or C

<220>

<221> allele

<222> 69324

<223> 99-20499-221 : polymorphic base A or G

<220>

<221> allele

<222> 69181

<223> 99-20499-364 : polymorphic base A or T

<220>

<221> allele

<222> 69146

<223> 99-20499-399 : polymorphic base A or G

<220>

<221> allele

<222> 76458



<223> 99-20473-138 : deletion of TAACA

<220>

<221> allele

<222> 78595

<223> 5-249-304 : polymorphic base A or G

<220>

<221> allele

<222> 82159

<223> 99-20485-269 : polymorphic base A or G

<220>

<221> allele

<222> 84522

<223> 99-20481-131 : polymorphic base G or C

<220>

<221> allele

<222> 84810

<223> 99-20481-419 : polymorphic base A or T

<220>

<221> allele

<222> 89967

<223> 99-20480-233 : polymorphic base A or G

<220>

<221> primer\_bind

<222> 988..1006

<223> 99-20508.pu

<220>

<221> primer\_bind

<222> 1509..1529

<223> 99-20508.rp complement

<220>

<221> primer\_bind

<222> 5039..5056

<223> 99-20469.pu

005250-1129460

```
<221> primer_bind
```

<223>..99-20469.rp.complement

```
<221> primer_bind
```

<223> 5-254.pu

```
<221> primer_bind
```

<223> 5-254.rp complement

```
<221> primer_bind
```

<223> 5-257.pu

```
<221> primer_bind
```

<223> 5-257.rp complement

```
<221> primer_bind
```

<223> 99-20511.rp

```
<221> primer_bind
```

<223> 99-20511.pu complement

```
<221> primer_bind
```

<223> 99-20510.rp

```
<221> primer_bind
```

<222> 19986..20005  
<223> 99-20510.pu complement

<220>  
<221> primer\_bind  
<222> 29529..29547  
<223> 99-20504.pu

<220>  
<221> primer\_bind  
<222> 30041..30061  
<223> 99-20504.rp complement

<220>  
<221> primer\_bind  
<222> 42268..42287  
<223> 99-20493.rp

<220>  
<221> primer\_bind  
<222> 42732..42752  
<223> 99-20493.pu complement

<220>  
<221> primer\_bind  
<222> 69026..69046  
<223> 99-20499.rp

<220>  
<221> primer\_bind  
<222> 69525..69543  
<223> 99-20499.pu complement

<220>  
<221> primer\_bind  
<222> 76323..76343  
<223> 99-20473.pu

<220>  
<221> primer\_bind  
<222> 76771..76790  
<223> 99-20473.rp complement

095231-062501  
T05290-TE260



<221> primer\_bind  
<222> 9475..9493  
<223> 99-430-352.mis

<220>  
<221> primer\_bind  
<222> 9495..9513  
<223> 99-430-352.mis complement

<220>  
<221> primer\_bind  
<222> 1431..1455  
<223> 99-20508-456.probe

<220>  
<221> primer\_bind  
<222> 5235..5259  
<223> 99-20469-213.probe

<220>  
<221> primer\_bind  
<222> 6211..6235  
<223> 5-254-227.probe

<220>  
<221> primer\_bind  
<222> 14711..14735  
<223> 5-257-353.probe

<220>  
<221> primer\_bind  
<222> 19174..19198  
<223> 99-20511-32.probe

<220>  
<221> primer\_bind  
<222> 18985..19009  
<223> 99-20511-221.probe

<220>  
<221> primer\_bind  
<222> 29605..29629

09590 "T.E.29260

<223> 99-20504-90.probe

<220>

<221> primer\_bind

<222> 42507..42531

<223> 99-20493-238.probe

<220>

<221> primer\_bind

<222> 69312..69336

<223> 99-20499-221.probe

<220>

<221> primer\_bind

<222> 69169..69193

<223> 99-20499-364.probe

<220>

<221> primer\_bind

<222> 69134..69158

<223> 99-20499-399.probe

<220>

<221> primer\_bind

<222> 78583..78607

<223> 5-249-304.probe

<220>

<221> primer\_bind

<222> 82147..82171

<223> 99-20485-269.probe

<220>

<221> primer\_bind

<222> 84510..84534

<223> 99-20481-131.probe

<220>

<221> primer\_bind

<222> 84798..84822

<223> 99-20481-419.probe

005290 "T E 3 9 / 0 0"

<221> primer bind

<222> 89955..89979

<223> 99-20480-233.probe

**<220>**

<221> misc feature

<222> 3698,12593,13035,21712,27644,27655,31143,43084,43129,64585,66950  
67301..67302,67926,75425,98821..98822

<223> n=a, g, c or t

<400> 2

ctcaagcttg	aatacttgaa	tccaaacttt	catgcttaga	gtttacccca	tctgttgaag	60
gatgtgcaat	ataatgactg	caatagaatt	cactgtggag	cctccaaatt	agaaattatt	120
gtctgtgagg	gccaggcacg	gtggctcacg	cctgtaatcc	tagcactttg	ggaggctgag	180
atgggaggat	tgtttgaggc	caggagtttg	agaccagctt	ggtcaatata	gcgagacccc	240
catctctgtt	tttttttttt	aaagaaatta	ttgtctaaga	accagtgtca	tcttccaagg	300
agaaacttct	agatacttgt	tttaagataa	ataagaaaca	agtcattttct	aaatgtgaat	360
tatttttttaa	atgcaatttt	ttaaacattt	tattttaatt	atggcaatag	acgtggaaaa	420
gactcttttt	tgatagtagg	ggagagcaga	agaaacattg	aattaagtac	acagagattc	480
ttcagacctg	ctttaaaaac	acatgcatac	aaatgcactt	ctgtctctta	ggatctacta	540
actgatgctg	cttgcttttag	tcttttagct	aatattttct	ttctttcttt	ctttcttttt	600
tgttggagac	agagtctcgc	tctgtcgcca	ggctagagtg	cagcggcaca	atcttggtct	660
actgcaacct	ccgcctcccg	ggttcaagcg	attctcctgc	ctcagcctcc	tgagtagctg	720
ggactatagg	cgtgcgccac	cacgcccagc	taatttttgt	atatttagta	gagacgggggt	780
ttcactgtgt	tggatgggat	gttctccgtc	tcttgtcctc	gtgatccgcc	tgccctggcc	840
tcctaaagtg	ctgggattac	aggcgtgagc	cactgcgcct	ggctcatatt	ttctttatat	900
atcaaaaaca	ttcagcttgc	ttcactttta	tgaaagcttt	attatgagtt	tgaaagcaat	960
tctgcatttt	cttaacattg	taactggtgt	tgagttgaag	gcaggcccct	gggagcccct	1020
tgtgggcaat	tcccttcact	ctggaggctg	cctcgagcct	ggacaggcac	ttacacttgg	1080
tcagtgattg	cacagaaccg	gttgcaacag	attctgtgca	cctccctgtg	gcgcgtagca	1140
tttagcaggc	acttggtcac	tatttgctga	gtgagtctgt	taccttaggc	gtgtatttcc	1200
cgtggacctg	cctggggatc	attgctcatt	cactcatttt	gaacaagcca	atattacatg	1260
tccagggtac	gctctatagt	gtgaaacaca	aaggtaaattg	atagttcccc	ttctcaaagg	1320
aattttctaag	gtagtagcca	ttcttttgat	gcatattctc	attctcatag	agagtccaat	1380
tatggataat	tggacaaaagc	tgaatgtcgc	ttttatgaga	atccattctt	tctcttttat	1440
gcyttgaaaa	atgtgtagca	ttcattagtg	aattaggatt	tcattattca	agaagacat	1500
aaggctcttcg	aacagcagat	gactgaataa	aataatacct	aacagcagta	gaatgagggg	1560
aggacatatt	caaggaacat	tttatgccca	ttagattggc	agaaattttt	aaaaagtgac	1620
aataccgtat	aaagggtgaac	tttcctatac	tgatactggg	aacatgaatt	tgtaccattc	1680
agggaagaga	aacttgataa	tatctggtgt	agtctgaagg	ggcacagtcc	ctgtgaccca	1740
gtgaggacat	tcctcattat	ttcccttgcc	aaacatttca	catgagtcta	taaggagctc	1800

[illegible]

tatataagag	aggctcactgc	agcctccttt	gtaagagcaa	gaaaaaaaaag	caaataagtg	1860
tttaacaata	ggaacataga	taaattaggt	tatgcagtga	atatttgcac	tctgactaaa	1920
gtgagtgaat	caaaaaaaaaat	ttgtcaacag	gaataaatct	caaaaaataat	attgaaagaa	1980
gaaagctaata	ttacagaagg	atgtgtacag	tatgacacca	ttcatttagt	ttcaactaca	2040
tatctttttat	ggacacatac	atataaaaagc	agaaaacatg	aattgatagg	ataaacacca	2100
aatattttctg	catatggcca	ggtgtgggga	agtagtggtg	attaagcttc	aaagatgtct	2160
gcagtgggttc	ccattaaaag	tagaaaagtag	gctgggcaca	gtggctcacg	cctgtaatcc	2220
cagcactttg	ggaggccaag	gcagggtggat	catttaaggc	caggagtctg	agaacagcct	2280
ggtcaacatg	gcgaaacccc	atctctacaa	aaaaaataca	aaaattagcc	agatgtgggtg	2340
gcgcacactt	gtagtcccag	ctactcggga	ggctgaggca	tgagaatcac	ttgagcccag	2400
gaggtagagg	ttgcagtaag	ccaagatcgt	accactgcac	tccagcctgg	gtgacagagt	2460
gagactccat	cccaaaaaac	aagcaaacaa	aaaaagctca	tagagtaggt	aatagtcagt	2520
atatctgatg	tttttttgatt	gtctggttta	cattttttat	ttttattttt	tgagacaagt	2580
ctcacgtgt	cacccaagct	ggagtgcggt	ggtgcgatgt	cagctcactg	caatctctgc	2640
ctcctgggtt	cgagcgattc	tcctgcctca	gcctcccaag	tagctgggat	tacaggcgtg	2700
caccaccaca	cctggctaata	ttttatatatt	ttaatagaga	cagggtttca	ccatgttggc	2760
caggctggtc	tcgaactcct	gacctcaagt	gattcatctg	cctcagcctc	ccaaagttct	2820
gggattacag	gcatcagcca	ctgcacctgg	ccttggtata	tgtgttttaa	tttgtattca	2880
ttcattttaag	cctcatgaca	gctctgcgag	gaaagttcac	tatacgtctt	caggctgcag	2940
gtagaggacc	tgaaagggac	aggaggtaac	agtctggcca	agaccacaga	gccagggaat	3000
agcagaggaa	catttcacct	gggcattgca	ctccagagct	gggcttctca	ctgttctcaa	3060
cccctggcaa	atgctcactt	gaacaaagcc	aggtggtgat	acaaagggtat	ttgttatatt	3120
agtctctaca	cttttctgtg	tgcttgaaat	aactgcaaca	aagaatatat	cagtatttag	3180
agtaatgggg	gatttgcttg	tgtgtgtttg	tatttttgag	atggagtctc	gctctgtcgc	3240
ccaggctgga	gtgcagtagc	atgatcttgg	ctcactgcaa	cctccggctt	ctgagttcaa	3300
gcgattctcc	tgctcagcc	tcctgagtaa	ctgggattac	aggtgtgcgc	cactacaccc	3360
ggctgttttt	tgtattttta	gtagagacag	ggtttccccg	tgttgggccag	gctgatctca	3420
aactcccagc	ctcaggtggg	ccaccacact	tggcctccca	aagtgctgag	attacaggca	3480
tgagccactg	cgcttgccg	tttttttttc	taacaaaatt	attttctaac	agaaagcaat	3540
cagggtgagaa	tccacataag	aaacaattta	attcagagat	ttttgttgca	tattaaaaaa	3600
aaaatgtacc	ttcggtggg	tgtggtagct	cactcctgta	atcccagcac	tttgggaggc	3660
tgaggcaggt	agatcacttg	agctcaggag	tttgaganca	gcctggccaa	catggtgaaa	3720
ccccgtctct	acaaaaacta	caaaaaatta	gctgtgtgta	gtcccagcta	ctctgggggc	3780
cgagggagaa	ggattgcttg	aacctgggag	gtcaagactg	cagtgaacca	tgattgtggc	3840
cctgtactcc	agcctgggca	acaaagttag	accctggcac	cctgtctcaa	aaaaaaaaaaa	3900
aaagtacctc	cttgtaaata	agtaacacta	agacttcatt	tagtggttgt	caagcaaact	3960
ccattgtatt	tttattttca	gtttttatgg	ctagtagtta	agggagagaa	gcttggttgc	4020
agagaagaat	gaaaggatga	tgggaaaata	aaagtaggag	agaggaaaac	gcaagaaagc	4080
aagagatctg	tagaaaggga	tgaagggaatt	gtataggcag	agagaatagg	ttctttaatt	4140
gagaaattta	tgttgtctca	ccttctgaaa	tgcccccaaa	ggtaagttat	tgttttattt	4200
tgaaaagcta	atgatagcta	cctttctacc	acgctgtgtt	caatgtttta	cacactttac	4260
ctgtttgagt	ctcacaacac	agtgttatga	ttcgatcttg	ccattgggtct	cactttactg	4320

T 0 5 3 0 " T 2 3 4 6 0



aagaggaagt ttgaggctca gaaaagtaag aaactggccg aagaccacgg ttagtgaaga 4380  
cagatctctg atccagttgc agagtctgag caataaacta cttcaactga ttggtttcaa 4440  
agcacatttc gtcattttac ttggggtaat caaagcaact ctctgaggca aaattatttc 4500  
ctggacttgc agccatgtca ctaaggagca gatgagggtga gatcacagac aggatcagaa 4560  
tgatggcctg gtgccaaaaa gatgtgtcct agagattttt cattccttta agaagcagag 4620  
aagggagcga taaatgactt ttcgtttttc acttttttag acatcgaga tggcagcaga 4680  
gaatattgga agtgaattac caccagtgac cactcgattt aggctagata tgctgaaaaa 4740  
caaagcaaag agatctttta cagagtcttt agaaagtatt ttgtcccggt taagtagcat 4800  
aatttctcct gatttaagtt aaatcacttt ttaggagagt gtaagattga gttctatgct 4860  
tttattccat caatgttcat cataaaggta aaagtataaa accttttttt atgtttttctc 4920  
aggcttataa cagtattatc tacattttta attgttttta atttggccta ggtttaaaaa 4980  
aaatattcct tactcttttg tattatatcc aatgggattt ttttgccgct ccaaagaata 5040  
tttggttagcc agtccctata aagagcatgc attagataca ctgaagtgtg gcttctgttc 5100  
tccctactat cactatgtat aacttaaaaa acagttactg tcagctgctg gtgttagcta 5160  
tctaaaaggc tatatagtag gggtcagcaa actatgccc tgggccaat tctaccacc 5220  
tcctattttt gtaataaaag tttgtgygaa acaccgccac atccattcat tttccagtta 5280  
tctaaggctt cttttttgca gacttcagca gttgccaaa acactatatg cctcaciaag 5340  
cataagacac ttactatctg gccctttaca gaaaaagttt gccaaatata gctctataga 5400  
aagaacaaag tacacatgta catcaatctg ggagttcttt aagaaattat cctccctcc 5460  
catgagtgtg aatagcctga tggcacgtct gagaaatcaa atctgatttt cctcagagt 5520  
ttcacacctt tctggagtgt gcagtatctt attatagttc tttttgattt tatggcacac 5580  
ttcttttgaa acatctgatt ttattttatt ttttaattaa ggaaagttaa attttatttt 5640  
cttcgaagat gtttctgaga attttgcaat atcttctgag atcatgaaaa acagttgatt 5700  
tacaaaacca gagttgggag gggctgcatt tgagagctcc caaagggata gagtgctgtc 5760  
cgagtgcacat gcggctggcc gttatgatga cttgtgacct aggggaggga gttagttgct 5820  
gagtgggctt gagcacttga ttttcttat agacgaattg tcttgtcttc ctgcctatca 5880  
ctcatgccaa attacttagc caccaggtgt tttggaacgt ttaggttagt gtcttattta 5940  
ttttttaaaa aaatgatgga aatgttgatt attttaatgt acaaataatcc ttagtagcat 6000  
ttctcagtag ataacatttt tttcttgagc ttatttaaat ggaccaatct gcttctagct 6060  
gatgcctttg caaaagcctc cagagtcata actcgactgc cttttcttta tgtagggtaa 6120  
taaagccaga ggctgcagg aacactccat cagtgtggat ctggatagct ccctgtctag 6180  
tacattaagt aacaccagca aagtaagcac atttctcttt atcgacacc ctgaagaaac 6240  
caacaaatag gtcttgctca tctcctgtct acatactcc aatcataaaa cgtttgctgc 6300  
ttgcaaattt cttggcacag gtggaggact ggtcatgcag ttctatcata acataaaagt 6360  
tttacataaa agagcagatg gggctgggtg cagtggctca acgcctgtaa tctcagcact 6420  
ttgagaggcg gacgcggcg gatcatgagg tcaggagatc gagacgctcc tggctagcac 6480  
agtgaacccc cgtctctact aaaaatacaa aaaattaaaa attagccggg cgtggtggcg 6540  
ggcaccgta gtcccagcta ctcgaggagc tgaggcagga gaatggcatg aacctgggag 6600  
gcggagcttg cagtgagcca agatcacgcc actgcactcc agcctgtgtg acagagagag 6660  
actctgtgta aaaaaaaaaa agcagtagat tttcctatta aaaaaataat taatattggg 6720  
aaaacatcag aaagtggatt tgtgaattta gagaagtata cagcttaaat ttttcttttt 6780  
ttaagaaaat tttatttttg atttgggggt acatgtgcat gtttattacc tgggtatatt 6840

gcatactggt ggggattggg cttctagtgt acccatcacc caaatagtga acattgtacc 6900  
cagtaggtaa tttttcaacc ttcacacccc ctttcattct cccccacttg tggggaaatt 6960  
aaattttctga aactttatcc tgtagctggc tctatgatta taatgaaaca ttactgtttt 7020  
atttaataa gcaagtatct atgtccttct ttaataaact tgctttctag acatttaate 7080  
atatttaagc ctggtcagtt caactttata actcctgaaa agtggggttg ggttttgtgc 7140  
tagggaggcc agctttccct tctgctacca gaggactctc tttggcagta gtgaggagg 7200  
gagtgtttgt ggaggccagc tcttaccac aggcagggtt tacagtcctc tgccatccct 7260  
cctagacata tggtttcag aatttttcta acctacagta agaagcacat ttaacattgt 7320  
ggcgtagttc acaaacacac atacctacac attcacacac aaaattaaaa gtccacaaaa 7380  
caatatttac tgtgaacaac atacaataca tactgatatt ttgttctatt ttatttttaa 7440  
aatgctcatg gcaaaactact cagttgtacc acctactaac atgatagagg gaggcgttg 7500  
agaaacactt ccttagatgg atgagtgtt ctcaaatttc aggtgtccg cctcccgggt 7560  
tcaggccatt ctcttgctc agcctcctga gtagctgaga ctgggttaaag tgcagattct 7620  
ggttcagtag gcaggcgagg gggagccctg aaatgctgca tttctgacaa gctccaaggc 7680  
aatgtgtctg ctctgggtct gcagaccgcc tctggggagt gaggtcctag acagcagtct 7740  
tgtaaagtgt agtttctgag ttaaaatcca gggaacata gtgtcgtcca gcctccatct 7800  
aatacacact gatccacccc tgcaattcat tgcaagtgtg ggaaggctat ttgcttattt 7860  
gttgtgtaca gatgaaccac acaccgccc tttcatgtag gaagttacct aggaggagag 7920  
agatgacaga tacagaaaca gcccagcat caagcagagt gtggtaggag cccagaagtt 7980  
acaaataaga gacattggtg acttcagtgt cagaagagca aggggaaggg aagttagggt 8040  
tggtcagtgg aaccaggga aaggtgagg gtgagggggc agtgatgctt ttggactaaa 8100  
tcttggtgta ggaattgtgc ctatggaagt gaacagagaa gaaggcattc tagacagacc 8160  
agtgtcaata gacataccat gaagacattc atgtcactca gtgggctttc cagtaagcct 8220  
tattgcttgc tttttatttt tttccaaaag gcagatctag gaatatatac atattcattc 8280  
ttcaggactc gatagtgtg aagattcttt taaaaggatt taaaagtctg tctaagattg 8340  
caatttctag agtcattcta agagagatgc aacttttcag aagctgcttg tatgtattgt 8400  
atatgtttaa gtgtacttta catctttctt ttattcatct tgaattgaga aactactata 8460  
ttctatttta tgtaattgga tcccttctaa aaaattgatc acctaggagt tgcaaagaaa 8520  
ccaaatagcc ctgaaacttg acaaatgaaa atggcccttt cagttgtcca attaagctaa 8580  
gggttagctc tttgatatga tttggaggga tattagtaag aatttagatc aacagggttg 8640  
catgatggag attgtgttct gtgatgtatt gtcttagaga gactttttaa tccttaaaag 8700  
aatcttcaca actgttgagt cctgaagaat gaaaaacttc agttatgaaa gtaatcaata 8760  
tttcatagta tggtgggaat tttttcctaa ttcttataca attaaatgta tgtaacttct 8820  
ccctttggta aacacatttc tttttttttt tttttcaaat taaaaccctc aatacttgtt 8880  
acctaaaagg cactcaactg tgtaaatgaa caggtagaat tcagagtctc cagtccactg 8940  
ttagatgcat tcattcttgt ttactctatt cctgttgatt tattttttct cttccaacaa 9000  
tttcaatagg agcaagctgc tacaattcct ctttttgaat attttgaata tattaaaaat 9060  
atattggcca ctagccacgt cctgggtgca gtgttaaaca tcagtttgct tgagtggtag 9120  
tagttcattc ctttgaaaaa gcgtgcatcg tgaaggcata caactttaaa atattgtcat 9180  
gattctcaac aaatgtttga gcactcactc catagattta ttgcatacct aataaaacaa 9240  
taacttatgt ttgtgtaaca ttttacaaca taaaagtac ttttggttgt atcatcttgc 9300  
tttgttcttg aaactcagat acattttttac tttaccctct tacagaagaa attgaggtgc 9360

agaaagaaat tatttgcctt gaattgcagc agtaagtgcc tacagagtga ttttccatat 9420  
 tctaagaata ttgatacagt tcttaatctc aaattatgaa gtcgaatctc aacagtagat 9480  
 cagattcgga gagagcctta aaatgtgggt ttaacatgag tgaacacatg tggcaaagat 9540  
 aaagaacttg gttaagcagt ggagacaagt tctccagcac tcacaccctc tagaagctgc 9600  
 agtaaacagt cctgttttct agagagaggg cactattcat ggcgttggtc agaacgttac 9660  
 agattgtggc ttatgtcctt cactcctgca cttggccagt ctccccattt ctccagcaag 9720  
 ccagcagtggt gtccttgagg agcgggcatt tatttaattgg accttcattt tcttctgctt 9780  
 ttggtgggtg cttctagatg gcattataat cagaacacat acttagatac tgcaatgttt 9840  
 gcccgtagcag gaactagaga ttataaaatc ccacatatcc cccatgggtg gtctgatctg 9900  
 ctgtgtgttt gctcccagga gccatctgtg tgtgaaaagg aggccttgcc catctctgag 9960  
 agctccttta agctcctcgg ctccctcggag gacctgtcca gtgactcgga gagtcattctc 10020  
 ccagaagagc cagctccgct gtcgccccag caggccttca ggaggcgagc aaacaccctg 10080  
 agtcacttcc ccatcgaatg ccaggaacct ccacaacctg cccgggggtc cccgggggtt 10140  
 tcgcaaagga aacttatgag gtatcactca gtgagcacag agacgcctca tgaacgaaag 10200  
 taagatttgt ttaaatttgt tgcataaata gctggggcat atctgtgact agccagggtat 10260  
 gtgcatccca ggtatgttta ttgagtga gaaatgagtc aggcctttact cttggtttgg 10320  
 agataaaact ggaagcagtg acatgttcgt tcgagctgct tgtgagtata caagcaatgg 10380  
 gtacttgtat tgtcaggaag caagtgaag tgagcaaaaa tggtagcctaa catgcatagt 10440  
 cattactcct caaacaaggt aagagacggt gttgactgtg gaactttgct gctgtgagga 10500  
 agagggcaag cggatgagtc tccccatctg aagccctgga gcagggttat aatgggaggg 10560  
 agaggcgctg atccttacag gcagagcaag agaggtatgc tggcctcata gggtagcagg 10620  
 ggtgcttcag cttctggtcc tagctctgcc gtgaactaat tgtgacctgg acgaatttgc 10680  
 taaattctct gaataacaaa attggagtag atgttttcta aaatctctca ctgtaagaat 10740  
 tctagattct tctaacaaga ttattcatt gtaatagttg gggtcctgtg accagttaga 10800  
 atcgtctggt tatggagaag agtaatcaga agttcccccc attccttcca agtgtccctt 10860  
 agtgattcat ttaattctgt gtgccagaga ctataaatgg acacagttat ctttaaaaac 10920  
 aactttaaac aattttaaaa atctctcacc taatatgaat caaggtcaca cctgtgtaca 10980  
 gtcgctgctt tcttctgacc agcagccgca gaagtcccag gacctatgtg ttcgtgtagt 11040  
 tcatacacgg atcattgaga gtgtgagtta gtacagaagt gtttggaatg ttctgagtaa 11100  
 agaagtgtga gcattaacag tcctggatga tggagcagag cctcccagct ttgttttctg 11160  
 tcagccattg gaaagagttc ttggttcttt ggaattcagc ggggtagtgg tgatcccaaa 11220  
 agcaggggac atgtcagaag gtactgctta ataaatacac gcttttagag acacacatcg 11280  
 ttgggttgta gctgtgtaag tttcttgtgt ttaacaccct gtctgcacat tacttctgtg 11340  
 ctgcctcacc actgcctgcc cactcctctg ttgttggcgt tttcagtgat cattgaaaca 11400  
 ttctgtctg gagagtccta gttctcttgt gaagtctgct gtttctcaaa agccagagtt 11460  
 gataggactt agtatcagta cttttccttt ctccatgaag aatgtagctt tataatagat 11520  
 gatgtcacac atccgtaatg ggagggatga ggagatgctt gtctgtctgc ctctctagca 11580  
 tggcccatte tgctttcttt ccccttctgt agctcttttc cgatttatct acaggaaata 11640  
 agacattgaa attcagggca ggatattgtt cattttaaaag ggaaatgtat tttttaaaag 11700  
 tcagtttttt ttgtcttttg ttataacttt aattaaaaat ttttttctc gccagttcct 11760  
 gaaaaagaaa atagagaaaag aaatattatt gttcctgggc gaagtggctc actcctgtaa 11820  
 tcccagagct ttgggagact gaggtgggag gttgcttgag gccaagagtt caaggtcaca 11880

gtcagctgtg atcgtgctac cgcactccag cctgggtgat agagtgagac ctattaaaaa 11940  
 aaaaaagtat tgttgaggagc ataaacacgt gggaaatggg caagaacggc cgtcaatata 12000  
 ctctgttttt cactgaaaac tacctttgcc agagagcgag cagagatgag gaaaaggagt 12060  
 ggaagaagtc ctccactctg atagtgttac tggaacaacg agacaaaagc ggtgtgctcc 12120  
 ttccacctgt ttgctccgtg tccctgtcgg cgccccctct cctgctaacc ccccggtgct 12180  
 ttctctgatt gctgtttagt gtggatcctt cacctgtggg tgagtctaag caccgcccag 12240  
 gtcagtcttc agctcctgct cctccacctc gtcttaaccc ctccgcctcc tcgccaaact 12300  
 tttttaagta cctaaaacat aattccagtg gagaacaaag tgggaatgct gtgccaaaga 12360  
 ggtgagcaca ctacagtggc aagtttggtg ttgtctgttt tccctggggag ttcacactga 12420  
 tgaggatgtg ctgaatgggg ggaatgtcca tgcaggaagc agagccactg tgtgtgtgtg 12480  
 tgtgtgtgtg tgtgtgtgtg tgtgtgcgcg cgcgcgcgtg tgtctttgtt tatattttgt 12540  
 cttattttca gctgtcattt gaaccaagtt aattttacta ttgatgactt ttnttaagat 12600  
 tattatgaaa acagatctta atggcagatt ggtttgtgtt tgtgtttgtt tttttttttt 12660  
 ttgagacagg gtctcactct gttccccagg ctggagtgca gtggcgtgat ctcggtcac 12720  
 tgcagcttct gccttgtggg ttcaagcagt tctcctgcct cagcctcccg agtagctggg 12780  
 actacaggca cagccacca tgcccggcta atatttttat tttttttgt agagatgggg 12840  
 tttcaccatg ttgaccaggt tgttcttgaa ctctaacct caagtgatcc gcctgcctca 12900  
 gtctcccaa gtgctgggat tacaggtgtg agccactgca cctgctgca aattgttttt 12960  
 ttatacttat tttcacattt ccttgcccta gtggacactt acatgcatgc gtatatacac 13020  
 acacacgcgc gcgcngtgcg cgcacacaca cacacacaca cacacacaca cacacacaca 13080  
 cacacaggat aacatctgtg tttgatcatg tacactgcaa tttgtgccat atcagaaact 13140  
 tccctgattga tttaggggaa ttatttttcc cagtttgaaa ggaagagtta tttggaaaat 13200  
 ggatggattt tcttttttaa aaaattattg atcccattca tttaaaatca aattttattg 13260  
 gtgaaaatga aaattaaatc tcgttcgtga actactttta atttcttacc tagttttctt 13320  
 ttcttagcat tagaacaaaa atgtttcttt tattttgaag cttatatatt atactttgtg 13380  
 tttttatgtt tctttatcct aaactctttt ttcaaccaa ctcttagcat ctccactagt 13440  
 aatgccctgc ggaaaaaact tcattcttct tccctgtgct caaattttct aaaatttctg 13500  
 gtcctgttag atgaaaaata cacctctgat tttatgaaca caaaaaggta gggcttaatt 13560  
 tagatatatc aagcctgggt gttactaagt gttgaatata attagatata caagggtgtt 13620  
 ttaattacta ttttgccatt taaaaaatca tttcagctaa atctgttgta tcttctttct 13680  
 tatacttttt tcttactgaa tgccattttt aaaaatgtgc aaccaacctg ttctctagt 13740  
 ttgacgagga ttagtttaag tgttgtctta agaaaagtct ttgccaagtc tctgagacca 13800  
 gtgtttctgg ttagtgagca tatgtctgtt tcaaacagg atgtctgatc tgttcaggac 13860  
 gtctaactctg taagttgagg ggattgctta cttacaggta cataacttgg gtataaattg 13920  
 gaagggcctt ctcaagttgt cctgtgaata ggagaaaaca tttatgattg tgtttatata 13980  
 ttgataactg tattttgtag tttaaaaaat acacacgtta aaacaattat catcatcaag 14040  
 tgactgcata gttattgcct tgctggttct gtgtaattaa attgcaagtt ttttcatttt 14100  
 ttgtgggaat ccttgagagc atgggcctgt gctgagcaga tattcccatg cacagaagag 14160  
 ggcagaatgg ggccccctgg catcaccccc ttccccctt taggcagttt ctctttatca 14220  
 aagtggcacc aagagaggcc caattggaac tatgatatgt ggaacatgtt tcttaatctc 14280  
 tgttacaatc gaaatcactt aagggcattg aatctttctc ttttcatgaa aagaattctg 14340  
 taagaaagca gttctttagg aatgatgacc cactgtgagc ttgatataac ttctgtgatt 14400

T0520 T0520

gattatattgt	ttatacaaag	atagttgata	atttagtgat	ttgtttaaaa	aaatgttaag	14460
ctaacaaaat	cccgtgaatt	cctccccact	agtcataaat	caatcatctt	ataatttttag	14520
ggactttgaa	tccaaagcaa	accatcttgg	tgattctggg	gggactcctg	tgaagacccg	14580
gaggcattcc	tggaggcagc	agatattcct	ccgagtagcc	accccgagca	aggcgtgcga	14640
ttcttccagc	agatatgaag	gtaaggccgg	tacctgaaat	gaaacctcaa	agagagcacg	14700
ctgacagagg	accctgggag	ccycatcata	ttggtaagaa	agcagagcgc	cgtcctcttc	14760
agtattggca	ggctctgaggc	aatcacaaaag	gtaactaggg	aggggaattta	gaggttaccc	14820
tccattttctt	aggggaaggaa	tttaaagcta	atttagggta	acctctccat	aaacaggagc	14880
agagctctga	tgttttagagt	ggtcacagt	ttaaccagcg	gtgaatccag	acaggtctgc	14940
ggcaacctca	cttcttgccct	cctaggacat	aaggcaaaaag	gagagactga	ggcaagtttt	15000
agagcagcag	tgaaggttta	ttaaaaactt	cagagcagga	atgaaaggac	gtcaagtaca	15060
ctttgaaggt	ggttaggcgg	gcaacttgag	agatgaagt	tgagatttgg	ccttttgacc	15120
tgggggtttta	tatgctgccca	tacttccggg	gtcttgcggt	ccttcttctc	tgattcttcc	15180
cttgggggtgg	gctgtccgca	tgtgcattgg	cgtgctagca	cacggggggt	gtgggggagc	15240
gtgcgcaggg	tgtttactgg	agttgtaggc	gtgctcactt	gaggcggtct	tcctgtcca	15300
gtctagcatt	cctagaggaa	cgatcatgcac	caggtaaatt	ccgccatgtt	gcctcttaat	15360
gcgcattgctt	gagccctactc	gccagctcc	cgagatctta	ttgggaagct	gcagctcccc	15420
agtttttaggt	gttttctatc	tactgggagc	ccgcccttcc	ttggtgccc	ctgtgaccaa	15480
cgatcacttt	agagaaacag	ttgacaactg	cctgaccaac	acctgatggt	cgctgacat	15540
tgctgggtgca	tatctggaaa	gggccctctc	ctgccgtcct	catgtctgac	gagctaccg	15600
ctgtaaccaa	agcgtgggct	tccgagctctg	ctttcaaatac	ccagcttttc	cccttaggag	15660
ctgtgaacta	gaataaactg	tctaaagtta	ccacctataa	cctgggatta	attatgcctg	15720
ttgccacact	gatagagaca	aggcagcatg	atatcattac	tgatacattt	tttttaaagc	15780
attcaaaatt	catagtactg	gaaagaaaat	cagtgatgcg	aatgtttcca	gggtaatgtc	15840
acctcccatg	ctgtggaagt	ccttcgggtg	agcctggccc	cttgcttctt	ttgccccagc	15900
ctttctatgt	gggggcacca	tggagctgcc	actcaccagc	accttttttc	cctcaagtag	15960
tttgtacct	ttaaagtattc	ctgccgtggg	tggccccctg	gtggagctgc	tgagcctagc	16020
caggggtttga	tttctcttcc	tgccagtgtg	agccagatgg	ccacatctct	cttccccctg	16080
cccggtggaga	ggctctgctta	ccgcaaagaa	gggctcttcc	tcccaggctc	tgtagcacc	16140
tgttagaggg	tgtggagtgg	agcagtggga	accagagcca	ccagagggag	gccctggagg	16200
aggaacgaag	ctgattcatg	tctgaaaggg	gtgccagaac	ccaagtttcg	gtgtttaata	16260
aagagtgcct	cgggtgttgcg	gtggccatac	ctcacagggc	atggctcgctt	ggaaatttct	16320
gctcggaaat	gctttgtgca	gtggccagga	tgcgttaggg	gccacagatg	actgcttgct	16380
ccatcataga	acagttccaa	gttttcaaac	gagcattcac	agactgagcc	gcacctgcc	16440
tcctgtcct	ctgattcctg	gcttcttctc	tggctctctga	agccacacgg	aaatgtgttt	16500
gcacctgttt	cctgcccttc	agatgacaga	ggaccatgga	agctgctgcc	tccttttagct	16560
ctcttctcca	ggggaattgc	cctcgtcact	gtttgggaac	ccctgggtccg	agtcctgtcc	16620
tccgaagagc	ctctgccccct	cctggagctc	tgagttgaac	ttggtgttca	cttggcctct	16680
ggctctggca	gtgtgttgct	ccttcggttg	acctgccact	gctctgttaa	tgcagattga	16740
tcttcataat	ctgtttctgc	tttaagtgat	taactcaaac	attcttggtc	cttattctat	16800
cttgtccttt	gggatatgaa	ccattattta	aatttggtgact	ggtttcctgg	cttggcacag	16860
ttgaccatgc	ctgtaagctc	agtgccttgg	gaggccaagg	caggaggatc	cctggaggcc	16920

aggagttcga ggccaccctt ggcaacatag tgagaccctg tctctacaaa aaaataaaaa 16980  
 ttagctgagc gtggtgttgt gaagctgtag tcctagctac ttgggagact gaggcaggag 17040  
 gattacttga gcccaggagt ttgagtttac agtgagctgt gatcacaccc ctgcactcca 17100  
 gcctgggcaa cagagtgaga cttgtctcg cgggggtggg gagtcatgtc tatacttgag 17160  
 aagttttttt ccctcgcata gttgtacct ataaagtatt catcagtttt gagcagtcct 17220  
 tttgcgattg ttttgagtct tgattctggt gaccagtaag ttgtatatat ttgcctgtca 17280  
 agtggacaaa catggccttt gtgcctttaa gtaatggcta aaagtaccaa acagaacagg 17340  
 gcctggcata gatgctgtc ctctgttcc taggccgtaa tcaccctga ttcacagac 17400  
 cccaaacaag tcagctctc tccctcctgt gccccaccac ccaatctcct gcaggaagat 17460  
 gtctggagac ccctgtcagc gctaggcaga gatcaccatc catgtccacc tttcctctga 17520  
 tgcaggtcc cactagcccc tctggcttgt gccatgccag ccatgaactc accctcatgc 17580  
 cccacccgag ccctggcaca ggctattccc tctgcctgga atgctcttcg tcagtatccc 17640  
 catggctccc tccctcccct tcccttgtat cctgactctc ccatagcagc tctctccctg 17700  
 taacacatgt tcacaggttc atcttcgtca cccatctccg gcagctcctg caggcttgat 17760  
 ggctgctaaa ggcaggcaag tcagtggctc agattcttgc aaacttagtg attagtgatt 17820  
 tctcaactcc ctctcatgc ctctctgtc tctataggca catattattt cttatctctt 17880  
 ctcaagaacca agccgcctga atttctgaat aacattgttt aagtgttctg tgtatgcaaa 17940  
 agaaaaacga gaataaaagg attattaagg aagaattaat ataataatag ccacatatta 18000  
 tgctcttttt atactctgct aagtgttcta catgaattat ttcctttaat tagacaatct 18060  
 taagaacatc gacattttta tgaagccat tttacagggtg ggtgagtgga ggctgggagt 18120  
 ggcttaaata actttcccca aaccaggag ttagtgggag ccagaggcag gacctgagct 18180  
 cgcggtctg agctccaaag ctattctct gaactgtgca cagcactggg ctgcagccag 18240  
 agatgcagga cgctgcggga ccctctggag gtggtcctgc ctgtgcttcc ctctcccccac 18300  
 aggaagctcc ctataggcat ctgtgttggg cgtggactct cagtgtacct gcatgtctcc 18360  
 ctgttggcca gacaccaaca ctgaatggaa aacatgtttc tgggcatttt aatgtacgta 18420  
 cttgccttca gtcàatctcc tccgccccct tccatcctga ccgcctccct aatagttagc 18480  
 agtgggactg gagcttgáat ggcaactgat ttctgtctga gaggacaaat caggcatctt 18540  
 tgtcctctgc cactgtctgt tccccatcct taggatgcac gatgccagag ccctccactg 18600  
 tggctctgtga ccactttgac ccacactagc aggtctccat atgttccctc cagctgagag 18660  
 acatcacatc caaagacagt ttagagctct gaggtttctt tccccagagg tccctgcttt 18720  
 gtgcaactg tctccagcca agcgtgcaca agactctgtt cctgatattgc ctgggcggct 18780  
 gagccatggg cagctgagcc tgcagccgt ggactcactg cattcccact ctgactttgg 18840  
 catgaaagac acacaagtgt gcttgtgaga aatagatctt aacagtacct ttaaacacct 18900  
 atttcagggt ctcaaaatga ctgcctgttt tacattttata ttctggcagt gcaaacttca 18960  
 attggacagg aaatcttaca acctctcttc cagggtgraaa agcgaggcag ggatgtttat 19020  
 acagttccat ccatgtcctc ccacttggaa gatactagta aaacacacca acagtaatac 19080  
 aaaaaccatg gtgtttgcaa tagtgataat gttactagtg aaggaaaaat agaaactttc 19140  
 tgtaatttgg agattctaatt ttttatagggt gggctaaaaa aaaaayctgg agagaagggt 19200  
 gttaagttag taaggagtgt gtctctaact aaatatagt taaaaagaga agaaaatata 19260  
 aagtcaggca cagtggatag aggtggatag tctaactctc aatagtataa tgggcaaaat 19320  
 tgtctcaaac aaaattagtc tgccctcttgt ttactcaggg atgtgtgact gttttctatg 19380  
 cacaaaatcc ccatgaaata attaagttgc aagaatctga actttatatt ttggaaacct 19440

atctgaggta ggtaggaagt taattttatat ttagaaat tttt gcttgcatat gtctagtagc 19500  
 tccaggacaa atattcccaa atcccagact attttttttt cttttttaa tcaacagtga 19560  
 ccagttgggc tcttgtaaga attacagcct taagttagca aagtctaaga gggctgggtt 19620  
 taatcctgaa cctcagaggg tccctgcttc tcaaatacta agtaggtcac gtgcacagca 19680  
 ggtactacat tgaagggaaa ttgtatgata aataggaaat cagcgatttt tacttgga 19740  
 cttggcaagg caaatgtttt tgtaataaaa atagatcgtg aaatagaatc ctgaaagctg 19800  
 cctgttttaa tgtaaagcaa atggcttttag tgatgcttta agtgtggcag tcacttctgg 19860  
 ctgccgcaga aactatagaa agtgcattct ctcttggtgc tgtgggttct tagggatga 19920  
 gccttggtg acgctgagta tgtggaagga ccattcattc ttggtaacta tacactagga 19980  
 agaggggtggc gttagcgaag ctactgcagg ttgggtgtgt ttaagatttg gatttatttt 20040  
 tcttttaatt tttattttta gttccagggt acatgtgcag gatgtgcagg tttgttacat 20100  
 ggttaaactg gtgccatggg gggttgctgt acctatcaac ccatcaccta ggtattaagt 20160  
 ccagcatgtg gttatttttc gtaatgctct ccctgctccc tgccgcccc caacaggctc 20220  
 cagtgtttgt tgttcccttt cctatgtcca tgtgttctca tgattcagct cccatctatg 20280  
 agcaaaaaca tgtggtgttt gggtttctgt tcttgcgtta gtttgcagata gataatggct 20340  
 ttcagcttca tccatgtccc tgcaaaggac atgggtctcat tcatttttat ggctgcagata 20400  
 tagtccatgg tgtatatgta ccacattttc tttatctagt ctatcattga tgggcatttg 20460  
 gggtgattcc atgtctttgc tattgtgaat agtgcagcag tgaacatatg catgcagata 20520  
 tctttgtaac agagtgggtt atattccttt gggtatgtac ccaggaatgg gattgctggg 20580  
 tcaaactggt tttctagttc tagatctttg aggaattgcc acaccatctt ctacaatggt 20640  
 tgaactaatt tacattctca ccaacagtgt aaaagcattc ttacttctcc gcaacctcac 20700  
 tagcatctgt tgtttcttga ctttttaata atcaccgttc tgactggtgt gagacagtat 20760  
 ctcttctgtg ttttgatttg catttctcta atgcatcagtg atgttgagct ttttttcatg 20820  
 tttgttggtg gtatgaatat cttcttttga gaagtgtctg ttcagagag agacatat 20880  
 gctcctctga gtaaagggtg aggatgctta cgtctgtgtg acagccttct ctctttttca 20940  
 gaacctcact gtggatcgcc atcgttggtc tgtactgaag gtaaagcaga tagaggcagt 21000  
 ctcatctgtc agatgaagac ctcatacacc tgttgattaa gaggttttct tcagatcatg 21060  
 gtttagagcg gtgttttaca aacttgacgt gcttgagtc tcttggaat cttgttaaaa 21120  
 ggcagactct gttgcagtag gtccgggtgg gttctgaagt tctaacaagc tccccagtga 21180  
 ggctgatgtt tcaggtccac tgtgaggagg cagggttag aataaacaac cgtggggaat 21240  
 ccagtcaga tctttgatgc atcctaggtt aggcctgtct gtcaggctgc cctgggtctc 21300  
 tagtgatgga ctccaggagt ctctcaagtc tcaaataagt ctgagtcac agggatattt 21360  
 tttgagaaga gttgtgctgt ctgaagaagc aaagagtga tgtgatgggg aaaatgcagt 21420  
 gattaaaaac atggtaaggt ttaaagaaag atttgaccat atgccagggtg aacccaaatg 21480  
 tatgggtgctt tgcgtctttc ctgccctttg gttttccagg gaggcaaggc cttatctctt 21540  
 atggagcaca ggagacacag tgtgggcgtt tgttttctca gccgtgggct ctaacctaat 21600  
 tgtcaagcct acaaaaaaaaa aatgctgaaa atcaacttct gactagatat ctggtagtac 21660  
 ataactctca taattttctc tctgggtgta ttatgcaaaa gataatcctt tngttattaa 21720  
 gaaacaattt ttaaggcaac tccaacttt gaaacgggga aaaatcattt tatttacctc 21780  
 tatgtgctag ggaacaatat taaatttagt tttatacttt tcttttaagc atttcagatt 21840  
 atattgtgca tttcaccaac aatagaagct ttcagacttt atatgtcttg taaaaaaaag 21900  
 cctaatatag ataagaataa tttattgatt tgaaacccat tgtataagaa atagtcaggt 21960

gaaacttaag	ttcaaagt	tttttgtctt	gtggatgtag	ctatgtcaat	atgcctagtt	22020
tatagtaaca	ttaagtctag	tggattagat	attagatatc	aattgagatg	taagcagtaa	22080
taaacagtaa	tgcctaaact	gaagtatata	atctgaatct	ttatgatgac	caatttatat	22140
tattgtgaaa	aacttaggaa	ctgatttgaa	acatgattta	catgttttac	atgaaacatg	22200
atttacatgt	gtcatatata	gttttcaata	atttacgtac	cagcaggaaa	ttttagtggg	22260
taagtggaa	aaactgcagg	tgaactttg	ctggaaaata	caagcatagt	gacatctgtg	22320
caccaaagc	acctggggag	attttttaaa	acatgggcca	gacatgcctt	cctggctgtc	22380
tccctcactg	tcagtgagtg	tggggatggg	gtctgggcct	gaattctttc	tttttagact	22440
cctcaggatt	ctgattgctg	ccacgttgag	aggggtgacc	tcaattcgga	cctcagaggg	22500
tgacttgaga	aactgtcacc	acttggtggc	agtgttgctc	cccgcacctt	gattgccctt	22560
gtttctttcc	aatcccggaa	aagtgtgctt	gttttttttt	ttttccctgc	gtgtttttgt	22620
ttttgatctt	gctataatat	ttatatcctt	tgctcatttg	caacttat	gaatggagag	22680
ctactttctg	aaatctagat	gtttttcttt	ttctacaggg	ttttagggca	tgggcaaaac	22740
acggaagaaa	aaagtgtgtc	tcagttggca	gagacgtgga	tttttaagat	tgttcttaat	22800
ttactttctg	tataactttg	cttttctgtg	gtgaacaaa	accagggtca	agataaaaata	22860
ttgcaagcca	agaatctgat	tgttcatgga	tttctatggg	ttaaagatact	tgatcacctc	22920
cccatccgcc	ccctaccca	cccaccctgc	gccgccccca	cacccattg	tgcttcttgg	22980
cttgtcattt	caaaagtcaa	ggaagtcaca	gtgaatggca	agattttacc	tcgacttgct	23040
atttttgtgc	ctgttaacaa	ttgtgagtta	acactgactg	agcttttcct	agtgaacctc	23100
cggcgtttaa	acagccagtc	cataacactg	tgtgagggct	ggagctaagg	ttattgggtga	23160
cacaagatag	cacctgagcc	agtgtgtgct	ggtaggaggg	ctgaggggaa	gagggctgag	23220
ggcttgagtg	ctgagatgct	agagtcacat	cgcctggatt	tgaatccctg	ccctcctgtt	23280
ctgataccag	ctgacccatg	acgatgctac	agcacctgac	agcagattcc	tccttagggc	23340
tggtctaact	ctagagtgtg	tgccctgtgtg	cctgcaggag	aatgtccaaa	gtgggtgatc	23400
ttgatctgtt	aacctttgaa	ttttaaccta	taccaggggag	ccattgaaga	gtttaaagca	23460
agtgaatgac	gagtagtttg	aaaatatttc	caggtggata	gaatttgtgg	acatacatga	23520
acatgagcag	cctcaaaatc	agggctggga	ctagagttag	gccagcacgt	gtccagggtg	23580
caaaatgtaa	ggaggcattc	actttcaggg	cctggcaggt	gtggaccctg	aacttcagg	23640
accttgagag	tgagtgtctc	ctaaggatta	cacctggggg	gcctatttgc	ctcatcctgg	23700
tccctgggtcc	tctgtgtacc	ctattgcctg	cttcagtaaa	caggcagccc	tgcaaggggaa	23760
ggaaggggtg	gatcagctct	gaggaggggag	ttttttttaga	aggatagatt	tgttttgttt	23820
aaaaaacagc	tttattgaga	tataattcac	atcctatata	gtttgttcat	ttaaaatgta	23880
caattcaatg	ttgtgaggtt	attttttggg	atatccacag	agttgtgtga	acatgaccac	23940
aatctaattt	tttttatttt	tttttttttg	agacggagta	ttgctctgtc	gcccaggctg	24000
gagtgcagtg	gtgcgatctc	ggctcattgc	aacctctgcc	tcctgggttc	aagtgattct	24060
catgcctcag	cgacctgagt	agctgggatt	acaggcatgc	cccaccaagc	ctggctaatt	24120
tttatatgtt	tactagagac	ggggtttcac	catgttggcc	agactggtct	ccaactcatg	24180
gcctcaagtg	atccttctgc	ctcagcctcc	caaagtgttg	ggattacagg	cgtgagcccc	24240
accaccgca	gtctaatttt	gaaacatttt	ttgtccccct	agaaaaaaac	ctgtagttgt	24300
cacttgccaa	tctactgccg	tccacctcta	accatagaca	gcccctaata	tactttctgt	24360
ctctatagat	ttgcctattc	tgaacacttc	atctaagtgc	aatcatataa	tatgtgggtc	24420
tttgtgtctg	gcttctttga	tttaacatgt	tttcaaaatt	cattatgtca	taatacatac	24480



cagtaatcca ttctttttta atgacttatt aatattccgt tgtatagaga catcacatat 24540  
 ggttttatcct ttaccagtcg agaggcattt ggattgtttg cacttttggc tgttacggat 24600  
 aataccgctg tgaacattga tgtatgtgtt tttgtgtgtt gaatgtgagc tgggtgtggaa 24660  
 actcctcctc cagggggggc ttacctgtga ttctaccacac ggggatgggt aagccagcag 24720  
 ggatgggaag ggtttggtcc tgctggccct aggctttcct gcaggctgcc atgtgccttt 24780  
 cttctgccta ggctgaaacg gaggctgccc tggttttctgg cactgccctc gtgagtgtgt 24840  
 ggggaaggctg ggggaagcca agtctccatg gtgcctccat cagggaccct gcagctggga 24900  
 ggcagccaga gggccacagg ttggtagcat tcacacagag ctacatttct tttttttttt 24960  
 tttttgagac aatcttgctc tgtcgccag gctggagtgc agtgggtgga tctccgctca 25020  
 ctgccacctc cactccacag gttcaaggaa ttctcctgcc tcagcctccc aagtagctgg 25080  
 gactacaggc gtgcgctgcc atgcccggct aattttttgt gtttttagta gagacgggggt 25140  
 ttcaccacgt tgaccaggat ggtcttcac tcccgacctc gcgattcacc tgccteggcc 25200  
 tccccaaaag tgctgggatt acaggcgtga gccaccatgc ccagcctaca tttctttttt 25260  
 ttttttcttt gagatggagt cttgctctgt caccagggct ggagtgcagg ggcaccatct 25320  
 ctgctcactg caacctctgc ctctgagtt caagtgatc tctgcctca gcctccggag 25380  
 tagctgggat tacaggcaac tgccaccaca cctggctaatt ttttttattt ttatttttta 25440  
 atagagacgg agtttttcca tgttgaccag gctgggtctcg aactcctgac ctcaagtggc 25500  
 ctcaagaggc caatccgcct tggcctcccc aagtgtctgg attatagggtg tgagccactg 25560  
 caccaccca gcccgtagct acatttctgt cagctgtttg caaactgtgc ccagaaatcc 25620  
 cctggaggac ttgtagaacc accagttact ggggttacgcc cccaaatgtc tgatgtctgga 25680  
 gatgaattat cttgggtgga gccctcaagc cgcagcagct gataagcatg gggacctcct 25740  
 attctgataa aaattccaaa aaagtcctga gtgattaata aacagcacat tgaaaattag 25800  
 aaatgagttc tatggcaggg gatgaaacag gcaacaaaagc ctattttctt tgcaatgaag 25860  
 cgcacagat attaataata gccattgtaa ttatctttat catgtattaa gcattttgtg 25920  
 tttttcactt ttacacaatt agatgatccc cataggtatt accgcctttt tttttttttt 25980  
 ttttttgaga cagagtcttg ctctatcccc caggctggag tgcagtggca cgatcttggc 26040  
 tcaactgcaac ctctacctc caggttcaag ctattctcat gcttcaccct ccttagtagc 26100  
 tgggattaca ggcgcctgcc accagaccca gctaattttt tgtatctttt ttagtagaga 26160  
 caggggtttcg ccatgttggg caggctggtc tcgaactcct gacctcaggt gatccgcca 26220  
 cctcggcctc ccaaagtgtc gggattatag gcgtgggtca ccacaactgg acttactgcc 26280  
 catcttttaa gagatgagga cagaaagatt gagtgcacac gttatgtctc ctgcagctct 26340  
 tgggttcacat agccaggatt cgtatcaatc tatttagctc taaatctagt ctcttaatca 26400  
 cagtaatgaa ccgttgacag ttttacgagt aaattatcaa gagttttgat aggtttgctc 26460  
 acttaaatta gtgcttgtag agtaatgggc tgtgttagtg tgaaggaatg tatcttatgt 26520  
 tggaagtact ctagaattaa atgttaactc ttgctaataa agcatacatt tggggcatta 26580  
 ttagcaactt tttttttttt ttttttagcaa aattagaggc ttcctagttg agtggtttat 26640  
 gttatttata tttatttatt tgtttgtttg tgacaggggtc ttgctctgtc acccaggctg 26700  
 gagtacagta agtagcacia tcatagctca ctgcagcctc gacctcttgg gctcaagcag 26760  
 tccccgcct cagcctccta agtgcctggg accacagggtg cgcacacca cgccctgcta 26820  
 aatgtttaca gttttttag agacaggggtc tcaccatgtt gccagggctg gtcttgaact 26880  
 cttgaattaa agcaatcctc ttgcttcaga ctcccaacat gctgggatta caggttgtgc 26940  
 cactgcgcca ggctccatg tatttgaatg aaagagcaga catctcctgg aggtggcaaa 27000

gctatgcatg cccccctgg aggggagctg ggggctctgg gggtacagtg atggcacatt 27060  
cagggagctc tccgctttgt gagatcctga gataaagcca aaggatgcat taaactgctt 27120  
ctaaatgaac tttttccaag tgaatttggt atatcacttc tatataaatg aaaatatttg 27180  
cagcatgagt actaacaaga tttttttttt cttttacccc gatggagtct cgctctgtcg 27240  
ccgggctgga gtgcagtggg gcaatcttgg ctactgcaa cctccgcctc ctgggttcaa 27300  
gcgattctcc tgcctcagcc tcccagtag ctgggattac aggtgcgcac caccacgccc 27360  
agctaatttt tgtattttta gtaaagatgg gggttcacca tgttaaccag gatgggtctct 27420  
atctcttgac tttgtgatct gccgcctcg gcctcccaa gtgctgggat tacaggtgtg 27480  
agccacgctc ccggccaaga ttttaaacat tatttaccaa agtaggaacg tggtaattat 27540  
ggctttatat aattctgaaa atgatttcta gtaccaaact atgaatttta tacttgaaag 27600  
aatgatgggt ttttcacaga aagttgaagt tattatgggt tgtnttctcg ttcanggtgt 27660  
ttttgctgga gaatgttcga tgaacagcag ttctgggtgat aagttatgga tgtacacagc 27720  
tggtgtgggt tttaggattt tattttgcag cagcatcttc ctcaaacagt tgccagggga 27780  
aggctttcct tcttcttact ggtaccagcc tttctcttgc agacaaggca gtatgggagg 27840  
gttgggagac aaaacagaag ctggtgggtt cttcagcctg gcaaggattc agattgcagg 27900  
ttatagattg gaggccgtca gtggggatac ctttcoggac aaagtgggtgt ttctgcctgg 27960  
cactgcttgc cagagaagtt tcagttcttc attctcgtc agagaaaccc atatggacca 28020  
cattctgata gttttcttct gtttccctaa caccgaaggc tcagcccctg gtgcaggctc 28080  
cagtgtacag caggctgcat acagtttagc cagatgttct tgtagtacga aaagtcaccg 28140  
agtttccatt cacttggtgg tggcagggtat ggccctcctt acctcccatg gcccagggtt 28200  
ctctgtcctg ccgttttcac attttccagg ctttcacctc caggtaccaa aattcacatc 28260  
atthagagat tgtgtctgcc tgccaatacg cggatgtacc agtgagggat tgttctcgcc 28320  
tgacgagagg tctggatgat gagagagcag agctggccct ggggctcagt ggtgacaccc 28380  
tcgagcttgg ctgcttctgt tcttctgctt cctctgcttg gattccttcg cctttggctt 28440  
cccctccagt tccaagcaga acaaaacagg agatatcaag gaggaaaggg tgacccctct 28500  
atatctggag agcaaaactg tcgcggaaat ccctagtgtc cttccatttg tgtctcatta 28560  
tctgaaaccg agttacctgg ctggtcacgt gcagccacca gaggcaggaa ggtagtgatt 28620  
ctgcctgtgt ggaatgttct agcattccct ggtagctttt gtttcttcag gcagccatga 28680  
ctttgcatag atcatttctt tttgccagg acactcctgc tcgttttctc cctcctcac 28740  
caaaccaca gtgcattaac agcgacagac ttctcctcat cctctcaggc cacttggtatg 28800  
tcaccatttc ttctctttac ccctcaggcg tagtcagcct ctctgtgcct gatgttttat 28860  
ggctttgtgt atgccccgat ggagagcgtc ttactgtgtc ttcgggttat ttatctcaac 28920  
ctcgcactct tgctatcctg tacagtaacc aacagccaca tatcactatt taaaattaaa 28980  
tacaaactaa ttatacttaa atgtaataaa aatgtagccc ctcacattag cctcatttca 29040  
agagccacat gtggctaccg tattgtaagc agagctcaag aacattcagc aatattgtga 29100  
gagtggctac catattgaga gcagagctct agaacattcc cttatcccag aaagtctctt 29160  
tgacatgct gctcaagggt gtgaactctg agatctccag tccccccagc tccgtcactc 29220  
agaaccacaa atgtggcacc atcagccttc aggggtgggtc ttgtgttgta ctgtctcctg 29280  
actaaagaag taaacttcag gcagtcaaga ttttctacaa cccacactgc tcctaaaact 29340  
agtgttactg gatatgtaaa agctattgag ccagtgctt tcaaggaatc ctaaaagcaa 29400  
gtggggactg tcatgtatcc aggttccttg ttttgacaga gaagaaatag aggcttttagg 29460  
ggaaaggggt ccactcaagg tcatacaggc aaggtcatac agtcagtgggt agaattggact 29520

0976311 062501

ggaatcttga ttgttcatct cactccccctt tccattaagc cataactgat tgagtatcac 29580  
 caacctgttt gttttccctg attatgttcc cttctcrctt gtttaatcag gtttcagctc 29640  
 cttgtgagag gaagttgttt tcagcttctt agccccctac ctgtaggcgg tgctctggga 29700  
 acgctcagga agcagatgca tgtgagcctg tctcagccaa tgattgctta gttgcaagaa 29760  
 acaaaaaaat gactcaaact agcttaatca aaaggaggct tttaaacagc aagataaggg 29820  
 taaaggctgg gaactaggaa gttgtcagga accaaggctg tctctctgga tctctctttg 29880  
 aggccatgta attttttttt cctctcggtc tctttattct gcacaccagc tcaccagct 29940  
 tgcttggttac tatatgcagt catgccaaacc cccagatctg tatgacctgt tagcctcagg 30000  
 ggcaccaca tagctggctg ccaatctgtg ttttcttcca ggtttcgaga gggagaaatg 30060  
 attggcccag ctcagggtca cttaccaggg gagagggtggg ggaagtatgg aggcaccgtg 30120  
 gtatcagggg acccctgggc caagcttgct caaccgcgg cctgctttct ttcatttttt 30180  
 ctgttttttg ttttgttttg ttttgttttt ttacagctca tcagctattg ttagtgtatt 30240  
 ttatgtgtgg cccaagacaa ttcttcttct gctgtggctc aggaagcca aaagattggc 30300  
 caccctgtc ctaggccatt atctgggctg tgggagggtg ggagcagggt cagagctgga 30360  
 gggggagggg atagcctcca gccaccataa gttggtgtgt tcttggtaat tatattgctt 30420  
 gtcaaccgaa ggcagaatca ggacaatgaa agtaatgaga atccctagct ttgtaacagt 30480  
 tagtggttat ctaaaagtag gtgaaattgt acatgagtga gtggcatgaa tttcttatta 30540  
 ctaaagtgtc cagatagctg gctaactttc tgtcaaagat ccctctgcta ggatcaacat 30600  
 ttgattaata tattttatct gtaataagaa tttgggattc ttaaagcaaa atagttgtca 30660  
 tgtggctgac tacacaacca aagatggctc aggtgtcgct ggaagaggag agactgaaga 30720  
 gctgttgcca ggttcccacg tggaccttcg gcatgacctg gccatgggga ggcctcacac 30780  
 gctcctgcat cgcccacatc ttgccaagcc atggaaaaca cttgggattc atatctaaat 30840  
 cctagtttaa gcttgggtgag gacagtggcc tgggtgcagag tttgggtcat agatgggtgct 30900  
 tggtttcttt tgtataaagg ggtatatgat tttggaatat ttaccaaagtg tgggcatttt 30960  
 ttctataaaa attattgtat ctactgagat tatagtatgt aaaaaaaca tacacatgga 31020  
 gaaagaatac aaagagagca ttgatattct acagaagtgg caagaagatg tgggtgatagg 31080  
 tgatattttt gccttttgtt tcaattttgt attgtagtga cttttttggg agaaaaaact 31140  
 aantttctaa ttaagggaga aacatttgaa gtacatttag tctttctaga aaacctcatc 31200  
 ttctcataga agtttaagat ggagacatac ttccattgtg aatcatgatg ctaaccagta 31260  
 ttcagatttg ttggaaatgg actcagtttt aaaattgctt ctctcttggt ggctggaact 31320  
 gcaaatgatt gtttggggat ttttccccctt ttcttctatg gagttattca acttggcatg 31380  
 accagtgatt tgagctgaga acatggaacc cttgatttgc agaaatcaag cccccaagg 31440  
 tacagataca gtggtcattg tctgaagggt ttcttttgtt cttggccctc ctgtccctgc 31500  
 tcttactgtg gcagctgcag ctgcagggtc ctctgaagcc ttgccatcca tggtcacttc 31560  
 ctgcctgctc cccaccacc cctgggaaag agcccccaag tgtccaaaag cactgtgttg 31620  
 cctaagtctt gttgagagtc tacatttctc tagatctagc agaagtaaaa tttcagtttg 31680  
 ttatatttat agtttcagga atagtttggg aatggattta ataaaaaatt taaaagccca 31740  
 tcatttttat atctcttttt cgatatttga tggtttaaaa gacatcaaag ttatcttctc 31800  
 ccattactca tcctatacaa ttaaaacctg ttttttgaag ttgtaatagg taagttagcc 31860  
 ttaggtcacc ccatatttat gtaaaactcca gccactgcc acagctactt tgattgtgat 31920  
 ctgtcattgt gttaccact gtagggcaga aatggttcct gcctcatgcc gttgctgctt 31980  
 tactcttctt gaagtgggtg ggttctgtct ctgtagtctt tggcacactg taggttctca 32040

gatggcaggg tgaaaagttc ttctgtttgc ttaaatctct cataataccc tgagtctgtg 32100  
gagctcaata aattctactt ggtattattg atagattatt tggagccttt tatgttagaa 32160  
aagggattct taatccaatg ctccgtttta cagatgagaa gactgaggct caaagaccat 32220  
acccccagga gccatgattt gcaactgtatt taggaatagt gtctaggggtc agcacctggg 32280  
ggtggccgac tgcagagcag cctgggttagg agccctgggg ttgggcgggt ctgggctgct 32340  
gggtgccacag cagtctccct cccctggggac tttgggcctg ctaccacccc ctgttccttc 32400  
ctttgtgaga tagggctagc agtaactgtc ttgtttcatc agaggcagta ttgcataatg 32460  
aatgagagct ggggcctaaa ttaggcacaa gtgcaagccc tcagaaaact atgtacacct 32520  
agagagagag agagacacac gtctgtatga cagagaggca gggtttgga atgttctgat 32580  
ttcatgtttt gaattggtgt gacctttggg aggatctct tggaaatcgca gagcttcgtt 32640  
tacatcatga ctttcctgcc caccacatt ttctgagaag ccagagtttt aaatgtgggac 32700  
cccgtgagct tttctctgtt gcctcatttt ggccctgtggc cttttgtttt cttgggtatgt 32760  
catgaggcaa aataaaatga aactcagtgc tggtaataa ctcccatcat aatgtatatt 32820  
tctgtgaatg gctttttagc catttgagag gaaaaaggg catgtaaatt tcagaaagggc 32880  
ctgattggct ggagagtcag tgtagtgtca cagttaagag tatagattta aaaaaattt 32940  
tttattgtgg taaaaaacat aaacataata ctaccatcta aaccatattt aagtataaag 33000  
ttcagtagtg ttaagtatat tcacattgtt gtgcaatgga tctgcagaat ttttcatctt 33060  
gttaaactga aactctatgc ccaataaaca actcctatc cccctctccc agccctggc 33120  
aaccaccatt ctactttctg tttctctgag ttgactact gtagataact catttaagta 33180  
gagtcatatg gtatttgtct tcttatttct ggtttatttc gcttagcata atgtcctcaa 33240  
ggttcattca cgtttagca tatgacagga tttctctctt tttttccgc ctttttttg 33300  
agttatattc tgttgtatgt atatttaaca tttcttcat tcatctgttg acattcatct 33360  
gcttcacct tccacctttt ggctattgtg aagactgcag ctatgaacat ggggtgtgcaa 33420  
atgtctcttc aagatcctgc tttcagttct ttcggatatg taccagaag tgggtttgct 33480  
ggatcgatca tagtgtagtt ctgtgagtaa cctcatact gttttctgca gctgctgtac 33540  
cattttacat tccaccaac agtgcccaag ggctccagtt cctctacacc ctaccccaca 33600  
cttgtaatct tctggattgc agattttctg gatcaatctt ctggattaca cttgattttc 33660  
tgtgttgggc ctggatgttt agaacagtat cctcctttg gagtggtaaa tatgtaagtt 33720  
tttattataa aataatggcc atcctagtga gcgtgaggta atatctcatt gtgggtttga 33780  
tttccttcat agttaatgtg gttgggcac atttcatgtc ctggctggcc atttatgttt 33840  
catatttggg gaaatgtctt ttcaagtctc aagtcctttg cccatttttt aattgagtta 33900  
tttgattttc tactgttgag tatggattat taaatcagac tggcctgaac ttaaactcatg 33960  
gcccttccat ttttgacaa aagcagctgt gtgtcccat tgtgccttg cttcttcggt 34020  
gtaatgccg cataatgata gccccacct gtagttaaga gtgttggggc agtcagttag 34080  
gaagcactca ctccacagga gcttgttac taaggagaag gcagccggtc cattccta 34140  
aggggtctga aggaaggaag aagggtgaa ggaagtaaaa agagcctcct ccatgaatgg 34200  
cagccattct tgaaatccac cttggctgcc ttcattttta atgtcagtgg acttttaaga 34260  
caaccaaag gatgttcttg gatgaccaga gactgtggca gaggaggat ggtcacattg 34320  
ccaaggatct ctctcaacct cttggatagt gtgctgctgg tagtttgcac aattgcttca 34380  
gctttttggc aaagtacatg taaaatcctg aagtcactgc cagaggaaac ctggttcctg 34440  
agatagcagc ttgatgtcc tgcctcatcc caggtgcaca cctcactggg cagctctggc 34500  
tctgaattga gggacagcaa aaacctctaa ccaaccatac tgaaaagcag gcattggggg 34560

ctttagggga aggttctttt caaaactcat gatggggaga gaccaaagac tgggaatcat 34620  
 tgtaaagaag ttagtcatag atgcttctact ctttacaatc atcccaacac aagggttaaac 34680  
 aacatgcagt tttcacgatg tcccagaaag cgacgagtgc agtgaggtga aacgtggcca 34740  
 tctgagcaca caatgaccag gcttggaaag atcgatttcc cctgtgctgg ccctcagaat 34800  
 ttaaggcaca acttttaagc tgagtgtgca gactctgatt ttctatgttg ggcttgatg 34860  
 tttagaaaag tatctctcct tcagagtggg aaatatgcaa attttttact gaattacttc 34920  
 atttaataca agcagccgac ttctctgccc tcccctgttt ctgtcttggg gttgaatatt 34980  
 tgggtcccatg taacaactct tgattcttaa tgatgccaca tggaaagctgt gtgtgctggg 35040  
 atttgccata ttcagttatg gtcagtagag actttcttag tctctctctc tttttttttt 35100  
 tttttgagac aaattcttgc tctgtcacc aggctggagt gcagtggccc aatcttggct 35160  
 cgctgcaacc tctgcctccg ggttcaaatg attctctgccc ctcagcctcc cgagtagctg 35220  
 ggattgcagg cacgcgccac catacttggc taatttttgt attttttagta gagacagggg 35280  
 tttgccatgt tgtccagact agtcttgaac tcttgacctc gtcacccgcc tgccttgggtc 35340  
 tcccaaagtg ctgggattac aggcgtgagc caccgcgcct ggccacagtc agtagagact 35400  
 tttgaaagga aatattacct ctttaatgat gtttttagtc caagtaaatt gtggtaatgt 35460  
 ttaagaaatt tgcttaccac aaaaacagtt ttcaaggagc atttgaactt gtccacttta 35520  
 agtcataaaa tggattaaag tgtttgaaat ctattgggat tgtaaattta tgtcagtgtg 35580  
 ctgactttca agagatcttg atgatcatgt cgtctgtttt cattttctac tacatgagaa 35640  
 cattgaagcc tgaaacttaa cacaaacccg agtccccac ttgcctaaga gtcattggata 35700  
 cctaaaaagt atgctacttc ccaagttgat ttctttcagg atatgggccc ttcaaaggaa 35760  
 agcagtgagg ctgggggtttt ccaggtggaa aggtcacatt tccacatata actcagcgaa 35820  
 cattgtgttg ggttgggaga agaattggtt cactatttta aactttttgt ttcattctga 35880  
 ggacttcccc atccccctc ctccgcaaag cacaaaagta tttcctaatt ttaagtcat 35940  
 gggcttccct taatggattc tgaactcaga tcacgtccag ataagcattg tgtaatggga 36000  
 tgggtggggt tagatatttt agtcacagat gcatgagagg agggagggtg gaggacagca 36060  
 aagtttataa ctggagccta tagtagttta tctcttgtca tgggcccagg cacagagtct 36120  
 cacttcagga cagctgtgca agcagaaccc ccatcacggt tttcttgatg cctttgacag 36180  
 tcacctgtac atgcctctgg gacctttcct cctcctttct ctttttgttt tttttccctt 36240  
 ggtcacatgt ttcattctac taaatgtcta accagctctt ctctgtaaat tacagagctg 36300  
 tgatggcacc ttgcttgttg attatttctg gttgaatagt ttccaatggg acttctctgg 36360  
 agataagtcc tgtattagtc cgttctcaca ctgctaataa aaacatacct gagactgggt 36420  
 aattttataa ggaaagaggt tgactcacag ttcagcatgg ctggggagac ctgaggaaac 36480  
 ttacaatcgt ggtggaaggg gaagcaaaca tgtccttcac atggcagcag gagagagaag 36540  
 tgccgagcaa aaggggggaa agccctttat aaaaccatca gatctcatga gaactaactc 36600  
 actatcatga gaacaggatg ggggaaactg ccccatgat taaattatct ctgcctgttc 36660  
 cctcccatga catggggatt atgagaacta caattcaaga tgagaattgg tggtagcata 36720  
 gccaaaccac attaaatccc aagtgcgcct gtctggccct gatcccttta tgtgagactg 36780  
 gggcatgat cctccgcac ccgtcttctg agccctattc ctacttgggc atgcttaggc 36840  
 acttcagcat ctgcatccca ttgatgtctt aagggtggtt ccagaccttg gaggtacaca 36900  
 cgacacactg ctgatgaaaa cctagaatat agaatggaag ttacatttat tcatagagtg 36960  
 aaaatccaaa aatagaccag agagaagata tgaaaatctc aagaatgctt atcttaggga 37020  
 ggtaggatta taggtaactt ttttttctct tagataaata tatagataga tatattagt 37080

tttacagttt	ctctgccacc	aaccaaaata	tttttttcag	gaggaaaaaa	aaccccagcc	37140
agccaacata	cctaaaaacc	atctcctggg	cccagagagg	aaaaattggg	ctccttttct	37200
tgaaattgcc	atttgtgcca	ctgttggtatt	attttaccag	taactccaga	ttccaggctc	37260
ctgtatctga	gttctctctc	cttccacagt	ggagctcata	ccttcctgtt	tcctggctgc	37320
cactcagatt	taggctccgt	ttttcagacc	tcagtggctg	taatagctgt	tccttctacc	37380
tcttaggatg	gttctttctg	taatagcctt	tgtcatcaca	tcacagagg	atgatagctc	37440
ttaatgagga	tctaaaattt	gcaggtaaga	tatccctgcc	tctgacatga	gatagatgta	37500
ttgcatgcta	tttaacatac	aactatactg	agtgtgcagt	tgtatgtaaa	agcattgttc	37560
taggtattgg	gttgaaagtg	gatcaaattg	tagacaaagg	agcgtacaag	tcttgtaagg	37620
aagacagctg	ccaagagaga	agaaaggatg	gggaaatgct	gcgtctacta	agttcaagg	37680
tctgaattgg	aaagctgcag	ctattgagga	gaagagctct	ttaaaattcc	taaaggggtt	37740
ttgttatctt	ttattgatgc	aaatgctatt	ttgtggcata	aaccttaata	attttgggg	37800
tgaaactctt	atcaggataa	aatgatcctt	ttctatccca	agcttaataa	atattgttta	37860
agtacaaatt	aaatatatga	aatctgcccc	tctatattat	aatgtcata	tggcagaaat	37920
tataccttga	cttttggttc	tttcacaaaa	ccttaatttt	tttttttttt	ttttgccttc	37980
aatgaatttt	gtctgatttt	acattaaaag	cctgtaattt	ctcaagtctt	gagtctgggg	38040
agcgtctg	atcctttttt	ccctctctcc	ttgtcttctg	gatgttcaag	cgattttaat	38100
tagatgttgg	gcttttatgt	caagtgtctg	cattgcactc	catgataatc	cagggactcg	38160
gaagcacatg	ttatgctca	ccctgggttg	gtgcagtgga	actgggggtg	gttggaaagta	38220
gtattctaaa	tctgcttctt	gcgatgggg	aggtcagggt	gtcctgtgtt	gacaaggaag	38280
aagtctgggt	gaggaagcgg	gatgaaagca	gaccagacgc	tagagtccac	tttcaagtcc	38340
gatcccagga	cctggcttaa	agttaaagaa	cagcaaagat	gaaagggtgc	gcacagcagc	38400
acaggctcgg	ggccacgtta	atgacataga	aagcaagtgc	tgtgaattca	aaagaaagga	38460
cagctctgag	ccagagtact	tgggtgactt	gctcaaacaa	atccctttct	ggcaccacca	38520
ggccttccct	cccgttcaa	aaaaattctg	aattgtgcca	atccattgag	gctcagctca	38580
aggccatccc	atgcctttcc	atcgtaataa	agccttggtt	cctgggcttt	aaacatatcc	38640
cttttttctt	aggtacagat	tgaacttttt	taaaagggaa	gttgtcagag	gctctgtaaa	38700
acgttaaatc	aaacctgctt	tgttttaggg	atggggtagc	ttggaatcag	atttgctcct	38760
gctatggact	gaacatttgt	gtccccccaa	aattcctatg	ttgaagccct	aatgcacagt	38820
gttatgggtg	ttgaaggagg	gcccttggga	ggtgattaag	tttagatgag	attgtgtgag	38880
tgaagccctc	atgaatggga	ttactgtcat	ccccaaaaaga	ggtagagacc	ccagagcttc	38940
ctctctcttc	accctgtgag	gatacagcaa	gaaggaagct	ctctgcaagt	caggaagaga	39000
gagggtctct	actagaatac	acttgtactg	ccaccctgat	cttggaactc	ccctccagaa	39060
ctgtgagaaa	caaatgtgtg	ttgtttaagc	caccagctcc	ctatgatttt	attagagcag	39120
cccagactcc	attctccact	ccctggcttc	ctgcatggac	tttgcaacca	gagcttcacg	39180
gggtatagtt	taatagctgt	ttctctgtaa	cgtagccact	tttctcttct	caggtctagt	39240
tttgaccctc	ataacacttt	gttaggggag	atttgagggt	gaggaagttg	gcttgctttt	39300
cttttcacca	tgtctcagta	gaaacagaag	cagaaaggcc	ctgagatact	gagcccacct	39360
ttctcagcag	ggtgtgacag	cccggagtac	cctgggctga	ggaggccagg	gctggagggg	39420
aggctcccac	ggtggagggg	ttgaaagctg	ggttgtaatg	agctgctttt	ctgtagatgc	39480
ctaaatgatg	tgggttgaga	aatcgtgatc	ttagctttta	gtagtatatt	tttctgttta	39540
tgttaggtga	gtcatcagtc	tgtctctgac	tatgttcaga	tctggaagtt	ttctggaagg	39600

aaatttggtta ttgctgtaat agtgtagggtt gttgatctgg attagcaggg agcggcccct 39660  
 taatacatte ttaagaaaat ggtatttagt tcagtccttg gctttgaact ttgcctttga 39720  
 caaagatgaa agtgcgactt gactgggtgtt tgaaaaacat ggtgatatgg ccaggtgtgg 39780  
 tggctcatgc ctgtatccca gcacgttggg aggccgagggc gggcagatca cctgagatca 39840  
 ggagttcgac acctgacttg gtcaacgtgg tgaaacctca tctctactta aaatacaaaa 39900  
 aaattagcca ggtgtggtgg tgtgcacca taattgcagc tacttgggag gctgaggcaa 39960  
 gggaaatcact tgaacctggt aaggcggagg ttgcaatgag ccaagattgt gccattgcag 40020  
 tccagcctgg gcaacaagag cgagactcca tctcaaaaaa aaaaagcaag ttatattaca 40080  
 ttttaaaact ctatttaatg gtcagggtcat ccattccataa tgggtagagt cattgcttaa 40140  
 ttaattttaa acaatgtatt taaaagggtac ctttgttccc tagtgtcaca taacgtgaaa 40200  
 tatccaatta aggttaactgt aatgtaaagt aagtggctaa aaaagtgtg aacgccaaag 40260  
 gccagagatt caaccttttg tgtgcattag aatttcccaa ttgttcaa at ccaggttgct 40320  
 ggatctaccc cagagttttt gatccagtag gtttgggggtg ggaccaagaa tttgcatttc 40380  
 taacaagctc ccaggtgggtg ttgaggctga agctcgtgtg gggaccacat tttgagaact 40440  
 tctcccgtag actgaactca tgggtctaggt tctgtcagct gtgacctctg tgctgctgga 40500  
 gggagtgggc agatgtcctg acctctgtgc ccacagtgag gtccaagctg agtaggtttg 40560  
 accagcagct gtaatcacag agtgaacaat gtaaacgacc aatgttgggt ggtctgacat 40620  
 cttttaaaaa aaatccacgt ggatgagatc acagggttaa gtgtgggcag cagtcagggt 40680  
 aactccatgt ggttactgcc catgcactct ctgctgtttt tcacctcttc ttcagagtgt 40740  
 ggtcaggatg gtggccttgc ccagcacagg aggccctttt ccttctgacc acctgacctg 40800  
 acccacctct tagcatctgc aggcactccc tgtcccttcg ctgggccccg tggggaacta 40860  
 cttgcagtca tcaaattcat catgctgctt tcttttaatt cccacacttg ccaaggtggg 40920  
 actgccccgc atctccttcc cagtcgtgtg tcagaactca gcaactggacc tttccccctt 40980  
 cccactccc accctctctc accccgacga acgtctcact tgggatcatc tcttctgagg 41040  
 ttggacctgc acagccgccc tctgcactct cgccacctta tgggtgccc ttgacctctt 41100  
 ggcacacaga cctggaagt ggcctgtcga gctgtctct taggggtgga gcttgggttt 41160  
 ctttcatcac tgttctgcga tgaattgaat gcatgattgg tcacaggaag gtaggggagg 41220  
 gataaacacc ttatgatatg tttcttataa ggttttatat gtagaaagt atatgaaagt 41280  
 gtcagatata tatatatgaa gtatatgtga agttttatga tagttttgca taatttaaga 41340  
 ataaactctt taaaggagct gagtcccaat cccttgggtc gagagtgtcg tggctcccg 41400  
 ggctgtcttg tttccttcca ctctgctgtg tcgttgtgtg cccctcatag gctgtcccag 41460  
 acctctttga cttctctctt ttctgcccag tcttccctga gacgtctcag gctccctggc 41520  
 ctctgcttc tcggagcttc tcttgtgttt gttttctgtg ctgaggcgcc catggtgcta 41580  
 taggccacag aggaggcgtc tggggctcct cggggcagggt gcagcaggag gaagccgtct 41640  
 ccgagggcat gaccttgaa ctgagcattg acagaggaga gtcagccaga caaagaaagg 41700  
 ccaaaacccc accctctcc caccctattt ctacgtgacc atgggcccctg gacacagcaa 41760  
 gacggtgacc ccgggcctcc tattgttgcg aggagccctt gggaaaatgt tggcattttc 41820  
 ttcatagaac aggtttctct tctccagtat tcttcagtaa atcaactttc ttttttatcc 41880  
 ccaaccccag tctgattgcg aagaagtcta agcaacagaa agattttgccc aaatagatta 41940  
 tcttttttag aacaaaatag atcatgatat taataggaat tcagcactta ctcttgtcta 42000  
 agtactgttt ttaagtgtc tcaaggattt ttcattta at cccacaaca aagctgtggg 42060  
 ggggtggatgc tattattatc ggtgatttat gaatgaggaa actgacacag aggggtgggc 42120

gaggagcttg	cccatttcct	ggtagttagt	accagggctg	gcacatcag	ttgcctgctc	42180
cttttcctct	ttgcttttgt	gtccattacc	ccaaggcatt	aggatgagcc	agccaagttc	42240
tagtcctgga	ttcaccacct	aattagctct	gtgtcccatg	tcttgccgtg	gagggataaa	42300
accaattcct	agcttatccg	ttggtggtga	agatgaaatc	agtggggtag	ttgtaaagca	42360
cactgcccag	cacatagtaa	gtgcccagaa	aatgtgacgt	cggacctctt	taagcttcag	42420
tttccacatc	tgggaagaga	gggggagttg	agctaagtca	ttttccagtg	tccctttcag	42480
ctccatgttc	ctgtgagcac	tgacagtttc	cccacaatmc	tgaagaaaga	aggaaaataa	42540
gggccccgtg	gcgaaggtag	ccactgtgac	gtggctgctg	gtgggaagtc	cctggggagg	42600
caaggcccag	cttcccagac	acagccctca	ggtgtctcat	ctggtggcac	tgaccagggg	42660
ccatggtggg	cttttccacc	ccaccatgtc	tcataaaatt	acaagaacca	cagttgaaaa	42720
tcagtgttac	agaaatggta	ataggatagg	gcaaactggt	acaaagatca	gcacttaaga	42780
ttctggctga	ggcggaatat	ttgtttctct	ttagttttgt	tgtctttaat	caagaactga	42840
gagccctgac	tttcagctcc	tcaaaaaata	cagcttcctt	ccccttgtag	atgcaaaaac	42900
aaacgccact	tctttccaag	cataattttc	tcccattgct	tatctcctgt	ctacagcttt	42960
ttcttgatcc	ttctccagct	cctgtagacc	tcccatttag	agccaccagc	cggccatcac	43020
tggggctgct	cagagctctt	ggtgctctgt	gccctgggct	cggccaccca	ggcctgttct	43080
ctngcctct	tcctggttct	cttccctgga	cttcccactg	cgtgtggnc	ttcagtgtct	43140
ctctgagctg	ttgtcatgac	ctctaaccag	actgagtcag	gacttttttc	ttctctatct	43200
ctaagtcac	cttacacagc	cttgggaagt	taccctaaat	ggctattttg	ggagggagtg	43260
gggataaaga	tctgcaggcc	tcttgctcct	ggtccttggt	tctgcttata	ttggcttctg	43320
tttttaagt	tgtgtgcacc	tctttcctca	tcacaccctt	cccctccgta	tggtcccat	43380
ctcaggcaga	gttaggtgct	ctgttctgtg	tccatagctc	tttttcgagc	ccttcttctc	43440
actgtttggt	agtggccttt	catgtgtgtc	tgatccacta	ggctgtgcac	tccctgcttg	43500
ccaggatatg	gttaaagtgc	taaagaatgt	atatatgaga	tcacttttgc	ttaaaaaac	43560
cccaatcttc	tgggaattccc	aatttctaac	caattaatat	gtggattgac	tagaccttaa	43620
gcaaccaaga	gtcagccagc	cttgtcttct	atattcaggc	gcatactatc	tggtcgttag	43680
acaaaatggg	tcattatcag	tgatgagtta	ataattacct	gcacatcttg	tttatgtctg	43740
ttctttacct	aaagtggctc	ccatcaatta	aacctgtatg	gattttacct	gttcttccag	43800
aaccacccca	ctttccacaa	aaactgacaa	caatgatggt	aagaagaatg	gtagttgaca	43860
ttttattaaa	tgtttactgt	gtgcaggctt	gtttttttcc	acacatttac	ctacttaatg	43920
ctcacaataa	tcctatgaac	tagtcagttt	tatgcagatt	tcgcagatta	ggaaactaag	43980
gtggcaagtg	atcagataac	ctgtttgagg	ttgagtagct	agatcatggt	agagccaggt	44040
tcaatcccag	atacctggct	ccagggccca	tgtcttgac	cttataaacg	gctgaaattc	44100
atcttttttt	gctgaacttc	cagaacactt	tctttgtatt	tcccttattt	tggtagtctt	44160
gtacttctct	gctaccctga	ttcatacttg	gatttctagc	agcatgcctg	gcagtaggca	44220
acaacttaac	agtatttctt	tataccaaat	gaatgttgct	tttttttttt	ttttttttct	44280
tgagacagag	tctcgctctg	tcgcccaggc	tagagtgcag	tggaactatc	ttggctcact	44340
gcaagctccg	cctcccgggt	tcatgccatt	ctcctgcctc	agcctcccaa	gtagctggga	44400
ctataggcgc	ctgccatggc	gcccggctaa	ttttttgtat	tagtagagac	ggggtttcac	44460
cgtgttagcc	aggatggtct	cgatctcctg	acctcatgat	ctgcccgcct	cggcctccca	44520
aagtgtggg	attacaggtg	tgagccacca	tgcccggcca	tgaatgttgt	ctttaaaaaa	44580
ttctgttttc	ctctagctag	actgtcatat	aatgcaactg	taggaaataa	tcaggttctc	44640

095290-1-062504



tttggagtat tttccataaa agatccacag aagtcacggc aggggttgaga gtggacttgg 44700  
 gcaaatgaat ctgttcattc attgaatatt ccatgcatat ctgctgtttc ccaggcatgg 44760  
 gatatggcag ggaacacaga aatctctgcc tcctgggctc tgctttctgt tgtagtagag 44820  
 gtaaagctgc tcatactttg taaacaatat gacaacatta agtctacatg gtcattttac 44880  
 tttgtttttt tctaagaaat tttgagctgt tcgtaacaac agacgctgca gatgttaatc 44940  
 ccgttggtgt taacttttct ccagagattt aatgttcaat tttctccttt ccagaatcga 45000  
 tttatgttgt tcaaacagag gtttgagaat aactggaatt tttttaactt cttttttttt 45060  
 tttcgcacag agttcagaat tttcaagagg gatgaagaga gttataaaat gctctatggg 45120  
 gggtaacaca cagaaaaagc cagaaaattg gagaataagg atctgtctac tcgtttcctt 45180  
 ctagagctcc tctttcttac agggcactta acatgtgatt taatgtcgtg tctttaaaag 45240  
 gaggagaact gcagttcaga acttaatgtc agtgctttgt gaaagtgcaa gaaagaagcc 45300  
 ctgtattctg cacttgagag agccagatac tgggcagata ggaggtgggtg tgcacgttgc 45360  
 tttttgtctt tctcgatcat ggcattgatt ctgttcataa caatgatgca atgtcatcct 45420  
 cttccccaca catttggtgtg cagatagaaa gaatgcaaca gcacagagtt gttggggaat 45480  
 aatttggcat ctaaaatata gacataccag catagatcat atttatgact ctggtgggag 45540  
 tgtcacagca atgatttaat aggaggcagt tgtctccaag gcctcctgaa ttatgactgg 45600  
 ttttaaaatt cttagaacct attggaggtt attgtttctg aaaggctaca taatttaagt 45660  
 gctccacatc cgtcattata ggagatgtca gaatagtaaa atctaatact ggactaagtt 45720  
 gttatcgcag ccctttgggt tgggtggcttt gccgacttta taaatatgcc tgtcagtgcc 45780  
 tgtggtctct acagttgggc agtcggcggt gaatatcatt tctcacattt tacactgggg 45840  
 gactggaacc cagaaggcat atgttttccc aagaggcacc aacacagttg gcccatgagg 45900  
 tagagcagcc cctccttcgg ctcagcctcc gctgcactga gccaagccaa gcttcctaca 45960  
 ctggcctctg tgcagctgtc tctcagcaag aatgcaagtc ggggagagaa gccggatccc 46020  
 tgggattggt ctagagagta gaaacctcag agtagcctc cttagaccac ctaacgcatt 46080  
 gcatcgctgc atacatgtaa gggactcaat gctggtagga ttggcttagg aatgatgcaa 46140  
 gtgaaaacag tgcctcgggt tatcattaga acaagggtct tagctgacag ttgcctcaga 46200  
 ctttgatttt gttctccttg acctgccact ccactcgagt ccacatctct caagactgca 46260  
 cacgcctgaa ggaggactga ttacaaacca aagccttggtg cccagtctgg atctttttgc 46320  
 attgttgaga aagcagctta ctttcttttg actgattcag caggccaaat ttagaacaaa 46380  
 gatttttaac tatctcctt tataaattac tgagctatct ttagccagg ctactcttaa 46440  
 tatgaacaaa aaatattata caaatttggt gttaatcgta aactataaaa aaatcagtaa 46500  
 ttgttaccac gtgaaatgaa tttggataaa agagatacgt ttttgccct tcccagggtt 46560  
 taggagagac gaaatggtga gatttttagct ctgaatcaga gggtcttatt agaggtgggt 46620  
 ttgttcctcc tgacccttag gggatattta gcaatgccta gaggcattga tgggtggcag 46680  
 atgtactat gccctctgct aaacattcta cagtgtataa aactgttcct cctgacaaag 46740  
 aatcatccag ccccaaatg tcagtagtgc tgaggttagg aaacctcct ttaaactctt 46800  
 ggggtttatt gctgacctt acagtggatc agcttttatt tagttcatgt agaggtgaaa 46860  
 ttaatactag tgetcaaata tgtctttgta ttctggactt ggcctggatc ccccgaccaa 46920  
 atttgggaca agctcctgcc atgtgttagg gacctgaatt caggcagcta acaacagtat 46980  
 ttgaactgtg ttttcagtgg tgggagtga gggagatgag cgacgtgcta gcaagcgcac 47040  
 aggggttgcac gaggaaatag agagtaaagc tgcagcgtgg agccctgcta ttcagagtgt 47100  
 gcttgagaaa acagcagtgagg aggcattact ggggagcttg atggaaatgc tccctcaga 47160

cttgctgaat caaaatcttt aatttagcaa gatccccagt gaggcttggt catgtagaag 47220  
 ttagagaagc acggggtaaa ctcttctttt ttttactttg gaggaaaata cacctttttt 47280  
 cttattatgg ctctgacct tactagctgt gtgaccttg ccaagttata aaacctcact 47340  
 gcaccttatt tgttttagct ggaaaatgga gatcataata tcacctgtcc tatgagattg 47400  
 ttgtaagaat caaacaagct tatttatgcc aagaacccat atggtaaaaag ctcaacaaac 47460  
 tgtcactagt gataataaga aaaagatcac aaaagtagaa aacattaggg agacagctta 47520  
 ggtcttaaat ctacagttg tcgtcccaa acaatacttg ttttttgca gatccagttt 47580  
 ctctgaatac taaaataaaa ccggagtttc ataaacttct atagacagtg gtccttgca 47640  
 gtagcccaag tggcagagag tacatggatc tggggacaaa cagcctctac tgtaggaat 47700  
 gttccatcct cctggcctga gttacacctg ctcatgtga ttccgaattt gaaaggaaca 47760  
 cagtaggaat tttcaagacc ctgggaagag gaaggctgtg gtaaacagga aggatgagat 47820  
 tagaagaagg agtttaggtg aggtgagccc ttgttttact agtaggggtt aagaatatcc 47880  
 aagtcagctg gacatggtg ctacacctg taattctagc actttgggag gccgaggtgg 47940  
 gcagatcacc tgaggtcagg agttcgagac cagtctggcc aacatggtga aaccccgctc 48000  
 caactgaaaa tacagaaatt agcagggcat ggtggcgcat gcctgtaatt ccaactactc 48060  
 actcgggagt ttgaggcagg agaatcgctt gaacttggga ggtagaggtt gcagtgagcc 48120  
 aagattgggc caccacactc ccacctgggc aacagaatga gattccgtct ccaaaaaaaaa 48180  
 aaaaaaaga aaaaaaaaaa aagaatatcc aggtcaaccc cacctaacc tcagcggggc 48240  
 tcccttctgt tgctgggtg ggtcctgggt tctcttgacg cacacgagat tgtgagagt 48300  
 tatggaaaca ctgccctgc tatcaggaca gcgctgcca tgccagccag aacacatcat 48360  
 aggaattgca aaactctttt gcaaaccagt gagagatatg cttccaatgt gaggtaaagc 48420  
 agaactttaa tcacagctgc agtggtccac agaattccaa gagccaagat ggtaaaagaa 48480  
 taaaaaaaaa gaaaggaaag ggctcaaatt aaagacttca agctgcagaa taagattaaa 48540  
 taaaaggatt caattgaact gcatcatatt cagtaatgac taatcctaag tatacagggt 48600  
 ttgggggtga aaggatttgt aagtgtttt caggaaaata ttttttccat ctttcatttt 48660  
 aattagaata gatttgcatt attttttctt agtttttatt tttaaaatat ttattgccac 48720  
 aaatttagaa aatacaggga aaacataaat aacagtacat gtaaaccaat attttgtccc 48780  
 ttcttttggt caacagctat ttctcaggca cctgctgggt gtcagcagct gtgctcagt 48840  
 tgggtgacca aaccttggtc aacaaggcag caaggttcta acctgggttag ggcttacagt 48900  
 tgagtagctg aaattttgat ttcttttctg tgccctagt aaagatatga tagcaaaca 48960  
 taagagctat tttttttatt gtgttcttac tctgtgttg gccctgttct cagtgggtta 49020  
 tagcctatta actcagctc tttaccacca ctctgagggg aggtctctgtc ataccactt 49080  
 gacagatcgg gaagtggaag catcaggagg ttaagcaact tgtaaagat caaaaaatca 49140  
 ataattgacag agttttgatt agaatccag cagcctgtct ccagaacctg ccctattaag 49200  
 tgagtgcaa ctgtactgcc ttctataata tgtatcaaat tgagatgata ctttataatt 49260  
 tcaattcttg cttttctatt gaacagtaca cagtaacatc ctctataat gcatataaac 49320  
 cccaaaaga tgtagaattt taatttattc atttgtctga taggtcata atgaaataag 49380  
 actctataaa gctgtgtaatt ttagatatag gaaacatttg gattatagt gtatgtagtg 49440  
 ggaacaaatg gtcttctgaa tcagggaagac atgagttaga gtatgccgggt gtacctcct 49500  
 actcactgta tgacctggg caagtttctg aactttagtt tcctttccag gctaatatct 49560  
 gccttctgga cttgtcatca ggattaaatg agtctaccta tataaatgc ccagcgcagt 49620  
 gccagcacg tggtagaagg tctgctagt gttactgtta ctgctggcta ttaatacat 49680

tttaatcttc cttcagaata cctggccaga tagcacagtg gttaagaatg cacatgaaag 49740  
 ccagactgtt ggggtccagt cctggctcga ctcccttcta gctatgtgac attaggcaac 49800  
 ttacataaac tccttgttcc tcagtttgca tttctttaaa actgcatagt tatcataccc 49860  
 atgtcttaga gttttgtgag tgtaaattat tgtatataaa gctctgagaa cagtttggtgta 49920  
 cacagtaggc actgtatgaa ctttttctgt aattatcaat aatataatta tttaaataaca 49980  
 ttttcagaag gagataaaaa tattacacct taaaaagcag gtatctttaa attcttcctc 50040  
 agctactgaa gttttgctta ctatttgaca tatcatttgt ttcacgtttg tggctcagac 50100  
 gtggcttatg ccaatgcata ttaacacagg aatttttaaat ttggtgatat tattatatatt 50160  
 tatctgaatg aacagaattt gctgatttga cactgtgttt gaatgtgcat tttttgttga 50220  
 aaaatgacaa ttctggaatg cctctccct ttccagatta ttcagagctg ggagagcttc 50280  
 cccacgac tccttttagaa ccagtttgtg aagatgggcc ctttggcccc ccaccagagg 50340  
 aaaagaaaag gacatctcgt gagctccgag agctgtggca aaaggctatt cttcaacaga 50400  
 tactgctgct tagaatggag aaggaaaatc agaagctcca aggttggttt gccatcttga 50460  
 tattgaacag gcctgggtctt atcttggtc tgaagttaat cacatcagac ataagcatgc 50520  
 tgtcttaaaa atacagcagc acgatatgtc aatgtataca tctatctata tctgtttact 50580  
 ttttcagagt aatattaaca ctgtttactt tctggtgatc taatgatagt ttcaccaaca 50640  
 atattcatta ttctctatg gtcactgtta gtacagtgtt tagaacttct gagatccaag 50700  
 ctttaaatct aagctctaac acgctgaaag gtgcttttca ttttgttttg ttttccctc 50760  
 tgtctctctc tctctctcta ctttatcctc agccatgggc tgtgcctgtg tgttaggtat 50820  
 gaacttttct tgtgtaagtc attaacatac gtaacttcac tctgtgtgct ttttcagtga 50880  
 tttgcaagta atctgaaaaa aaagaattag ctgagttcta cctgtactga tatcaatagt 50940  
 gtcaaaatat gacatgaact ttgaaagttt agattttgtt catttcctgt ttccatgctg 51000  
 aactggaac caattaatgt tatcttcaaa gtagcttaag atgcaaagtt tacatactct 51060  
 ttggaaagag catgagtctt agggtatcta gagaactgcc cgggtgataaa gtagtgaaga 51120  
 ttttgagcag gaagtctgca taatctctt caaagggag atgtagcaga tggttcagtc 51180  
 accctgccat tgcccagaa caattttgga attacagtac atttcattca gcatcattct 51240  
 tgattgcaaa ttttgatctt ttaaaatgac cttgatgctt gtatagagct aaaaagtcatt 51300  
 taagacacca actctgagga ataagctcct gagaatgtgt tgcattctgt agtttcagtt 51360  
 gcatagctag tgcatacg agtgataga cgttctctgt gcatgtccct acaatgcttg 51420  
 tgagttatga caacactgtg tacgagcaac atagtttctg cagttgaaaa gtacgaattc 51480  
 atagaatgta aagagatagt gtctatatct tttgactgaa aacagaaaat gagatataaa 51540  
 ggaataagac ctttcgacat gaaagtaacc ccacagttgg aataggctag taagctttcc 51600  
 aacatgcagt tttgaagctg agaaagacgg gtccctctcat cagggtgctg tggagatga 51660  
 tagcacactg gggggcggtt agagcaggtg agtgctgttt tcttccaacc cagtttttct 51720  
 gccactttct tatgtttttg tgaaggtaat tttaaaagca gatgtctaaa agatgtttgg 51780  
 tagtgatggc attactgcat gtgtcatcag ttaaatgaca gctcgggagc acagcagtta 51840  
 tgttcgtgtg tatcttggga tttttgttga agaggaaaaa ggcagttatg ttcattcatgt 51900  
 aggtcaaaact ttaatgccaa tactggccaa tattcttgca aatgacagcc atgtaaaatc 51960  
 agggcatagc tataaaatgg gaacggtgct cacagctggc ttctttgtgg tgaggacagc 52020  
 tataattggt gaggcaaaac cagtgtgcca caaaagcaga atacattctg ctgtgcaagc 52080  
 aatgaccaga cagactagaa tgaaaaggca agagtttcct aaggttacct ggaacccctt 52140  
 gccaggtgtt gcattaagtt tactggccct tgccaacatt cttctaattgc ttctcattt 52200

catctggctt	cttggcagtg	ttcagttttt	gtgggtctttt	atttttactg	tttgacttca	52260
tttctcttct	tagctctgta	aagttccaca	tgtgtttatc	tttgtggtga	aaacacaata	52320
aacttgctta	atataatggt	ggaagtatta	atccattgta	ttagtgtgta	caggacctgg	52380
attgctgata	aaaaaataac	tagcaataac	agcctgattg	cttaaaaaata	tttagtaagt	52440
tttgtcgggg	tggattgggg	cagggcagaa	cttttacatt	aaatatagat	gcaagatttg	52500
ataagaatca	gccagagtg	acagtaagta	ttcacttaat	gttgccaata	ggttcatgga	52560
aactgcgaat	ttaagcaaaa	tgatgtataa	tgaacaaaat	tttactaagg	gtttattgat	52620
aaaaacaaga	gttaagttcc	tatggcatat	ttctgggcac	aaaaacatca	ccaaacttct	52680
aaataaagac	ccaagacact	tctaataatta	aatattgatg	taaacgtgag	atatgcaaac	52740
atttaagcaa	gattaataca	aatatgataa	ttattggctt	ggcacagtgg	ctcactcctg	52800
taatcccagc	actttgggag	gctgagacag	ggagatcacc	tgagggtcagg	ggttcgagac	52860
cagcctggcc	aatgtggtga	aaccctgtct	ctactataat	tacaaaaaaa	aattagccag	52920
gtgtggttgt	gcacacctgt	aatcccagct	acttggaaga	gtgaggcagg	agaatcgctt	52980
gaaccagga	gacgaaggtt	gcagtgaacc	aagatggtgc	cactgcactc	cagcctgggc	53040
aacacagtga	gactccatct	taaaaaaaaa	aaaaaaaaaga	aagaagtaat	tatttttcca	53100
cttattccac	ttcagggctt	cagggggcca	gaacctatcc	ctacagcttg	ggatgcaagg	53160
caggaaccag	ccctggaccg	aatgccattc	catcttgggg	tgactcacac	acacactcag	53220
actgggacca	tgtagacata	ctgattaacc	taatgtgcac	atctttgaga	tgtgggagga	53280
aactggagca	cttggagaaa	acccacacag	acatgaagag	aacacaaaact	ccacacagat	53340
aatggccccg	ggctaagaat	ccattttttt	cttgtcaaca	ttataagaaa	gcgacattga	53400
gcataaagac	attatttgag	gacctgctgt	actatgtact	tagagagata	ggcattctat	53460
cttgagttcc	ttttttttct	cccttcttga	aggaagggtta	aattgcatct	gagatggctc	53520
ttgaaattga	tcaggggttc	aagctgactt	gcatactctt	tgggaaagaa	tttagaagga	53580
tgtgtatgag	gaagttctta	tgggttaagcc	tgtttcctga	cttgaataga	tgaatcaaat	53640
attttttact	attctggaag	catcgcattc	tggaaagaac	catactatgt	catctcagtc	53700
tacctcactc	cattgtaggc	acttggaagc	tgaagttgtg	atttctccaa	aattagatag	53760
ctaattttca	ttgggtgttg	aacaaaaaagc	gctgcctctc	tttgaagaca	ccagtcctcc	53820
accgtcctcc	tctgcaaggc	cgttttcccc	cccctttttt	ttttttttga	gacagagttt	53880
tgtctcttgt	gcttaggcta	cagtacagtg	gcacaatctc	ggctcactgc	aacctccgcc	53940
tcttgggttc	aagcgattct	cctgtcttag	cctccagagt	agattacagg	caccaccac	54000
cacacccggc	taatttttat	tattagtagt	agtagtagta	gtagagatgg	ggtttcacca	54060
tgttggccag	aatgggtctg	aactcctgac	ctcaggtaat	cctccacact	tggcctccca	54120
aagttctggg	attacaggca	tgagccactg	tgtccagcca	atttttctgt	atttttaaat	54180
gaagatgtga	gcagccta	gtaagatcac	aacatgtgat	tcaatacagc	cgtggcttgg	54240
tgttgacatg	ttattaccag	ttgagcta	ccatgtaact	cagcatttta	tgctttacta	54300
agattaaaat	gatgtgataa	cattaaaattt	tgaattacag	ttgatgtttt	ttatttaaaa	54360
aacatttttc	ttagttaaat	aatacatgat	ggtttaaaaa	tcaaatattc	agtgcatttc	54420
ttctaaaatc	tctgcaagtg	tgggggtcat	ttaattgctg	agcctcccag	cctattagct	54480
ttccattctg	agctttcaag	agatggtggc	agctggcaag	gcagttttgt	ctgggaaagc	54540
cattgttaac	agagcagaat	tgggggatgga	gcagccatag	cccaccacc	agagtaggca	54600
caaatcagac	ctgaacgtta	tcacaaaagtc	caagttggct	cagacatttg	tgtaaataca	54660
taataaatat	tttagagaac	ttgggtgcaa	atttacattt	gatctcagtc	agtcctcttc	54720

ccctatctct	acaagcttac	aaaccgcatg	gggtgtgtggg	ggctcttattt	aatattgcga	54780
acagctgggt	cctgtatctg	aagttcttgc	cctggagcct	gggtgtttgt	tgtagtctctg	54840
cacatctgc	cttgggtgat	aaggcatttt	ggaggccact	gatttttaggc	agcagtgttg	54900
ttaggatacg	gaaacagcag	gatgtttgtg	gattgagcct	tttcagctga	atcttctggc	54960
cagttctttc	tggctgtgtg	aagttgtgtc	gactacagag	caggatgctc	atgttgccctg	55020
ctgggctctg	ttaggggtggc	cagacgtgct	tgtagcagcc	ttactgccag	aggaacgtac	55080
gttggcatcc	agagtccagt	gctgccgcca	gttgcagtgc	agcaaggcta	gccccaaacc	55140
tgatttgctg	caaggattag	ctcaactcta	gtgacattta	ttgtgttttc	tcatagccca	55200
aatcacagcc	aaaaaaaaaa	aaaaaaaaat	ctagggttga	cattttttaa	aattttttta	55260
aaaaacattt	ttcttggtta	aataatacat	gatgggttaa	aaatcaaata	ttcattgcag	55320
ttctaaaatc	tctgcaagtg	tgggggtcat	ttaattgctg	agcctcccag	cctcttagct	55380
aaaaaatcta	gggttgacat	ttttaaaaat	gtattcaaca	gagtacgagg	gaaaagatta	55440
aagatggtgg	atggaaaacc	ataaaagctg	agaggaaggc	agcactgggc	ttagagtcac	55500
ttggcttccc	tctagctagt	aaataaccag	caccaaatac	cctgatcctc	ctgaacttca	55560
gtttctgtgg	ccatgaaata	agagggtggg	tccaggaatc	aatgtaaatt	gtcaatttaa	55620
catttccttt	tattgatatt	actccccctt	gggcttgata	atttagttat	aattcttcat	55680
gcagctttag	gttgagtaag	tttgggtggg	aacagtagct	ctcttcatat	atttgagaga	55740
tgtcatttga	aaggggtaga	tttattcagt	ttaactccaa	gaagcagaaa	tgggacctat	55800
ggtagaagct	accaaatgga	ggtttggctc	taaataagaa	aacgatcttt	ggagtgcctc	55860
tcctagttta	gatgaaaaaa	attgcatcaa	gttgtaacca	tgctagtcat	tgggaattttt	55920
attaacaaca	cgtagctcct	gtcctgggga	ggctcatagt	ttgatagggg	taagatggaa	55980
agaattgggc	agatgtggat	tatgtcttag	cagtagagcc	aacagagtat	gttgggggtg	56040
aaggggtaag	agaaatcaca	tacctcctag	gttttttagca	ttttccaaaa	tgaggaaaat	56100
gggtagaggc	atggacagtg	acttatattt	agacgcgtta	agccagtgtg	aactgcttga	56160
cgtctcagcg	ggataacaag	taggcagcca	tgttgtgtaa	tggaaattcc	atagctgtag	56220
cctttactaa	tgcgctctgg	aatgggtctat	tccagcctct	gaaaatgatt	tgctgaacaa	56280
gcgcctgaag	ctcgattatg	aagaaattac	tcctgtctct	aaagaagtaa	ctacagtgtg	56340
ggaaaagatg	cttagcactc	caggaagatc	aaaaattaag	tttgacatgg	aaaaaatgca	56400
ctcggtgtgt	gggcaaggta	agcttcattg	ggaagcatct	agtcaacctc	accttcattt	56460
ggtgattggg	gagaagtgtg	gaattaaaaa	aaagtcaagt	ctaatttttag	tggccatctc	56520
ccttcttttc	atcacatctt	aatctatttc	catatacctt	acttaataga	catgagtttc	56580
accacctttc	atgattcctc	ttaattaaaa	ttcccagaag	gccgggaaat	aggaagaaga	56640
cagaaaaacc	caagggtttt	gttgccctata	aactagataa	tgatttgatg	atatactttg	56700
aattaaatta	taaactagaa	actaattgta	tggcttgtct	ctgggtactc	tagggagaca	56760
acatagtgtg	gggagcacag	acttcagaca	gggtgtcttg	ggcttaaatc	tcaggcctgc	56820
cacttacttt	gcagtgtgat	cttagacaaa	tgcctcacc	tctctgagct	ccagtttcta	56880
caagtgtgaa	atgtgggtgc	tgacagtgga	tgttgtgagg	agcacacagc	atgtgtctgc	56940
tatactgtaa	ggccttagag	agcgggcagg	attcactgtt	ttttcagtga	gatctgccag	57000
cccaaactgt	tactgggtcca	agaagagata	agtacagaac	ttgaaactaa	gctttttggaa	57060
atgtttccag	caatgtgaca	cagtgtatct	aattaaaaat	gtggacttat	attttgtcca	57120
tctgtttttt	tttaaatttt	gtttttctac	taattttatt	ttactgtatc	gtataaaaaat	57180
atcagcctgt	agtagattgg	aaaattttta	aaaagaaaaa	aaattgatgc	ttcacagata	57240

gtttgagaac	cgctattttg	aagcttacct	tcagtcatta	ttagtgttct	agtcaaacaa	57300
tgatttcttt	aaaaatatat	gttaatgtct	tctggcaaga	gtaaaagcct	gagtctaatac	57360
tgattctatg	ctactgagtt	ctgggtgagc	tcacatgaa	taaccagggtg	ttctgaataa	57420
gggtttcaag	tatgtataga	atgggttttt	cctgagttta	tcagttgtgc	agtgggaaaa	57480
cgttgatat	gcactttttc	ttttttgaga	tgtagtttca	ctcttggtgc	ccaggctgga	57540
gtgcaatggc	gcgatctcag	ctcactgcaa	cccctgcctc	ccaggtttaa	gctattctcc	57600
tgcctcagcc	tctgactag	ctgggattac	aggcgccccg	caccatgcct	ggctaatttt	57660
ttgtgggttt	tttttttttt	ttttaagaca	gagtccttgct	ctgtcgccca	ggctggaatg	57720
cagtggcgtg	atctcagctc	actgcaagct	ctgcctcccc	ggttcacacc	attctcctgc	57780
ctcagcctcc	tgagtagctg	ggactacagg	tgcctgcacc	catgccccgg	ttaaatttttt	57840
ttgtattttt	agtagagatg	gggttttact	atggttagcca	ggatgggtctc	gatctcctga	57900
cctcatgatc	cacccacctt	ggcctcccaa	agtgtctggga	ttacagggtgt	gagccaccgt	57960
gccccgccaa	ttttttgtgt	ttttactaga	gacgggtttt	cactgtgttg	gcaaggctgg	58020
tcttgaactc	tggacctcag	gtgatctgcc	tgcctcgccc	tcccaaagtg	ctgggattac	58080
agatgtgagc	cactgcaccc	ggcctgcata	tgcatttttc	atctctagga	gcataaatgg	58140
aacaaagcag	tgttttttac	tatagttttt	taggcatttt	taaccttttc	tgaattttga	58200
catcaatttt	agtaatcatg	ggaagtattt	gtttgttacg	cattttccct	ttctatggat	58260
aaggaaactt	gggcttagag	cagttgaata	gtggcttagg	gccacagagc	tgggttcaca	58320
ccaccgtact	gcactgcctc	ctgttgaaca	ggatctccag	gtgcttatct	cagaacacgt	58380
atgcagtgg	gaagaccgaa	gttctggatg	gacaccagct	ttcagtgtga	ctttagcagg	58440
tacctctttt	ctgggctctt	gcccccttac	tgatagaagg	agagacttgc	actgagtaga	58500
ggatcttgg	gctgtcttgg	agttctaata	ttccttgcac	ctgtactttt	tcttgagggt	58560
tacctttaca	ccaaatgacc	ccaaattgct	gttttgaaaa	gggagaaagc	agagaaaaga	58620
atgagtctgt	tcttccccca	ttcacagttg	cctagatgat	caccttcagg	tgtctttgct	58680
tctgcgaaag	gcaaattgca	tgggtctgtg	acagctattc	caaataattg	agcttcttag	58740
aagcctggca	cctggatatt	tgtttttcac	tgggcatatt	ttgtgggggc	taatagaaat	58800
actctaggaa	tctggacctt	gggtagtga	agttgggcac	agatgattga	gcattctgta	58860
tactggagtg	agctaaggct	gacctggaat	ttccttatgt	gttgctgac	tttgccacat	58920
cactttttac	tgcagaagct	ctaaccataa	agggggcttt	gtcagtcagg	tggttttaac	58980
acattaagat	ttaacaactc	caaacaaatg	agggcggtct	attttgtgg	tcagaataaa	59040
aatgtgaatc	aaaaaatttg	agcctaaatt	tgaatcatat	ctttgacctt	tgaagtagag	59100
gccaactcac	ctcagagacc	ttgtaagaga	ggacagttgt	gtggattaag	aggcccttcc	59160
tcatagtga	ataaaagacc	ctgaagtgat	ggaaataaag	gaatttataa	aattttccca	59220
gttaaaatta	gatgaggggc	caggattagg	gtatcaattt	aggagaagat	aacataatcc	59280
tatgacatta	tagataattc	agtttagtac	acatcaaaat	gatttctcta	aagatatcta	59340
gatagaacct	tataagctgg	aatgtctttt	ttaggaatgg	gattgcagag	gggctgcctg	59400
ggctgctgac	agtagggg	agatgcaaac	tctgcttgcc	tttgaccgg	caatgccatt	59460
tataaaaact	tactctagaa	actaatcagc	caaaaatgta	ctgcagtaag	aatgcttatt	59520
gtgacattgt	ttaatagtca	aaacaaaaca	aaaaacccaa	catgtgacta	tcccatgtca	59580
tattccttga	aaatgacacc	ataagtagat	ctgtattttac	tgacgtaaaa	gatgtctaag	59640
ttgttaaagt	aaaagagtac	agcatgggtct	cctgtactgt	tgatatttcc	atgtgcgtat	59700
acatggaaaa	acaccacga	tgcagatgtc	caggttatag	acaggatgac	catagggccc	59760

aacctggcat agccctgggt tatgactcct gtcctggcaa aattattaat agccttcctc 59820  
 ctcttcactg tcaaaagctt cctgctttgg atggtaaata tatgcttatt ctagtattgg 59880  
 gtgggttttc actttcttct ttatacctct tgcatttcag aggttttttt gcaccacttt 59940  
 taaacagtga gtgtatatta ttttttaagt gagtaagaag ctatttacat gggggatgga 60000  
 ggaatggcct cctgccctcc cagaccctgc ctgcaagccg taggtgggct ccactgccag 60060  
 gtttctcttg ggtaggagt gaaggcagca ccatgggtgg gaagggcatt ccaggccatt 60120  
 cttagcaaaa acattgggtc caacctgcat gatcctgtgc tttaaatac agaatacctag 60180  
 cttactcctg aataccacaa tatctggtac tgtccagtga cacagccaat attcttttct 60240  
 ttcaaaaaat aaaggctctga taagacaatg ggaatgattt agtaatagga aattggacat 60300  
 ttcataactt gggaaaattt cccagtttga gaaaaagtat tttgtgaaaa aaagccccac 60360  
 tataaatcac ttatcatgct gactgttttc tagcccatat ttacttctca tcagcatttg 60420  
 aagtatttgt ggggaggggtg tgcgtgtgtg tgtatgtacc caggatatat ctatgagctg 60480  
 gaatagcaga gggagacaaag aaatagaata atagtagaaa gcagagatca gggatatatt 60540  
 gcttcctgtt gctaccataa caagttacta caaaaatagt gactaaagca acagaaattc 60600  
 ttctctcagt gttctggagg ccatagctcc aaaaccatgc cgtagcttg tctgtacttg 60660  
 gcctcttcca gcttctgggt tctgtcagct tcttagactt gtggtcacgg cactccagcc 60720  
 tctgcctcct tggtcacatt gattccccct ctcatctcct cctctgtatg tctattataa 60780  
 gaatgcttgt cactggatat agggcccatc tggataatcc aggatgctct cctcctccca 60840  
 aagtccttac ttaattatat ctgcaaagaa ggtaacattc acaagctcca gggattagga 60900  
 agtgaacaca tctcttttga ggggacacca ttcaactcac tctacagggt cattatatta 60960  
 atgctgagat aaaattacag aaggtagatg atgtgggtcat gggttacagg ggcctgtat 61020  
 ttcttctaca ggccaactta aaaaaaatga tacgtgaaag ggaaagaaga aagtacttac 61080  
 tacacagtaa gtatttccaa gaggtggccc agtgagactt ttgaatctgt taataaaatg 61140  
 attactattt gggtcaaact cacagatgggt tattttatca ttaattgcaa gataggaaca 61200  
 caaaaatatt tttctctagt cccatttga gtagcagcct tgtttgacat ttctgacatg 61260  
 gaggacacca agagaaaatg gcagtcagca tccctgggct gtcactcacc ggcctaataga 61320  
 cctagggcaa gggacctgtt ctcactgctt ctcttttctt taccatgagg ataactatgt 61380  
 ttcccttaga gggttatgag tatggtagtg gccaatcac ataactgca tggaaatggcg 61440  
 atgggtgcata gtggcctcgc aatcagtgct atctgctgct gctacctgcc agagcagaaa 61500  
 cttttcccaa aggtggccag agacagaaac cagagaaacc atccttctgg acaggctgtc 61560  
 tgagtggcag ggcagggtac aaagcggcca ctttttttcc cggatggaaa gaaagatcaa 61620  
 tgcctaactt ggaggettcc tttctcccaa aagacaagaa agacttggca tcttattctt 61680  
 cagtcttctt gctctccccc tttccacctt tttggccttg taatagctga gtaatgagct 61740  
 aaagaatttt ggttcaaaac tgtcaccttt taaaattagg tttgccctaa ataacatcct 61800  
 tgactttaag agaattttct taagtttttag acatttttaa tcaactgtgag tattcaaatt 61860  
 aatcacatgc aaagcattag ttagaggctc ttggacattt tctgttttta gagctttgtt 61920  
 ggatgctcac atggcaatgt ctgtgcagtc agttcctacc cagcctctgg gctcttcttg 61980  
 cagcttatct tgcagaaaga agcctcatca gaattccaga atctcagcta tgattagctt 62040  
 actccacctc agctcagaaa catgcatgat tccctggagc taccaaactg ggggcagggt 62100  
 tcttgccgtc aattttgcct ctcaacaata ccttccagc cttcttgcca gctgctctct 62160  
 tccacatgca cccttggtgcc tgaggcaaac tgaatcactc tcggttccct ctctcttgta 62220  
 cttttctctt ccttttccct catccttaag gctcggctca aatgaaggat tctgtggaac 62280

cttgattgct cagttagaaa tgagcaaaact gtcgcaagga cagaaaaacca aacatcgcat 62340  
 gttctcactc ataggtggga attgaacaat gagaacactt ggacacagga aggggaacat 62400  
 caaacactgg ggcctatcgt ggggtggggg gagcggggag ggatagcatt aggagatata 62460  
 cctaattgta aatgacgagt taatgggtgc agcacaccaa catggcacac ttatacatat 62520  
 gtaacaaacc tgcacgttgt gcacatgtac cctaaaattt aaagtataat aataataata 62580  
 ataaaaagaa atgagccagc ttctctttca tctgagctct acttcctttt gattctctct 62640  
 gctttctgag atcacatctt acatgacaat ttttcatact tggtctttatt tccctagaat 62700  
 gttgttaatt ggcaccaggt tggagctcag gtcgtatact ttattccttg cagagtctga 62760  
 cagggtcaga acatgataac acatttgaga agtgagaaga agggaggaag gggccaggga 62820  
 agtgagggga gaataggggg tggagtagg ggaagaagca aatagggcaa ggttttagtt 62880  
 gcctcccttc tgttcttatg ctgttaatta ataatggaac cagtggccag gcatgatggc 62940  
 tcatccctgt aatcccagga ctttggaggc tgaggcagga gtatcgcttg agcccaggag 63000  
 tttgagacaa gcctggacag catagtgaga cctgtctct acaaaaataa aaaaaaaatt 63060  
 agccaggcat ggtggtgggc acctataatt tcagctactt gggaggccga ggtgggagga 63120  
 tcattggagc ccacaagggt gaggtgcag tgagatgtga ttgtgcctct gactgcagc 63180  
 tgggtgaca aagccagact ctgtctcaa aaaaaaaaaa aaggaacaag aatttgata 63240  
 aatggaacat gaaacacaat tcatttttat tattaagttg tattctgtgc ataaattatt 63300  
 tccatgtctt ctctcccttt taaagggtg ccacgtcatc accgaggtga aatctgaaa 63360  
 tttctagctg agcaattcca ccttaaacac cagtttccca gcaaacagca gccaaaggat 63420  
 gtgccatata aagaactctt aaagcagctg acttcccagc agcatgcat tcttattgac 63480  
 cttggtaagt ctgtgccatc gattggagat gacaatggaa gtttactca catgaaaaat 63540  
 ctgaagagac tgtccaagtt atgtattgac ctgcctttag gtttagcaat caaaatttac 63600  
 tactgagact tttaatttaa aaagccctag ggtaatcaca aatgtcatct tcaagcatat 63660  
 aaaaatctct gtattttcac tggggagctt gttaactttg cttggcatgg agggagggtg 63720  
 ttcattaagg ctgcagtcac aattgtggtt cagtccagta actcaaatat tgataggagg 63780  
 tttttacagt caacegaagg aacatcctgg aaaacgtata gatgttcaga accgaggctt 63840  
 ggtttaatta caggagccac tcctcgttt ttactgtc caaacagaaat tcacagaaa 63900  
 aattgtagaa agcagtttgt gtgtgtgctt tgaatgattt tattttggaa actgggtggc 63960  
 acctgtctc ttgaatagtt tttaaaataa gaagatggga acaatataca gtcagccctc 64020  
 catatctatg ggttctgaat ttggggactc aaccaacctc agatggaaaag tatttgggaa 64080  
 gaaaaatcaa tgaaaactaa acaataatat agattttaaa atatagtaac tatctatgta 64140  
 gaatttacat tgtattaggt gttataggta atctagagat gatttaaggt gtgtgggagg 64200  
 atgtggccgg gcacagtggc tcacgcctgt aatctcagca ctttgggagg ccaaggctgg 64260  
 tggatcatga agtcaggaga tcgagaccat cctggctgac acggtgaaac cctgtctcta 64320  
 ctaaaaaatac aaaaaaatta gccaggcatg gtggtgggag cctatagtcc cagctactca 64380  
 ggaggccgag gcaggagaat ggcgtgaacc caggaggcgg agcttgagc gagccaagat 64440  
 catgccacta cgctccagcc tgggtgacag aatgagactc tgtctcaaaa aaaaaaaaaa 64500  
 aaagtgtatg ggaggatgtg tgtaggttat gtgcaaacat agcaccatct tatagaaggg 64560  
 ccttgagcac cgtggatttt ggtgntctgt ggggactcct gcaacctatc ccccgaggat 64620  
 gccaaaggat gactgtattg gatagatttg cagttgccac tgtgaaggac ttgttgaact 64680  
 ggggtgtgat tatgatgcac agagggccct cctgacttgt cagtggccat gcacagggcc 64740  
 aggtggcaat gactccctgt ttgcctgccg cctatcacc aagctgctgt ctctactggt 64800



ggtgagctgg ctgatgtgg taggagatgg gccctgctgc ttttagagca tgtggccctg 64860  
 cttccagaat acctgttctg gttgcagctg ctgctgctga aggctccaca gaacacacag 64920  
 tgctttgggg cctgcggtg gcccggttct ctgattgttc ctgcagccac gacagaggat 64980  
 gcagtgtgag ccgcatcagg cagtatgaag tcctttcttc tcaagccacg tagctagcct 65040  
 taaagggttaa tttcataacc cttaagggtta tttttttttt ttaatttttt tttttgagac 65100  
 ggtgtctcgc tctgtcgccc aggctagagt gcagtgggtg gatctcagct cactgcaagc 65160  
 tccgtctcct gggttcacag cattctcctg cctcagcttc ccaagtagct gggactacag 65220  
 gtgcccgcga ccatgcctag ctaatttttt gtatttttag tagagacggg gtttcaccgt 65280  
 gttagccagg atgggtctca tctcctgacc tctgtatccg ccgctctgg cctctcaaag 65340  
 tgctgggatt acaggcgtga gccaccacac ccggcccccac ttaagggtat tcttttagctt 65400  
 gaacatcatc tctgagaaac tttccctgac tgtgggtctcc tctccacct caagactgga 65460  
 tgagggtgtct tgctaagccc cctgtagcac ccacactct ccccatgggt cgtatcacat 65520  
 ttctcatcat caccgttata tgcttattat catcactgct gctgcctaac ttcaccttgg 65580  
 gccaaatggt gtgcaaaggg acttaaactc ctttctttta tctttacaac atgatcaggt 65640  
 agatgttggt ctgtttctct ttagagttga gaaaatagaa acagacaggt tacgtaactt 65700  
 gctgaaagtg acacagccga tttgccgcta atcagtgtga cttcggaagc tgcacttttt 65760  
 tttcaacttt tatttttagat tccaggattg cgtatgcagg tttcttacia aggtgtattg 65820  
 tgtgatgctg aggattggag tgtgattgaa cttgtcacc ccaggaccaag catggtaccc 65880  
 aataggtagt ttttcaacc ttgccttct cctccctct ccacccccca ggagtccctg 65940  
 gtgtctgttc tcatctttat gtccatgtgt acccagtgt cagctctcat ttctaagtga 66000  
 gaacatgtga tgcttggttt ctgtttctga attagtttac ttagggtaat gacctgcagc 66060  
 tgcattccatg ttgctgcaaa ggacatgatt ttgtcccttt ctatggctgc agagtattgc 66120  
 atgggtgtcca tatatcacat tttctttatc cggttcactg ttactgggca cctgggttgg 66180  
 ttccatgtct ttgcaattgt gaatagtgt gtgatgaacg tgtgagtaca tgtgtctttt 66240  
 tggtaggatg atttattttg ttttgagtat aactcagta atgggattgc agggtcgaat 66300  
 ggtaattcag ctcttagcag aacctgtatt tcttactcca cctccccgc ctgtccttag 66360  
 tatacagcag tggtctctta ttgccttttt ccttatagg ataagccct ctgctggactg 66420  
 ggctggggct gtttgcccat tataccctcg gcttctagga cagtggctgt gacacagcag 66480  
 atgtcaaaag aatatcttta agattcagag tgtgagacac tgcactagca ccgccatctc 66540  
 atgggccctc acaacagccc tgggaagggt gcctgcaccc tctctaagaa atgaagaaac 66600  
 tgaggtcaca tgttgaccat ggtcacaaag tcacctgagg ggagggtgaca ggaactgaac 66660  
 ccactgtcac tctgtgtttc cctgggaccc tctgagcgca ggaggccctg gttgctgtgc 66720  
 agtggcaggc caaggcaatg ccttgggtgga gctggggccc atttgccca ctgacctgag 66780  
 gaaagcagtt ttgtgaattg gcagttagct catttgctga catgggtgagt tacaggaaat 66840  
 gccatcatgt tctatcatg tgaaacaaag tgagaaatag gttcagggtg ggaggctgaa 66900  
 agggaggaat gcagacagcc ccgctcccca cacttgctcc aaggctgggn agggaggaacg 66960  
 ggaagggtgc tccctctctg gattcagtea ccttcttctc ttcattcccc tgcagtatcc 67020  
 cctcattctt ccacggacac gatcagcccc tgcttcttgt tgctcagatg tcatcacttt 67080  
 tctgcagagg gaaaagaaga gaccagatca gaacaagggc ctggcggtgg ctgtgcactc 67140  
 cgaaggcact gtgtgtgcct gagccccacc acggcctccc ctgcagggtc caggcagcct 67200  
 tccttgagct ggcattgagg ctgtgggagc ccggtccact ggcagggtct gctgcattca 67260  
 agtctctctc atccctgct ctccccccc tctccctctg nngccccctt ctctgacagt 67320

095234-06504

gctgaccccc ctctctcttc cccactcttt cccatcctcg cctggcctcc ggtttgatg 67380  
 ctgtccacac acttcccagag ggccctgagag gacctccgtg tgaggcaatg catttcccag 67440  
 gtcacctctg tgtctctctc caggcttttt ccaggggactc cccgggggtca gtctcctctc 67500  
 cccactggaa cggggaaact gggattggcc tagaccggc agtggagtcc caggtgccct 67560  
 gcctgcccgg ctgactccgc ccaggggaggc ctcccacaga agctcctcca gactccacct 67620  
 gttacctccc cactcctct cacccaaggc tgtgctgtgg ccaagtcagt tgtttagtct 67680  
 acactttctg tttagtctac accatggcta cctcaaggcc cagtgaagggt gtgtagtata 67740  
 aagcaaaatc aaatccatat ttcagttttc cttaaaaagt gaccttcata ttcaggccag 67800  
 aagaacagaa tgggtgggtg gatataatttt gagttttcat gggtttttgt tttcctgcct 67860  
 cttgttatac tttctgaaat tggcttttag tctaaacagg tttttttttt tttttttttt 67920  
 tttttnggca atgtgttttc ctccaaagag taagaataat aggcctcatg gctgggtcgt 67980  
 gttctacagt ttgtgaatat tttctcaacc tttgtcaaat ttcactctta cacatcctgt 68040  
 gtgaaattgg gcacgtgccg ttatttccaa cttagagggg atgaatgagc ccttaagagc 68100  
 ttgagttctc tgcccacatt gcgagttact cagtgccaga aggagttctg gaaccagggt 68160  
 ctctcagtc tccataccac atcccttcta gggcaccatg ttgcttctgt gtttcttgcc 68220  
 tctgcccact ccatgccagc acaactctcc ccaccctgc tttgggtggaa tcatgttctt 68280  
 ttggggtaga tcacaccagc cagaggcaac tgctctcagc ttagcagatg gtactcatca 68340  
 cattattctt gaagccttgg gtcaggagcc tgcccaccc atctgcatcc atttgtccag 68400  
 ccctcagaca attgccactg ttttcatgtc tattctttga ctctctatcc tgggtagaca 68460  
 acatggactg cccagcatcc tgtcttctgt ctggggctcc cactgtcgtc ctgaccacgc 68520  
 tgggggctgc cagtgcactc gggaaactcc cgagggaccc ccttcagggt tcacatcatc 68580  
 tgctccctcc ctagcatccc agcctagaac actttccagc catcagctgc attccccagt 68640  
 gaggcgtgca gcctctccca tgataggagg gcttcagccg aaagaacact tcaacaggcc 68700  
 cagaaaccca ggagcaccat tagatcagaa agcagaagca agaatgcac taatctcccc 68760  
 cacatcaatt gctatagttt tattaatctg catattatag gtcagtaagg ggatggcaca 68820  
 gtttataatc cctgcaagag tctgatgatc ttttggtgac cagaagtgcc attttttgat 68880  
 gggcttctag agatcctcca tcaggggatac cagacatgtt tggcatgcct gtgctgccgc 68940  
 gagacgctaa gcgtgtgtcc agactacacg tgtgggtcat ggggtccagca gcagagctgt 69000  
 catattgatt gtttgcctct actaaatgta taaagcctgc ctggtgtcca gaagaaaaga 69060  
 aactataatc caatttttta gaatccataa aaggtaagaa gtaggagaac atttagaatc 69120  
 cacaaaagat gagaagtagg agaacrgttg gatttttttag aatccataaa agatgagaag 69180  
 waggagaacc tccaaaagga aggaatcagc tgagagtatt gaagatgacc aagtacaaac 69240  
 aggcagaggg gagcgcttcc cttctctctc tcccaggcgg tgggctgcct cgctcggcca 69300  
 ggacacacag agcagcatcg tgcrttttga ggggcagggt gagctgctca tcaactagcag 69360  
 ggggtgctggc ggggaccaca gtgttctctt ccatctttga gttgaagtcc tgtgtgagaa 69420  
 atgagaaacc ttcattggca aagacagaaa gggacctaga atgtaacatt cagcagctct 69480  
 gttatctcac gcacctgtct gtccagttgg ggacgttgct gtatggagggt cagttgaaca 69540  
 atcacagttg aggagcctaa tgaattcttg caccaccagc cacacacatt attctgaaga 69600  
 gtgagccatt gtctctgatc ttatcaggat cacatcgtgg gatcatattt atttggtcat 69660  
 tctgaatata ccttttaagt ccaaagtga ataactaaat gtcgttgata aaaggaaaga 69720  
 ataaagtggg gtatgatttc ctttcacaga ggtctggaat ctctcgcct ttttcaagtc 69780  
 agtcggtggg gctggcaaat gtttaataac cagctcctct caccctcag aggaagccct 69840

096241-06501  
 10590-123260

tggtgttcag	tgtttgcaga	tttccattgt	gcaactagtc	ctccacacacc	ccatttttaaa	69900
ctaccactt	gatgtcactg	gtcatggagt	tggtgtcaca	gagccagtgg	gagtcaactg	69960
gagcagccac	tggactcatt	caagtgtttc	ccaaaacaat	ctgctcctag	aaggactctc	70020
ccttaatctc	ctaaccctgc	cattcaggat	gattccctgc	actctgggaa	gcacacgttc	70080
tagtggaag	actgatactg	ggcaactgat	aaccaagtga	cttaaacttc	tgagggttac	70140
aaaggggtgtt	tgtatcctca	gtgtctcatt	tcagattctg	ctcagagcta	aatgcaacaa	70200
tgtgagaaga	tgttagtata	ccagatcttc	atccaggaag	gaatcttaga	gatcattagg	70260
ttgtagggtt	tctcttctgc	agaggagata	gagggtcggt	gtcagattgc	tggtttgcca	70320
gtaccactcc	ctggagaaaa	gagcaaaaaga	aagaaacttg	ttagtcaact	gtgcagagcc	70380
accgtgagac	tgaatagctt	tgtgggtggc	cccgtgtttg	ctgcaagaga	cctctggcct	70440
cttgtagcag	ctgccacatg	gtaaacagag	ccgagatata	aggagtctcg	ctgaaaatgc	70500
agtcagatgg	gctctgaata	gaggaaggca	ggacactctt	gagatgggat	ggggtttctc	70560
acagcaccgt	acagggacca	cctgcaagat	ctcttgaggg	gcttgtgaaa	aacacatccc	70620
tgagggtcacc	attcttgacc	tgtctgttat	tgagtttctg	atgcctggga	tgtgcagggt	70680
taacaagccc	ccagatgata	ctaataggat	tcctgcctga	aaattgctgg	gtgaaggctc	70740
ttccccctcc	aagtataaaa	gaaggaaaag	attgatcctg	gaagaacatc	cgtagtagta	70800
gcaaaatttt	gtggagcact	tcatgaagag	gaattactag	gtcatttaga	aatatgtttg	70860
aattgtggat	catctttag	gcctttcttg	catatttctc	cacttagatc	cacaagacac	70920
atcgaatgtc	tttttataaa	ggggtttttt	aatgcccata	tttgacctc	tccacttaac	70980
agtcccatc	tcattttata	tgtgaaggta	atctgcttta	cagaaaaatg	taaaggacct	71040
gcacttctct	gctttgtggt	aagttgtaaa	atgcagttta	aagaggcagg	cctcatatcc	71100
tgatagattt	gtaggaagga	ttgcacagtt	ttaccagct	tcctcagagt	ttggcagaaa	71160
ttagctttcc	ctgagctggt	gtcttcccga	gctagcatgc	ttctcctatg	gggtgtgtgg	71220
ccttctctcc	tgtctttttg	aggcagagct	tcaatctaga	atctgttcac	aaactgaaca	71280
aatgcaacaa	acagtaaaca	gtcttttgc	catagttaag	gtgccttgag	ttgggtgtga	71340
ggggctgagt	gtgttctcag	gggtgctctg	cccacggctc	cggccaactg	ctgcaggtgc	71400
gcatcatatg	ggtgggtctt	gtggaatgcc	atcagcacta	gcttagtacc	tcctaaatgg	71460
gagctggagg	gctacagtgc	tcaacactgg	attatacgaa	tgtggattgt	ccaggaaatg	71520
cttttaatcc	ccctcatcca	ctctctaccc	acgtgacctg	cctctccctc	tttacttgggt	71580
gtttactcag	gaatgtgggt	gagttgtcgt	gttagcctag	aacagccatt	cccaaacttt	71640
gatggaagga	tgccattcac	tttgaaaatt	atcgagtagc	ccaaagagct	tctgtttcca	71700
tggataattc	ctatcaatat	ttactatatt	accaattaat	taaaactgag	attagtattt	71760
atttgattcg	tattttattt	tacatagcta	taataaactc	atacatataa	aaaattatta	71820
aaaaatgact	gttttccaaa	ataaaattag	ttagaaatgt	gacattgttt	ctacattgaa	71880
aaaatctctt	taatgtctga	tttaatagaa	tccgctgaat	tttatttgct	tcattcatcc	71940
tgttttgctt	gttatttgaa	gcatataaag	caaaagctga	ccttgccacag	atctatagta	72000
ggaaaagcag	ggggagggcc	tcatggaccc	ctaaaaggat	ctcagcgacc	tccaggggtc	72060
ctcaggctga	ccaaacattg	agaactattg	acctggaaga	atgtaaaata	ggaaaacagt	72120
gtctccccc	atagaatttc	gtgtaaaacg	tggactgtgt	tacaaagtca	gatgggtgca	72180
gttgtcctgc	ttaaccgcta	atcaggagct	gaaggccaga	gactcacagc	tgttccccag	72240
cctggtagtg	aaccagagg	cctgtcttgc	tgtgcagtgg	gacaggaagt	tgcatttggg	72300
agtctcatag	aacacactgg	aagatgtgtt	ttagcttggc	caggttcatg	caggacagat	72360

tttctgcata	aagaaaatca	atgacagttt	ctgaaactgc	atcctggaag	ccttgaccag	72420
tttgggtaat	aacaagagat	ttgaaagtgt	ggggtgtaca	gggtgttttg	tgaatctagg	72480
tgggtgggt	gattattatt	atgttgaatt	cagctttcag	ttctacctgc	ttgtgagttc	72540
caaactttgt	gaaaattagt	tgcttggacg	aaacttttct	ttgcctctgg	aaggctgtca	72600
gaaagcgaga	tttcccagct	tatgtgcagt	gttatagtta	atagagtaat	ggctctgcaa	72660
agttgttcct	ttactttaaa	tgtaatttat	tttgcatttg	tgctacagaa	cggtcataag	72720
tgtgcccttt	tgtcctcttg	tttggaaact	gggtttttat	aatgtgtgtg	gtctatccga	72780
agattattgc	ccattattga	acaccattca	tagcaaccat	ttgcattagg	cattgtacgt	72840
gtactctcca	ctctgcaaac	tatgtgttct	gtcccttttt	aaaaagagga	agctaagggt	72900
cagagaagct	aggtagtcca	ttctgagctt	cacgtgccag	aggccatttt	gtacttactt	72960
caaatgccat	tgaaataaat	gcacatcaga	gaattgttct	tagcataagg	ggcgctacat	73020
gtaacttttt	attagtgaat	tggtgatgtg	tcaagggtcg	tggttgatta	gaaaggcgct	73080
cagaccctgg	ctccagggac	tatggagcag	aactcgaggc	cagtgcctgt	cgagcgggtc	73140
cccacactcc	atctgtgtga	cctgactgtg	gatggcctgg	ctctgccgtt	agattgccac	73200
gggtgccctcc	tctggttgaa	cctttctcga	gaagtgcctg	ttggaggctt	gagtgcagag	73260
cctgtgagaa	gctctatgtg	gttcctattg	cctgtcagct	tgctgataaa	ggtcattgggt	73320
ttggcaaaat	ttggcccaag	gtttgccttc	tcataacata	ccactcggtta	gcaaggctgg	73380
gaggaagggtg	gctatagcta	tttctggaag	ctgcttaggg	ggctgcctcc	ccctaaattg	73440
gtacataatt	tgcaaggcct	attgcaagat	gaaaatgcag	aaccctttct	tgaaagatta	73500
ttaggaattt	caagacagag	acaacagagc	atgaagcctt	gtgcaaggct	cttctaagca	73560
cagagccagt	gtgaccgcac	agaacacaca	cccgtgaagc	cagctctgcc	cccaccatct	73620
gaccactctt	gagtggccaa	ttagcatagg	tcactcccca	ccctgctagg	cccaccctct	73680
taggaatggt	gtgaggctta	aataagaaat	agccactcta	caagcgggtg	caattagcat	73740
gggctctggt	ttctgtgtga	ggtagtgtgc	taacatgaga	gggtatctga	ttagctaaaa	73800
cgataacact	gacagattaa	attcagaata	actaaacctt	ccctgtgttc	ctttatgcca	73860
catgactcct	gcatattctg	ctaccagcac	ctgtttgata	ccagacggag	gggtccattt	73920
gggatgggac	aggagcatca	gcagaaatgc	agaagtgggg	aagtgtctca	tcttcttgga	73980
agctgagctg	gcaagggtta	tggaatgaaa	gagattgtga	atatttttga	gactatgagg	74040
aaaccagtac	actggtgttg	cccagtagag	aagccacatg	tggtgtttaa	gcacttgaga	74100
tgtggctact	ccaaattgag	gcgtgctgtc	agtataaaga	acacactgga	tttcaaagac	74160
ttggcatgaa	aaaagaatgc	ctaattgtct	agtattttta	tattgattat	gttgaagtga	74220
tagtattttg	tgtatgttgg	gtgtaacaaa	atatctaatt	aaaattaact	tcacctgtta	74280
ttttctaatt	tgggtgctag	aaactgttac	atcctgcatg	ggggtcacat	tccagttcag	74340
ttgcatgtgc	tgctacccat	tggtctacac	acacacacac	acacacacag	ctgcacacaa	74400
cctagagggg	tcagagaccc	caggagcccc	tgcttctggt	gcccaggcta	agcgctggag	74460
tggagataaa	agctgggagg	gtgggttaag	aggtgagtgc	acggagctcc	aggctaacag	74520
agtggataat	ttgttctttg	agcactgggg	agctatggat	tgcttactag	cagcaagggtg	74580
acttgtgcag	ggtatatctg	ggggagggtt	actgggggaa	gagatagagg	aggcaagaag	74640
tgaatacaga	acgagaaatc	aggacagtgg	ttaggagacc	gtagctttcc	tcttgagtca	74700
agttcagata	acacatctgg	actgatgaaa	ttctttttca	ggaagctgag	gaagagccca	74760
tgaaaatatg	ttcctccctg	tgctgagacc	gaataattgc	agtgaacaat	taacgtgtgg	74820
cctagatcca	ccttttgcct	tcgctgatcc	aagcagggtc	ataattcttg	cctggggcca	74880

095211062001

agcttggccc tggctgccag ctgcctggct ccagatgttt cttaatcggt tcaagtactt 74940  
 ctctgctccc tggaaacagg cactcccac agtcacattc cagaggagga ggaagaggaa 75000  
 cttgacaagt atcagctaca aaagcctcct gaacaaaaga aatcctttta gcctatttga 75060  
 ataacagttt tttgtgaaaa taatcaggat gttgagagct tttttttttt tcttttaaac 75120  
 tcttttttga aggttaacttt tgtgaaaaga aaacacctgc tgctcctcag gctgtttcaa 75180  
 aacactgcct atagtttgaa agtacggaga tatgcatgtg gtatgaagca tttgcaggca 75240  
 taatatgtgt agtctgggaa aagcagatcc agagagtgtg ttagtaagg cgaggccttt 75300  
 tagctgcatt tagatgatgc tgggattggg gtgggtgcag ggtgcagcag tggggaggaa 75360  
 gaactgtgtg tgttcctcct gagaataggg gttatgtcta gaggattaac agttttcctt 75420  
 tttcnttttt tttttttttt ttggagttgg agtttttctc ttgtctccca ggctggagtg 75480  
 cagtggcatg atctcagctc actgccatct ctgcctccca ggttcaagca attctcctgc 75540  
 ctcagcctcc cgagtagctg ggattacagg cacctgccac cagcctgac taattttttc 75600  
 tattcttagt agcgtggggg tttcgccatg ttgggcaggc tggctctgaa ctctgacct 75660  
 caggcgatcc tcccgcctag gcctctgaaa gtgctgggat tacaggcatg agccaccaca 75720  
 cctggccaac agttttcctt tttcgattga agttcagcta tttgcaggac cgaaggtagt 75780  
 tctgattact ttcacctgta cttccaccaa aaaataaata aaacaacat gagtaattgc 75840  
 tgatttttaa ttgaaagcat tattccagga ataactggtg gacttcgttt gcagaggaag 75900  
 tggcaaagac tgattgatat tatgatccag cttctaaaga ttttgcgtgt taatctgaag 75960  
 cacattggat ttctggttca ataggctttc tttttttgtt tttattatta caactaatat 76020  
 gtattctttt cacagggcga acctttccta cacaccata cttctctgcc cagcttgagg 76080  
 caggacagct atcgctttac aacattttga aggcctactc acttctagac caggaagtgg 76140  
 gatattgcca aggtctcagc tttgtagcag gcattttgct tcttcatatg agtgaggaag 76200  
 aggcgtttta aatgctcaag tttctgatgt ttgacatggg gctgcggaaa cagtatcggc 76260  
 cagacatgat tattttacag gtatagagtg ttccttatgt ctttaataca acaaaatgct 76320  
 aagaatgttt cttatccctc tccagatgtg cctcaggagc tttttcaccg tcaggtaaca 76380  
 ttgtaatagc tgctactgct gataaaggac tctgtgctag gcattattcc aagcgcttca 76440  
 tctgcacttc cctctaataa caggaagaca ctgttcatcg tctcaatttg tagattggaa 76500  
 aactgagtct ccaagagatt ataaattggg ccagtcaca cagctagcaa gtgtcagagc 76560  
 tggactggaa acccaggcct ctctgactct agggccttcc ctcttgcccc catcagccat 76620  
 cagatgatct cagacctacc tcccagctc tgcactgtct cttcctctgc ctcaccccca 76680  
 cccttgctcat ctcagggttca gctcaaatat cacatcctgg gagaagctca ttctgactac 76740  
 cctgatgttg tgttccccac ttccaccttg ggcacactgc gtcacgttat cctggctgat 76800  
 ttcttttgca caacacagcc actgccagaa atgatcttgt ttccataatc atctccctgt 76860  
 ctattttctg atttttcata gcctgtgaac tttaggagag ggaagggatc ttacgggtct 76920  
 tggagccgag ttctagttt ctgaaacagt gcgtgggttg aagtaggcac ccataagta 76980  
 tttgttgaat gaacaattct gtcagagaaa accaaacaca gtagcgatt gcaaatacca 77040  
 cgtgctgctc ttgctgcctg tcagagggaa aactctggat cctgcttcag gaatattcct 77100  
 aaatgttgca gcacatgttg atatgttcat ttactaccag taagatacta tgccttcaga 77160  
 gctctagaga gtatcctggg agggaaataca ttagagccaa ggacttgctt tgagagcacc 77220  
 aaattatgtg attcaaaatc ttttcacctt gacctgtgaa catggaccac gtgaatgcaa 77280  
 atatcataga aggaactcat tcaactgaaag attttgacca cataaacctt tccacatgta 77340  
 ctgtgaggtt ctctctacat tccctttatt aactttaaag acagtgggtca ccaggcagtg 77400

0976231-063504

gaatttttga gttttctata atttatgtaa cacacaactc ttttggggtg gtgccttttg 77460  
 ttgattagac agtcttcgat atgggagagc cacagctggt gcttatggga ttatattatc 77520  
 tgagcctctg aaaacggttt tgttttcttt ctctcagttt agataggaca tatccaactt 77580  
 ggtggatctt agcggattct gaccctctgt aggttggtgt ttcttttaggc tcaggccgtg 77640  
 gcactgctca gatctgggct ggctctcggg cctctgtgag cctgtaactc ttgggtggcac 77700  
 tactaggaac tggcatgaga tttctgccag aatcatgtca ttctgtgaag ttggagtcc 77760  
 acttttagttg gaaaaagttt ttatttcac ctaagatgca cacttgctt cttgttttaa 77820  
 cttgccaggt atctggatat tccatatatt atacacccaaa agaaattatg cttctcctgc 77880  
 ctattgagta atttcagggg tccagagggg acttgctgag tgaacatgta caatggattc 77940  
 ctatggaatc ataagatgcc cctaattcag tcttagtaaa gagactggct tcttatttct 78000  
 aattcctcca ggcttgagtt gtgcaaagag tatgtatttg taagagaatt tatgaaatgt 78060  
 ttgcacaaga cagattttta gatcttctta gtggaggaat acaagggaac aataaaaaagg 78120  
 aagtggcagt agaagaccca gcgttagcgt cctgggccta caccagcca gtgcctggca 78180  
 ccagcaggca cttgggaagc acttggttga tgaattagta gctgagctca gtggatcgca 78240  
 agccaaatcg aatgtttaaa gttctagtaa gtcttctctt acaccacccc tgtgagcagt 78300  
 aggcataact ttattgctgt ggcagatccc taattctcag cccttggtggc tgtcttctg 78360  
 cagatccaga tgtaccagct ctgaggttg cttcatgatt accacagaga cctctacaat 78420  
 cacctggagg agcacgagat cggccccagc ctctacgctg cccctggtt cctcaccatg 78480  
 tttgcctcac agttcccgtt gggattcgta gccagagtct ttggtgagca ttagtaaatc 78540  
 tgtttgccag aaccagcctt ctcttattag aggggaaaca tttcctgtct ctccrtggtg 78600  
 attcttattt ttatacctgt agctcttacc agaacagggg attgtttgat agtctaagat 78660  
 tagtcagggg tgggttttgt gactttggag tctccttaa cttctgataa tcacggggct 78720  
 tcccttagat gccttcact tgtgggatgt ggatccgac cgtgtagatc cgatcgctca 78780  
 ccatgagggg ctccctagag cagacatttg gaggacttg ctgaggagcc acaggtgtat 78840  
 gtttctcatg aattgccttc ctccagcact ctgggttggt agtattgact gatgctgact 78900  
 gtgggcctct gggcccttct tagattccct tggcatctct tctccctt tctcttcttg 78960  
 ccctgccctt ggctctacac tttctcccaa gtcactgtct tggagaccag tgtcaggacc 79020  
 ttgagtaaca cctccgtgtg gatggctcgc tctccccgct cagccttgac acttcatgaa 79080  
 ggcctcttgc cctgagccc acatgtcaca gccactgcca ctcccgtgcc cccgctgtta 79140  
 accttgggtg gttcacatgt aaaacctgcc tttatattct tgatttactt tttgagaaca 79200  
 ttgtcaaagt taggtgagtg ttcatacaca aagccttcaa cctgccttca tatgcaggga 79260  
 tagggctgtc cacgtgcgca tcaggaaccg agtggaatgt tgtgagcatg gtcagttcgg 79320  
 gcacagtttg ttttccctac tgcagaataa aagtgatatt tttgacaatt caggttcttt 79380  
 tttttattgt aaaggaggag gctactaaaa aaatgatagt tattatatat caaatgtttt 79440  
 taagcatcac ttgacagctt aaaaacatgt gatctttaaa aaatttggtt ttatgattag 79500  
 agagcatctt aagggaatg ttcaaagaca ttgatactac ttcagacatg ctttgggtaa 79560  
 acatctttaa tatccaaatt ctagaaatcc taaaatttgc tttttaatat aagtgagcat 79620  
 ttacccttct tctctctttt cctttccccc caaatactag atttttatta ttcactttta 79680  
 tctacaagaa cctttaaaga gtttccatt ttgctttact ataagaattc atattccttc 79740  
 ttttctgtcc ctgaaaaaat aaatcacta aattaaaata gatacaaaaa gctatctcct 79800  
 ggttgagcat atcttttagtg agagttcatg aaggtttata ccatgggttaa aaaaaaaaaa 79860  
 agattaacta aaagcctcaa aattgtgtgc ttagtttatt aacaaaagag ttacagaaac 79920

taaaatctca	agctctaggc	tttaagcttt	cttgccaata	acttctatgt	ttttgacttc	79980
tctaactctg	gaaattaaaa	gaaaattatt	aatctacctt	ccttacattt	tctccacatt	80040
ttagctatga	ttttcataca	gggtcatgaa	gaggagtga	gatggaaatg	gggaggagg	80100
agcgctgttt	gttaatgggt	tgtaaacagc	tcaggcatta	aattacttgg	ttagtgaaga	80160
aaattctacc	aaggcaacca	ggctgaccac	agactggagg	gctgaggggt	catcactgag	80220
tcatctctgc	cctggggccc	caggcactgg	agctgctgct	tgcagaaagt	tctggggctc	80280
tggaagagaa	atttttccct	cggtcataa	atgggtaaaa	agacgttaac	aaacaagcag	80340
actccacttt	ggaaatgata	gcccttctat	tgcagagtaa	tttgaagctc	tctgaagctc	80400
actctaata	cttcataaat	caaagctgca	gcttgtaaag	gtaagatatt	tttctgtaga	80460
ctttgtaggc	agtgggtgag	ctcggagttt	cataaacatt	atgcatagag	atgccagtg	80520
ctacatcatt	cctggatccc	acagactcct	gctgtgctaa	gtgggtcggt	gtccagctgg	80580
ccaagggctc	ctgggattag	aggcgggaag	tgggatctca	aggccgcact	ggcttgtgat	80640
gtcactttgc	aagaactgct	tttttttcca	cagtccatcc	catctttcag	tacttaaaaa	80700
cagaaaagat	aggtttttac	aaaccatttc	tatttttagc	actgatgact	tagagaatgg	80760
tgatggagat	agcttagttt	tatatttcaa	agcctgcat	tcagtcacta	tagtcttttt	80820
tggcctaggg	cccatttcat	tataagcctt	taagtctgga	taaactctaa	aaacatgtag	80880
aactttgttg	actaagtaaa	atatttcagt	ttgcaccacc	ttagctcata	tattagttaa	80940
taggttacct	ggcattaagt	atgtatctgc	tccttgagg	ggcggtgcc	agtgatgtgt	81000
gcgccttaac	cttcacatga	tactcacacc	ttgctgaatg	gcagttcttc	tacctggtgt	81060
cacataatgt	caatagcccc	ttcctgtatt	tttctagctt	gagtacagca	gggccctggg	81120
agagacgctt	gctctgttca	ctttctcatc	acatctacct	ttgggggaaa	aaaaatctaa	81180
aaaacaggac	ctggcttgct	cctgatggag	gaggaggctg	cagtgttcag	cctctgatgt	81240
ttttctatag	gacatgctgc	caaatagatg	agggaggagg	aggagtataa	aaactaagg	81300
tttggaacaaa	aacacagaag	ccacctgcaa	tatagtgaag	gcttcagaga	gacttttagga	81360
tgaaaataga	ctgaaaacaa	gattgtttct	gtggccagga	aaatctccag	ctattcaggt	81420
gacatgatgc	cgcgtgatga	tgagtgtgtc	cagtctgtct	gtgctgttgt	tctgcacagc	81480
actgtcatca	gccttcagcg	tccccctttac	ccgttactca	tagaatgtag	cggagccacg	81540
actggaggac	cgcagccttc	cagaagaaag	ttgagaaggc	tcagccttga	caaagacaaa	81600
gggtggctgga	aaaagagatg	cagtgcacac	tcacatagga	agattgcact	ttgagatcat	81660
gaagtttagt	ttcaaataga	gttccaatac	acagtaacgc	aataagagg	ttgctgaaat	81720
gtcctcaaag	aaaagcagtt	cctttgtgtt	gttcccagcg	aatacagtgc	aaagtaatat	81780
agtcagctga	atttaagatt	cctatttcct	gccggataaa	acgtcttgcc	tgtttctagg	81840
tggcttgaaa	aaggagagga	gagaaggaag	aggcaggaga	aaagtccac	tgaaggagc	81900
tgggctacag	tgtagtgagc	taggctactg	cctcactgcg	ctgggcggct	ccaacagttc	81960
actgtctcct	aggagaggtt	tcaaatgcag	gacatttgct	cacttttcca	aggagagtta	82020
ttgttttttt	gttttggttt	gttttggttt	gttttaaaaa	attccagaat	gtaaatgtat	82080
aagataccgg	aagacaggca	aaataaaaaat	aattgggtttg	gggcagtggg	tttataggta	82140
acatttttct	ttactatrrt	tttaaaatta	gatgtgattt	aaaaaaattt	ccaaagccaa	82200
aaaacggtac	aaatgcttta	aaggatgaag	atgttggtccc	caagtgtcat	cagacaaatt	82260
taggaggccc	ttcctcccaa	gcaaagcttc	ctgcagtcct	tccttcaact	ctgaattcaa	82320
gcacattcct	gcattgtgca	cccaaataat	ctcccgattt	aagacccctt	gtgtctcaca	82380
gaagcttctg	gggctgaact	ttctccggcc	ttggagggtt	ggacgctttg	aatgggagga	82440

gtggtggtga gtggagcatc tctggcagca ggcatttggg agtctctggc aggaatcaat 82500  
 cagcgtagtc tccaaagggtg gcctttctct gacactaact agcccttgca ggggtcatac 82560  
 ccataacctg catctcatta acatcatctc cttaccagtg cactgacctt gtgagaaaaag 82620  
 gaacaacaag cattcagcga ctctgcgggt gctccagggg aagttagaat tgcttggctg 82680  
 gggcagaggc ccctggtgat ctggacctgc gtgcccccat ttgcccacct tctgcccctg 82740  
 acaaccagtg ccctgcctt gccagccaga ctgtttttca ggctcctgca cacctccctg 82800  
 tattgacacc ctatttttctt tttattcaga gtattaatcc tgaggctctga cctaggaaat 82860  
 tttcattggt tcttcaagca gtcaccttct tgtgggcctt ttctttctctc tttgttctcg 82920  
 taacaccctg ggcataactc taccgaacca gaactccttg gtgtctctgc agcgtgttct 82980  
 ttgtgttttg ctcatggctt aatctccaga gcctaataca gtgcctgatg tgtattagat 83040  
 gctcaataga tgctcattaa gttaaagtag aagacacctc tcagcagagt tctcttaagg 83100  
 tgttgatgaat agcattggga aagaacattt attttttaat tacattaaat acaaacagat 83160  
 ataataaaat aaatcatatg ccagtgcta tgtcttaatt ttttaacata tcaataaaga 83220  
 gactttaaaa cacataacac caccctctcc cctccaaatt tcttttccgg gaaagtctcc 83280  
 ttttggaatc ataggaagca cttactaagt tgatttattg taaaaaaacc aagatcctaa 83340  
 taaatctcag aagatctcct gttaacctaa agagaccact gatgtggatt ctgtatttgg 83400  
 ttgtgctgac aaaagtttcc cagtaattgt ttattttaat tggcgtagat gtggtactgt 83460  
 acctaattta aggcacttgt ccctctgaga gtagagacca agctatagaa aatcactggt 83520  
 gttgtagggg aagcctttcc ccaggatccc tgcaaaaaag gtcttgattt ttattctgaa 83580  
 agatgccctc attttttgtt cagctataaa agttcatata ttgaaaggag gtctaggaag 83640  
 tctcactgtg taaaccactg aaacttcaaa tttactttag agttttgttt ctggaaatgt 83700  
 catttctgtt taaaaataca tctttgttat agtattattt tagatctttt tattttctgt 83760  
 agtggggaat tatacaggta gactacattt tataaaccag atatttcaga ggaatattct 83820  
 tcaattggcc tgccttggtg tatgtaacac ttaccctgaa aagctctgat ttcaaagaca 83880  
 cagttagttc tctagtatat cttcccagcc tcaacaacca gacttaagaa ggaagtgaag 83940  
 gattcatctt tcccactttc ctgcggccac cctgagccat cagtagttgt gatgtttgtg 84000  
 gaaagagtgt ggacctgag ctgggtggga gaagcaggct gatctcagcg ctggcatggc 84060  
 ttagggtctgc acccatctca gctcacatgg ttaattaagg gttttgtggt gggttacagag 84120  
 gatctcgagg gctatcccag ccagcgggct cctgggtctg tcatccctgc ctgtgctttg 84180  
 ttcagaaact acagggattc agtttcccat ttgcacagca gcaccagtc tttgcttttc 84240  
 tgtttctctg tggcttttaa atgttatcat attaacatc tagagaggca ccctgcaagg 84300  
 ttattcctct cacctgcttt tgctttctct gatttgatga aatttacagc ttctttctct 84360  
 cttccattat ctttcagcca aaagaaacag agaaaagaaa tactgacact tgccctccat 84420  
 tatatttcta ctctgatttt taaaattgtt tttttcttat attattattc tagttattag 84480  
 gtaacctgcc tcagtttagt caaccaataa ttagttatcg tsgctctgct ttaaccccag 84540  
 gacatcagac tcttttttcc cccagcagct tcaactctat gaggaagggt agacagggct 84600  
 ggggttgctg ctggccgct tgccttggcc ggtgccctcc ctcttattct gcagtctgta 84660  
 tagaagttgc atccatttgc cagccactct aagaacaaaa tatggccaga actaggaagt 84720  
 aaccttgaca gagttcttga actcctcaga gggaaaaatg ttctttattc cattatcatg 84780  
 ttaaaaatca gtaaaactgt atttaacaaw gtacttctgc agttgtacag ctgttgatga 84840  
 gtttttaaaag atctttgaat tctattcctt gtttcaaac agaggaaaca gagacacttt 84900  
 ttcacttact ctatcttaat ttctgatgct ttatctataa aaatctttta gtgtgaccca 84960



taaaaacatg	tttttagtatc	tccttttaaaa	cccaggagca	ttcctggaaa	aatagactaa	85020
taaaaccttt	ttccctttcc	cagtttaact	tttgaagcat	gtttgaattt	tatttttcaga	85080
gtaaaacata	atttttaatg	tttatgtact	tttatttgca	atactgtctt	gacaacactg	85140
tctgagatat	caggctctta	aaaatgaaat	aaagtttgca	atgtggggct	atgtctccca	85200
cactcctgct	ctgtgatgtg	tggaaaaggc	aatggaatgg	tattgcgtga	gaaactggtc	85260
tggtttaact	ttctgcattt	ctgtgttttc	tcagatatga	tttttcttca	gggaacagag	85320
gtcatattta	aagtggcttt	aagtctgttg	ggaagccata	agcccttgat	tctgcagcat	85380
gaaaacctag	aaaccatagt	tgactttata	aaaagcacgc	tacccaacct	tggcttggtg	85440
cagatggaaa	agaccatcaa	tcaggatatga	gtcagtccaa	accttgcaaa	tgcttaagcc	85500
atcctagata	tgtagaaact	taaatctctc	ttgagcagga	actgtttcct	accactttgt	85560
gttctgaacg	gcattctgca	tgatgcctgg	catggaggag	gcataacaca	aacgtgtgga	85620
atgatcgtga	gtgtgtgtgt	tgtgagcgtc	atggtgaaat	gccacatgga	aacatggtgg	85680
caatgttttag	ctgtagaaac	cagcacaggt	tattagtagt	ttcttacatt	taagagactt	85740
cagctctagt	agcttgttct	tctgaaacat	atataatatt	atgcaatgat	gcaatgtagg	85800
gttttgtaca	ttgagtgcct	tgatttgtgt	gtgtgtatgt	tgaatggttt	taattggaat	85860
tttctccaaa	taattctttg	ataacaaaagt	tatgataggg	aacatatatt	ctatgaattt	85920
gtttcatgat	gtgtgtgtgt	ggtttttttt	taactgaatt	cagttcaata	tctgtggctt	85980
cattacctct	ggtttcagta	tacaatagaa	ccattatcct	ctgaagtgtt	agaggctgag	86040
aggtgagttt	tactgggaat	tacaactaaa	ctagatgggtg	aatgccctgg	gttgggcatg	86100
ggagcagatt	ttgatcctgt	acagtttaag	gaagaacctg	ccagtaatgg	ctgctgatga	86160
tggaggacta	tgtccttag	tagagaaaagg	ggcatcattg	gaagtgtcta	gaggctgggt	86220
agctacgaaa	gcactggaag	ggatttcatc	attgagtcac	tgcagagtca	gcagtcaagc	86280
ctctcataag	cctagaatct	gctccagata	gtcatgtgtc	acttaataat	agcaatacgt	86340
tttgagaact	gcatacttag	gtgatttcat	cattgtgcaa	acttcataaa	gtgtacttaa	86400
acccaagtgg	tatagcctac	tacacaccta	cacatgttgt	tcctaggccc	tacaaatctg	86460
tatgacatgt	gactatactg	aatactgttg	gtagtgtgaa	cacaagggtg	agtattttgtg	86520
tatctaaaca	taactaaaac	tagaaaaggt	acaataaaaa	tatgggtattc	taatcttatg	86580
ggaccaccat	catatatgca	gttcgccatt	gaccaaatag	ttgtaatgca	acacacaact	86640
gtgtaacgaa	agcatagagc	aatcaggcaa	aaacaaatgg	tgaataaaag	ctattttttga	86700
aaaatcctta	ctctggagat	ttctgtagtc	caaaaggaat	ccatgattcc	agtggattgc	86760
atccacatgc	agtgtttgtg	attttcattt	gcagccacac	cttaggtgtt	aagcacagaa	86820
aaagatgcaa	atltggcctg	caaaagaaag	aggtttcata	ccagttgtta	acttttagatt	86880
tctgtttgca	cattgcatat	gcccttatga	aagaacagtt	cttgtctgtt	ctgcactcat	86940
ctttaattga	gagcctctcc	atctcttttc	cttccttgga	acactcttct	tgatgtggat	87000
agctttggct	tgttggggcc	ttgctttttg	ccaggttggg	tgggcatcga	aatatacgca	87060
tgatccatt	cagtgcaggc	cgcaactcctg	agaggatata	aagtgggttca	ctgactgacc	87120
cacactcact	ctgctgcaaa	gtggaaaggt	aggggttcaa	actcaagtcc	ctcccacctc	87180
aaagtgcctt	agcagctctc	ctacactgcc	aagagcctct	ggaggtcatt	taatttagag	87240
tttttccta	ttttaccagg	attctaatac	tgacttctcc	acccttttga	ttcttttgatt	87300
tctggcattt	tcattcattc	tttctttcat	tcattctctt	cttacagctt	ttgttgcattg	87360
tacttactta	catttacagc	ttctagggca	gacccccgag	agccttgggt	acctagactg	87420
agggctatat	ccactacctg	acatgtcacc	ttgctcctgt	ccctcaggcc	atcccagctg	87480

acattgttta cctcctaagt attgagcctc agaaaaaaat cccattgtct cctattttct 87540  
gtaaaaaaca aaaaataaaa cgtattgaga atacttagga tacatcaggt gctgtttcag 87600  
tgctggaaga gtggaaatgg acacagcatg ggaagaaaat agctgtgcgt gtacctgggt 87660  
tgtttcaagc cgctatctgg ctatttgga gttgccattc atttttccac tgactttttt 87720  
tttttttttt tttttgagac agagtcttgc tctgctaccc aggctggaat gcggtggtgt 87780  
gatcttggct cactgcaacc tccaccttct gggctcaagc attctcgtgc ctcagcttcc 87840  
tgagtacctg ggactatagg catgtgccac cacgctcage taatttttgt atttttagta 87900  
gaggtgggggt ttcgccatgt tggccaggct ggtcttgaa ccttggcctc aagtgatcca 87960  
cctgcctcat cctcccaaag tgctgggatt acatgcatga gccactgtgc ccgacctcca 88020  
ctgacttaat aactctagga cataggtatt ataattccta tttttataga tgaagctgag 88080  
cagagagtaa catgcccggc cccctgtaga aaggcagggt ctgtgggagc cagggtgtgt 88140  
aggttgaggc tgaggtgttt gagtccagct ggacttaaaa gatgacctaa gatcggtctg 88200  
cagacatttt ccaaagaagg ccaggcagta tatatttttag gcttcacggg tcataacgtc 88260  
acaactactc acctctgcga cagctactca cctcagcctt tgcagcacia acaatccatg 88320  
acagcatgta aggggatgat gggctcatgtt gcaaaaaact ttatttttaa aactgtgcag 88380  
cgtgatgggc ttgggtccaca ggtggtggtt tgctgatccc tgaactaaag gatcacagca 88440  
tgtgagaagg tacaggaatg agagcagaga gcaattctca gaacctgagg tgttcaacat 88500  
tttgggggta tcgggagcac aaaaatttgc atttagggcc cgggttttta tcagtgggtcc 88560  
ttatagaaag tagatctacc cgcctctctc cctcttttcc ctctggtgtt tctatctgaa 88620  
cttgacatct gagtgttctc tgtcaggctt tctgctttcc cactgcccc tcccatccag 88680  
aggggtgctgt agtcttaagt tctacacaga aagcagacat ggggtccagc atgattcctc 88740  
tgcagcctta gagatcccc aggcccaa cttgggggtc ttcagagtag atgtaatgga 88800  
tgcttccatt gcagatggtg gagcatactg cctgtgtac agatggggtg gggcaggaag 88860  
tgagggcatc tgggtggtcc tctgtgatag ctgtggttcc atttcattaa atgccctccc 88920  
tgtatagata gtctcagcct gagcaactgg gacacagctg gatcctgggt ggaagagggtg 88980  
gccaaggaca gggctctgga gctctggctc aatgtggaca ggctgaaagc agccagagag 89040  
ggcaattcca aaggtgacag agcccaggac agaagacca ggggtgtctga agcctttgtg 89100  
gcagtgtact taccaaagga cctgtccatt gaataaacca tctacatctt ctgacccaag 89160  
aagaatggaa actttgggaa taattagtaa caaaggaaag gaggtcagtg ttgtattctt 89220  
gtcacagtgg gtgctctgtg gctgtgaagt ctcagctcag ttaaggaag aaaaaagga 89280  
aggggtggctg tgggaggcag ggcaaatagc ctattttcag catccttttag gctccactca 89340  
gagcatggcc tcagcccagc attgtcatca tatcatgtca gagcttggtta gaaactcagc 89400  
atctggggcc acacccgaga cctgctgaat cagaatctgc atttcagtga gatcaccagg 89460  
ggattcacat acacacaaac agctgagaaa cctgctgtg ggcaactctg ttagaaacac 89520  
aatgaacaaa ggagcccctg ttccagttga gcttgtaggt tagaaaccag ggttctgtga 89580  
ttcagaagac acacctcaa tcaggggcaa aggtgcctct tctgcctgtg ggggagccgt 89640  
cacttcttgg gcagtttgca ccgtggaaaa ggagtagttt tgtacgagga caactggtgc 89700  
cataccagga ggggtggggcg tggcggggag aagtgggtta ccactggcgt tgttgaaaat 89760  
tgctcacatg cagtggtaat aacaagcaga gggactttta gtgggtttga tgtttttgt 89820  
aattcactac agatagtgtg tgccccctt ttgctgatac caggccgact gttccactc 89880  
tccagccctt ggtatgacaa tgggaccagc agattggagg gcagggggtt aggaaggcgg 89940  
aagctctgtg gcgagttctg caaacrtca gggttcatga ctttattaat cagtgtccat 90000

ggactgtgaa gagaaatgct gagtctacaa tagcaaata gccaagaaca taaacagaca 90060  
 attcaccgaa gaggagatat ctagtaaaaca aatatctggg aaagtatttg gcttcatgtg 90120  
 taatttaaac ttatgtaaca tataatgctt tactctacta gataatagaa agacatttct 90180  
 tgatgccagt acccagcacc aagggtatac tgtatgcaga acattagcat gttgctgatg 90240  
 gcagtgcaca ttgattagt gctgttgagg gacaatttg cgaacacatat cccaagccag 90300  
 taaaatattc atacccttg actcagtcac cccgtttctt ggaatgtatc ctcaggaaat 90360  
 aatccaaaat atgagggag ccatatgtat aaggatattc tcctagactt gtcacttata 90420  
 ataacagaaa cttggaacta gatgtctaac acttgatgac tggattaata tgatgatggg 90480  
 aggttgagct ggtagaatat catgaagcca gttatatata gcgacatgaa aaagctctta 90540  
 tttgatacaa tgttaagtaa aggaaaagt ggtataggaaa ttttatgttg gttatgttta 90600  
 gaactagaaa aacatgcttt taggaaatag gaaatatagc tagatataaa agttgtattt 90660  
 ggtgatTTTT cttttatttt tcaagcttcc aataatgtag ctctattgct tcagtaactt 90720  
 aaaatagttt tatcttttcg gcaaaacatc gaaagtatgg aaatagtcac tcctactttg 90780  
 gcaaacagaa gagaaatttt cttcagtacc aaaattctgg aacttgactg aaaactatga 90840  
 agaacctaag agccaggatg acaggaaggc tctagatccc cagtaattac aactctagtg 90900  
 gaattgctct gagatgggccc agcaagaaag aagatgagag ccagtccccc ttgcagaggg 90960  
 gccaggatcc ttgcagcttt gtgtagtgc cagtgtcag ggaacggctt aggcaagacc 91020  
 ctgggggagg tgggcactgc actgttccag cctcaggagt gactcagacc agaaatgaaa 91080  
 acaccttaaa gtgtatatat cttgttttcc tatcaacacc tagtttttaa tttcgtctg 91140  
 ttttattcat ctgagacaac ataccaaagg attgggtttt taatgttagg ccttcctgct 91200  
 ctttctctgg ctgagaactg ctctggcag tggatcactt gtgctgtcta agtggtgcaag 91260  
 gacaggcgcc cctcccaatt cttttcttcc cccaagtaat tagcccaagg gctgaagccc 91320  
 tcgtccagtg accagggttc tcctttgacc accagcctca tattgccatg gtttggggta 91380  
 aattcagggg cataactgca gaatgaaggg cctaggagtc ttggcagtca ggagatcatc 91440  
 aggcaattaa gcagagatga ttgtgacca ggggtggttcc tagggattaa tggaggcctg 91500  
 gaagagttaa tggtcttggg tactgctgag agccattaac ttaacacaga acatcaatcc 91560  
 gtaggaaaag ccagagggtt tgttcccagg ctttccaggg taggagatca cttaaattct 91620  
 tgtgaaagaa aaaaagtaat agtgtacatg acattttattc agcaccatat ttataattat 91680  
 acatgagtgc caaacaatct cagttttaac atttgtggtt tttactgttc agactattca 91740  
 gaagttccat gacgtggcat ccattgatga tggttttgct gaggttgaaa tgtgagggtt 91800  
 ttgtggcagg tgtgtggaag cagatccctt agctaagaga gcgcctgtc aacctgccag 91860  
 gcgtgtgctg ggaacttcca ggagttgctt cgtttaattc tcacagccat cctgggagg 91920  
 aggcgtgtc tgccccagg gggaggccag gaactgtggc tgagagaggt taagtaccga 91980  
 gctcaagtca ttcataagat cgccagtggg gccgaattt taaggctgac tcaaagcctc 92040  
 tgagccagtt acctcggtc tcaaagggag ataataatcc ctgctgacct cagggtcgtc 92100  
 ctgaaggaca aaagagaatc agttcaatcc agtaaaacaat tctctctctc cctcttactc 92160  
 ccctgcaca catgcacaca caccacacag atataatgga ttttagtttt taggcatcaa 92220  
 atgacttttg tgccttatta atattatcca ctgaatcaaa aacagcaagc tgaaaaattc 92280  
 atctcaagga agagaaaata agattgttgg gaatggtgag aaaggaaaca tggtttttga 92340  
 aaattgattc cagggagat aggttagttt gaatgccagt agggagccat cagaagaagt 92400  
 agttttacac tgatttttaa caatattgga gttgcttaag gcaatgcaat agagaggcag 92460  
 tttctgccct tttaaagcct gacttcactt tctgaatgtg tgttctgac tagcagggtt 92520

0924 0650  
 10230 T22250

tttttttttt	ttcttttaag	atgggtcccag	cttgactgca	ttctcagatc	catcagataa	92580
acgttagggc	ttcactgctg	tgctgagagg	ccccagcccc	tgggggttctc	tcatagaaac	92640
aactggaaag	aaaggaaatg	ccttgggcag	cagcagcagc	agctgtcttc	tgattctgct	92700
ttccgccctg	ccttccttac	caagagaaaag	tacagacacg	gacggcttga	gtcacttagg	92760
cacttaggag	ttgtttttca	cacgtgtggt	gttttcgtca	ccattactat	tgtgggaaag	92820
aagacaactc	aggcatcggt	tcgtattcac	tcatctgtgt	gggtgacatg	tgggttttgg	92880
ctcatttctg	catatttgtg	tgcaaaggag	agttttttag	taaacagtcc	cattacttag	92940
ctgttcttgt	aactctgaaa	acccaactga	actataatta	aactttgact	tgggtgactct	93000
gcaaacaggc	tatgattctt	ttgtttcttt	tctcctttta	acccatagtt	gatgtatcta	93060
acctaacaga	attttcagag	aaaagaagtg	aaataagaac	taaaaataaa	tttttatgtc	93120
tttaaaaatg	agaggttttt	tttttttttt	tggcttttgg	aagggtgagta	tcaaaaacct	93180
gtacttaatg	ttaccttgga	attatttcta	gatgtttctt	atatcctttt	gtcccaagta	93240
aaattattac	cttctcagtg	cgtagttttt	cttatttatt	acttctagta	ccaagtgtag	93300
agctaagcgt	agaggagacg	cttcacaggt	gcgcattgtc	gtgattgcag	acgcctgcct	93360
gtacttgtgg	ggtttttctc	agtttttagta	cgtgatgact	tttctttcta	taacagggtat	93420
ttgaaatgga	catcgctaaa	cagttacaag	cttatgaagt	tgagtaccac	gtccttcaag	93480
aagaacttat	cgattcctct	cctctcagtg	acaaccaaag	aatggataaa	ttagagaaaa	93540
ccaacagcag	cttacgcaa	cagaaccttg	acctccttga	acagttgcag	gtagagcata	93600
tttataaagc	agcttcctga	atcacaaata	tatggtagtt	cattaactca	ccaaaggcaa	93660
cagcaggctg	ggctttccca	tgaccagagg	acctttccca	ccctgatctg	tttatagttg	93720
ggatcaaagg	tatcccggga	gaatgggtcc	tttttattat	ggagcagaca	gattgtcctt	93780
tgctaaggtc	aggcagtcct	agagctttct	gagaggctgt	ttctgcactt	aactctttta	93840
ggggacaggc	ccagagatga	acttggattc	aggatgccgt	ggcctgttag	ctgaatgccca	93900
gccgttgtca	ttactcaaag	agaatctaag	agcttttaac	ttctatgagc	aaaaccagct	93960
aggtccacag	agggatggta	aaggaggaaa	gtaacacaga	aataaatata	acaaaccaga	94020
agagatgata	attctttgtg	agtccttggt	gcatatacaa	agatttgatt	aatgaaggtc	94080
tcagttctcc	cctctagaaa	cttccatttc	aacacggata	tactcagggtg	aggacataca	94140
gaagaaagac	cagttgagac	tgtgcacgca	ggaggggtgtg	cagagcaagc	actgagggtgc	94200
agcacggaga	ccagagctgg	ccaggteccag	catcaccccc	acccccacat	caccagggca	94260
cactgcccaa	aagaacacct	aactgcggag	tgcagctctt	ttgtcaatct	gatggcatga	94320
agcaaccata	tgttctactt	ttttctactt	tttttaaatgt	cacaagtgtg	tagcagtgtc	94380
gtccctgtta	aggagttgtt	ttgaggggtgt	ttttaaaagt	tgtttttgag	tggctgtgga	94440
taaaaataca	tatttttgcc	gaaattttta	tgggtgttcct	gggctgtcct	gagaataagt	94500
tccattctga	tctaagcctc	tgatttttct	tcatagaaag	atgagctttg	cagacacaag	94560
cttggcagca	aggtgagaaa	ggccagccta	gtgagtcaag	ctatctgaaa	tgcattcctc	94620
ccagcgggca	ttccatccca	gcatacccta	tcagatatgt	gaaagagagg	aaccaagacc	94680
gaatgctatt	cctgcccagc	cctaataacc	actcacattc	tgaaatttaa	cttctttttt	94740
ttccctaaga	tagagatgtc	ctaactgaaa	atatgcctgt	atacaattta	ccctggaagt	94800
ctcagccatc	actcaaggga	agtctccaga	gggtgaagag	cctgtctggc	ctgtaggggt	94860
acacagtgtg	gggtggtcatt	ttaaatgggt	tccaagccaa	tgataggtcc	ctgaaatata	94920
acatgggtgga	aacttcta	aaagctcaca	tttgcattga	agtgttttagc	ttgttaagat	94980
aggcagttct	caaataaaaag	gtttgtttta	ttgggtaaat	gaccttgtag	ttttttgggtg	95040

acagagcata gaaagtaatt tcatgctgct cctgtgctat tgtttttgtg aagacagggga 95100  
 gctgtgaaaa actgcttagc tacctacatt cctcaataaa ggcatacagac agtaattggt 95160  
 gattacagat gttctccctg gaatggctgt tctcttgacc aagtagtcct acacttctgg 95220  
 aaggatcatt cagaactgtg gtctatgcca acccaaccagt agttcctgag tccctgcagt 95280  
 gctgagtgtc gggggccacc aagttgagta agacactgca gctctcaaag agttggatct 95340  
 aggattgtat tgtatcgatt tgtgggtgtt ggatatagtt tttccatgat cccctacgaa 95400  
 aatatgcctc tcatatgtaa gaatcatgcc tctccgtgt acacttttca gacactgaca 95460  
 aggaaggggtg ttcaatacag tactgaattt tcatatagct tttctggggg ggcāaaaata 95520  
 ccaaaatcaa cccatttcct acctttatct tgtccataaa attgttagaa atatcaaaat 95580  
 cccatttcat tccctgttaa atacatgtga acgttgtcta gacgtggag agcaaattct 95640  
 accacctctt ttgttcagca gtacatcaga cgattgcata gacgtgccag atggaaccaa 95700  
 ataataatgc acatggattt gtcataatcc gtacaagtca ttgacgcca cactgagcca 95760  
 ggtgctgtgg gagacaggac catgtgtgaa agagaagaca tgcttgcttc tataaaagca 95820  
 tcggtgttat tgaggagacc tgacattaat gcagaatagc aaatgaccat gcaaattaat 95880  
 tcaactactaa ctaagctgca ggttgacact cggaatgcag aggggcttca aagtgatgag 95940  
 ggtaggcct gagccaggcc ctgatgatgg gtggattttg aggatcagag agtacagctt 96000  
 agagagatac ccaagtggg accaccctt gccagtagg ctgacaaact aaggctcttg 96060  
 gtccctttt catattttgg gtgttctagt ggccagcca gagctagact tcgagtcag 96120  
 aattttctgg cacaagtgtt gtcacattca aaaaagtatt ttctttgtt gaaaaatgaa 96180  
 aaatatatat atatgtgtgt gtgtatatat atatgtgtgt gtgtgtgtat atatgtgtgt 96240  
 gtgtatatat acgtgtgtgt gtatatgtgt gtgtgtgtgt gtgtgtgtgt gtgtatatat 96300  
 gtgtatatat ataccatttt tcccaccta aatggagcat ggcaaactct gactggatta 96360  
 gtgagataga ccaagtcaca gagcactcca ggatgcagct gtgagctggg gaacaggcca 96420  
 gaaaggcctc agggacatca gcatacatgt tggagtcttct gcagttttct tagggaaccc 96480  
 tttaatgtca ctagagctaa cacacttgct acctgggaag caagcctgcc agagcaaatt 96540  
 agagagacga gggacagttg ctagaaagac acacctggaa gttctattta actagcatta 96600  
 attatgtgct aggtgctgag gatgtagact gagtgagatc ctcatctct ctctgtaggg 96660  
 tggaagagag gatattgctt gtctccatgg ctgtagtga acagtcagt agaccaggca 96720  
 cataaacctc ttagcagaac acttggcctt tctaaggact ccatatgtgt tccggggtaa 96780  
 atgcctgtgt ttcttgacgt agtgatgtct tgttctcta gacatcacta actttacaca 96840  
 gtagctttag atggcgtgga cgtgaataaa tgcaacttag gttttcttgt tggtttctt 96900  
 ttgagtatca ttgtgtttgt aaagaatttc agattagagg attgttacca cgtgggcctt 96960  
 caggaggaaa ctgttttgag tttttgtcag cccgaaatcg atttgtgct ttaagtatat 97020  
 gtgctcatca aaacaggcca ggctctgctg cagtaacaaa cttacaagtc tccgaggctt 97080  
 ggcacaacgg aagtctttgt gtcactgacg cccacttcag ctttgtgttg ctgaagcatt 97140  
 ctttggcct atccgagctg tccctctggt ggtgggtgct gggggtttgg gttccctctg 97200  
 tgttgatgc caccatctca gcacggctt ccacagcagc catagcagga gaagaaaatg 97260  
 ctgggggctc tcagggtgct ttaagggcc tggccaccga cctgcaagggt gtgcgagttg 97320  
 ccttctgtg tggccagaac tgatgataaa ctgtagactc atccctgctg aaactcggct 97380  
 ccagagtgt cccaaggctg gacagcgtgt gggcactgga tcccacctgt gttagcactg 97440  
 gcaattgtat tctcatttct tcttttatct tccaggtggc aaatggtagg atccaaagcc 97500  
 ttgaggccac cattgagaag ctctgagca gtgagagcaa gctgaagcag gccatgttta 97560

ccttagaact	ggagcgggtcg	gccctgctgc	agacgggtgga	ggagctgcgg	cggcggagcg	97620
cagagcccgag	cgaccggggag	cctgagtga	cgcagcccgga	gcccacgggc	gactgacagc	97680
tctgcaggag	agattgcaac	accatcccac	actgtccagg	ccttaactga	gagggacaga	97740
agacgctgga	aggagagaag	gaagcgggaa	gtgtgcttct	cagggaggaa	accggcttgc	97800
cagcaagtag	attcttacga	actccaactt	gcaattcagg	gggcatgtcc	cagtgttttt	97860
tttgttgttt	ttagatacta	aatcgctccct	tctccagtcc	tgattactgt	acacagtagc	97920
tttagatggc	gtggacgtga	ataaatgcaa	cttatgtttt	cttggttggtt	ccttttttgag	97980
tgtcactgtg	tttgtaaaga	gcattcacia	tacgggtgaa	tttcaaaagc	tggaagagct	98040
cgagatcatg	cctcaggcaa	aggcgtgggt	ccatcgttct	tccgagagg	tttgtgtggc	98100
gactacaccc	tcagcgtccc	tggaagggtg	cagttggctc	tgcgccattc	ttgttatgga	98160
aacctaaagt	gatcattggg	aagatcagtg	atcttgggtc	attgatccct	ggctcagagg	98220
atagcggttt	ccatcataaa	ccaagatgat	gagttcagcc	tttatccctc	gtggttccac	98280
tagatgtaac	ttaaaggagt	taacatttga	ggactttgtt	ctacatcaga	ttttactatt	98340
tgaatgttta	agatcacttt	attgaatttg	aagatcatca	aattaaataa	aatgatattat	98400
ttaatttgga	tatcctgata	actgtcaagt	gaaatggatc	tctctctttg	gtattttaagg	98460
aagtttgtct	ttaaaaaaaa	aatagagtgt	tttcatacat	ttttgcttat	cccataagta	98520
cagttgatca	aagtcatagt	aggtaaagtgc	tttatgggac	agctgacacc	ttttagaccc	98580
taccaggat	tgctagcatg	tgagctgcag	ttgtggggctc	tgagatattt	ctttgtggta	98640
gtttcatacc	catactatag	agtcattgtat	ttatttttgc	ctgttgtgtg	atgtaatgca	98700
atcatgttcc	tttgagtctc	catcccttgg	aaatctgact	tcttgacaga	ggagtaggca	98760
catcaagata	ttcaggggtg	ccccaaagag	ctgggacttt	caaaaaaaaa	agatcaggct	98820
nnaactgcag	tcagatttat	gacagctgac	agtttttcag	aggctgcaca	cagtgactct	98880
cctctctcag	gatgacgagg	acctgtgcct	tcaacaagca	aaatgctgct	cacggttgtc	98940
ctgcttgcag	ccagtcactg	tgtaaagcct	ctctgatgtg	cacttaagag	tggtgtgctt	99000
tctcaciaag	atgggggttct	gtgcagtcac	aggctacttc	cttgacaaca	caatcatttc	99060
tgatctttat	cactgtaacc	acgtcttcta	ttccatagga	gtttcttttg	attctctcag	99120
ttgcgggggg	catctctttaa	tcctgggggtg	aaaggagaga	ttgccatact	tagactcact	99180
gtgagtctcc	ccggccattt	cacgaggaga	ccacagtgtc	gccaccagtg	cctaaacagg	99240
tggtgtggcat	tcgagacttc	ctcctgttcc	ctgggtcaga	ggatagcggg	ttccatcata	99300
aaccaagatg	atgagttcag	cctttatccc	tcgtgggttc	gctagatgta	acttatagga	99360
gttaacattt	gaggactttg	ttctgcatca	gatcttacta	tttgaatgtt	tactgttgga	99420
ttttggggcat	cttattactg	ttactcaaaa	acattgactc	tgcatcaaga	agaaaacaag	99480
aaagcaataa	aacaagaaat	aattcatgct	cacattttta	tggtgggtttt	tttttttttt	99540
ttacttttgg	atttttgctt	ttcagcccag	gagtaaagga	atgccttatg	aacacctgtg	99600
gcctacgtgt	ggcatgacc	caaccatcag	tgagattatt	tgagatattg	gtgtctgcat	99660
ccagtgttgt	tatctgagtg	tttattacgt	aagttgtaac	acctctacac	agggtgtgag	99720
tttagcactg	atgagaccag	ctccatcatt	gtatgtggca	gtgagtcctg	ttacgagatt	99780
gggttgggca	gaaaggactg	ttgacatgag	cctgtggatg	taggttggac	agtctcagcc	99840
tgtgactgac	taggcaagga	gcggagaggc	aactgtgtga	ggattctcag	agccaaattt	99900
ttaagccatg	ttttgggtta	tatttcccc	aacactcatt	tgtgcacttg	gtgggtgtcaa	99960

95

cag	cgt	gtt	cac	aaa	ctg	att	cac	aac	agt	cat	gac	cca	agt	tac	ttt	512
Gln	Arg	Val	His	Lys	Leu	Ile	His	Asn	Ser	His	Asp	Pro	Ser	Tyr	Phe	
100						105					110					
gct	tgt	ctg	att	aag	gaa	gac	gct	gtc	cac	cgg	cag	agt	atc	tgc	tat	560
Ala	Cys	Leu	Ile	Lys	Glu	Asp	Ala	Val	His	Arg	Gln	Ser	Ile	Cys	Tyr	
115					120					125					130	
gtg	ttc	aaa	gcc	gat	gat	caa	aca	aaa	gtg	cct	gag	atc	atc	agc	tcc	608
Val	Phe	Lys	Ala	Asp	Asp	Gln	Thr	Lys	Val	Pro	Glu	Ile	Ile	Ser	Ser	
				135					140					145		
atc	cgt	cag	gcg	ggg	aag	atc	gcc	cgg	cag	gag	gag	ctg	cac	tgc	ccg	656
Ile	Arg	Gln	Ala	Gly	Lys	Ile	Ala	Arg	Gln	Glu	Glu	Leu	His	Cys	Pro	
			150						155					160		
tcc	gag	ttc	gac	gac	acg	ttt	tcc	aag	aag	ttc	gag	gtg	ctc	ttc	tgc	704
Ser	Glu	Phe	Asp	Asp	Thr	Phe	Ser	Lys	Lys	Phe	Glu	Val	Leu	Phe	Cys	
		165						170					175			
ggc	cgc	gtg	acg	gtg	gcg	cac	aag	aag	gct	ccg	ccg	gcc	ctg	atc	gac	752
Gly	Arg	Val	Thr	Val	Ala	His	Lys	Lys	Ala	Pro	Pro	Ala	Leu	Ile	Asp	
	180					185					190					
gag	tgc	atc	gag	aag	ttc	aat	cac	gtc	agc	ggc	agc	cgg	ggg	tcc	gag	800
Glu	Cys	Ile	Glu	Lys	Phe	Asn	His	Val	Ser	Gly	Ser	Arg	Gly	Ser	Glu	
195					200					205					210	
agc	ccc	cgc	ccc	aac	ccg	ccc	cat	gcc	gcg	ccc	aca	ggg	agc	cag	gag	848
Ser	Pro	Arg	Pro	Asn	Pro	Pro	His	Ala	Ala	Pro	Thr	Gly	Ser	Gln	Glu	
				215					220					225		
cct	gtg	cgc	agg	ccc	atg	cgc	aag	tcc	ttc	tcc	cag	ccc	ggc	ctg	cgc	896
Pro	Val	Arg	Arg	Pro	Met	Arg	Lys	Ser	Phe	Ser	Gln	Pro	Gly	Leu	Arg	
			230					235					240			
tcg	ctg	gcc	ttt	agg	aag	gag	ctg	cag	gat	ggg	ggc	ctc	cga	agc	agc	944
Ser	Leu	Ala	Phe	Arg	Lys	Glu	Leu	Gln	Asp	Gly	Gly	Leu	Arg	Ser	Ser	
		245						250				255				
ggc	ttc	ttc	agc	tcc	ttc	gag	gag	agc	gac	att	gag	aac	cac	ctc	att	992
Gly	Phe	Phe	Ser	Ser	Phe	Glu	Glu	Ser	Asp	Ile	Glu	Asn	His	Leu	Ile	
	260					265					270					
agc	gga	cac	aat	att	gtg	cag	ccc	aca	gat	atc	gag	gaa	aat	cga	act	1040
Ser	Gly	His	Asn	Ile	Val	Gln	Pro	Thr	Asp	Ile	Glu	Glu	Asn	Arg	Thr	
275					280					285					290	
atg	ctc	ttc	acg	att	ggc	cag	tct	gaa	gtt	tac	ctc	atc	agt	cct	gac	1088
Met	Leu	Phe	Thr	Ile	Gly	Gln	Ser	Glu	Val	Tyr	Leu	Ile	Ser	Pro	Asp	
				295					300					305		



tct cag ggc atc aga cac gtg gac cac ttt ggg ttt atc tgt cgg gag	1184
Ser Gln Gly Ile Arg His Val Asp His Phe Gly Phe Ile Cys Arg Glu	
325 330 335	
tct tcc gga ggt ggc ggc ttt cat ttt gtc tgt tac gtg ttt cag tgc	1232
Ser Ser Gly Gly Gly Gly Phe His Phe Val Cys Tyr Val Phe Gln Cys	
340 345 350	
aca aat gag gct ctg gtt gat gaa att atg atg acc ctg aaa cag gcc	1280
Thr Asn Glu Ala Leu Val Asp Glu Ile Met Met Thr Leu Lys Gln Ala	
355 360 365 370	
ttc acg gtg gcc gca gtg cag cag aca gct aag gcg cca gcc cag ctg	1328
Phe Thr Val Ala Ala Val Gln Gln Thr Ala Lys Ala Pro Ala Gln Leu	
375 380 385	
tgt gag ggc tgc ccc ctg caa agc ctg cac aag ctc tgt gag agg ata	1376
Cys Glu Gly Cys Pro Leu Gln Ser Leu His Lys Leu Cys Glu Arg Ile	
390 395 400	
gag gga atg aat tct tcc aaa aca aaa cta gaa ctg caa aag cac ctg	1424
Glu Gly Met Asn Ser Ser Lys Thr Lys Leu Glu Leu Gln Lys His Leu	
405 410 415	
acg aca tta acc aat cag gag cag gcg act att ttt gaa gag gtt cag	1472
Thr Thr Leu Thr Asn Gln Glu Gln Ala Thr Ile Phe Glu Glu Val Gln	
420 425 430	
aaa ttg aga ccg aga aat gag cag cga gag aat gaa ttg att att tct	1520
Lys Leu Arg Pro Arg Asn Glu Gln Arg Glu Asn Glu Leu Ile Ile Ser	
435 440 445 450	
ttt ctg aga tgt tta tat gaa gag aaa cag aaa gaa cac atc cat att	1568
Phe Leu Arg Cys Leu Tyr Glu Glu Lys Gln Lys Glu His Ile His Ile	
455 460 465	
ggg gag atg aag cag aca tcg cag atg gca gca gag aat att gga agt	1616
Gly Glu Met Lys Gln Thr Ser Gln Met Ala Ala Glu Asn Ile Gly Ser	
470 475 480	
gaa tta cca ccc agt gcc act cga ttt agg cta gat atg ctg aaa aac	1664
Glu Leu Pro Pro Ser Ala Thr Arg Phe Arg Leu Asp Met Leu Lys Asn	
485 490 495	
aaa gca aag aga tct tta aca gag tct tta gaa agt att ttg tcc cgg	1712
Lys Ala Lys Arg Ser Leu Thr Glu Ser Leu Glu Ser Ile Leu Ser Arg	
500 505 510	
ggt aat aaa gcc aga ggc ctg cag gaa cac tcc atc agt gtg gat ctg	1760
Gly Asn Lys Ala Arg Gly Leu Gln Glu His Ser Ile Ser Val Asp Leu	
515 520 525 530	
gat agc tcc ctg tct agt aca tta agt aac acc agc aaa gag cca tct	1808
Asp Ser Ser Leu Ser Ser Thr Leu Ser Asn Thr Ser Lys Glu Pro Ser	
535 540 545	

gtg tgt gaa aag gag gcc ttg ccc atc tct gag agc tcc ttt aag ctc 1856  
 Val Cys Glu Lys Glu Ala Leu Pro Ile Ser Glu Ser Ser Phe Lys Leu  
 550 555 560  
 ctc ggc tcc tcg gag gac ctg tcc agt gac tcg gag agt cat ctc cca 1904  
 Leu Gly Ser Ser Glu Asp Leu Ser Ser Asp Ser Glu Ser His Leu Pro  
 565 570 575  
 gaa gag cca gct ccg ctg tcg ccc cag cag gcc ttc agg agg cga gca 1952  
 Glu Glu Pro Ala Pro Leu Ser Pro Gln Gln Ala Phe Arg Arg Arg Ala  
 580 585 590  
 aac acc ctg agt cac ttc ccc atc gaa tgc cag gaa cct cca caa cct 2000  
 Asn Thr Leu Ser His Phe Pro Ile Glu Cys Gln Glu Pro Pro Gln Pro  
 595 600 605 610  
 gcc cgg ggg tcc ccg ggg gtt tcg caa agg aaa ctt atg agg tat cac 2048  
 Ala Arg Gly Ser Pro Gly Val Ser Gln Arg Lys Leu Met Arg Tyr His  
 615 620 625  
 tca gtg agc aca gag acg cct cat gaa cga aag gac ttt gaa tcc aaa 2096  
 Ser Val Ser Thr Glu Thr Pro His Glu Arg Lys Asp Phe Glu Ser Lys  
 630 635 640  
 gca aac cat ctt ggt gat tct ggt ggg act cct gtg aag acc cgg agg 2144  
 Ala Asn His Leu Gly Asp Ser Gly Gly Thr Pro Val Lys Thr Arg Arg  
 645 650 655  
 cat tcc tgg agg cag cag ata ttc ctc cga gta gcc acc ccg cag aag 2192  
 His Ser Trp Arg Gln Gln Ile Phe Leu Arg Val Ala Thr Pro Gln Lys  
 660 665 670  
 gcg tgc gat tct tcc agc aga tat gaa gat tat tca gag ctg gga gag 2240  
 Ala Cys Asp Ser Ser Ser Arg Tyr Glu Asp Tyr Ser Glu Leu Gly Glu  
 675 680 685 690  
 ctt ccc cca cga tct cct tta gaa cca gtt tgt gaa gat ggg ccc ttt 2288  
 Leu Pro Pro Arg Ser Pro Leu Glu Pro Val Cys Glu Asp Gly Pro Phe  
 695 700 705  
 ggc ccc cca cca gag gaa aag aaa agg aca tct cgt gag ctc cga gag 2336  
 Gly Pro Pro Pro Glu Glu Lys Lys Arg Thr Ser Arg Glu Leu Arg Glu  
 710 715 720  
 ctg tgg caa aag gct att ctt caa cag ata ctg ctg ctt aga atg gag 2384  
 Leu Trp Gln Lys Ala Ile Leu Gln Gln Ile Leu Leu Leu Arg Met Glu  
 725 730 735  
 aag gaa aat cag aag ctc caa gcc tct gaa aat gat ttg ctg aac aag 2432  
 Lys Glu Asn Gln Lys Leu Gln Ala Ser Glu Asn Asp Leu Leu Asn Lys  
 740 745 750  
 cgc ctg aag ctc gat tat gaa gaa att act ccc tgt ctt aaa gaa gta 2480  
 Arg Leu Lys Leu Asp Tyr Glu Glu Ile Thr Pro Cys Leu Lys Glu Val  
 755 760 765 770

095290-1-06501

act	aca	gtg	tgg	gaa	aag	atg	ctt	agc	act	cca	gga	aga	tca	aaa	att		2528
Thr	Thr	Val	Trp	Glu	Lys	Met	Leu	Ser	Thr	Pro	Gly	Arg	Ser	Lys	Ile		
				775					780					785			
aag	ttt	gac	atg	gaa	aaa	atg	cac	tcg	gct	gtt	ggg	caa	ggg	gtg	cca		2576
Lys	Phe	Asp	Met	Glu	Lys	Met	His	Ser	Ala	Val	Gly	Gln	Gly	Val	Pro		
				790					795					800			
cgt	cat	cac	cga	ggg	gaa	atc	tgg	aaa	ttt	cta	gct	gag	caa	ttc	cac		2624
Arg	His	His	Arg	Gly	Glu	Ile	Trp	Lys	Phe	Leu	Ala	Glu	Gln	Phe	His		
				805					810					815			
ctt	aaa	cac	cag	ttt	ccc	agc	aaa	cag	cag	cca	aag	gat	gtg	cca	tac		2672
Leu	Lys	His	Gln	Phe	Pro	Ser	Lys	Gln	Gln	Pro	Lys	Asp	Val	Pro	Tyr		
	820							825						830			
aaa	gaa	ctc	tta	aag	cag	ctg	act	tcc	cag	cag	cat	gcg	att	ctt	att		2720
Lys	Glu	Leu	Leu	Lys	Gln	Leu	Thr	Ser	Gln	Gln	His	Ala	Ile	Leu	Ile		
	835					840				845					850		
gac	ctt	ggg	cga	acc	ttt	cct	aca	cac	cca	tac	ttc	tct	gcc	cag	ctt		2768
Asp	Leu	Gly	Arg	Thr	Phe	Pro	Thr	His	Pro	Tyr	Phe	Ser	Ala	Gln	Leu		
				855					860					865			
gga	gca	gga	cag	cta	tcg	ctt	tac	aac	att	ttg	aag	gcc	tac	tca	ctt		2816
Gly	Ala	Gly	Gln	Leu	Ser	Leu	Tyr	Asn	Ile	Leu	Lys	Ala	Tyr	Ser	Leu		
				870					875					880			
cta	gac	cag	gaa	gtg	gga	tat	tgc	caa	ggg	ctc	agc	ttt	gta	gca	ggc		2864
Leu	Asp	Gln	Glu	Val	Gly	Tyr	Cys	Gln	Gly	Leu	Ser	Phe	Val	Ala	Gly		
				885					890					895			
att	ttg	ctt	ctt	cat	atg	agt	gag	gaa	gag	gcg	ttt	aaa	atg	ctc	aag		2912
Ile	Leu	Leu	Leu	His	Met	Ser	Glu	Glu	Glu	Ala	Phe	Lys	Met	Leu	Lys		
	900							905						910			
ttt	ctg	atg	ttt	gac	atg	ggg	ctg	cgg	aaa	cag	tat	cgg	cca	gac	atg		2960
Phe	Leu	Met	Phe	Asp	Met	Gly	Leu	Arg	Lys	Gln	Tyr	Arg	Pro	Asp	Met		
	915					920					925				930		
att	att	tta	cag	atc	cag	atg	tac	cag	ctc	tcg	agg	ttg	ctt	cat	gat		3008
Ile	Ile	Leu	Gln	Ile	Gln	Met	Tyr	Gln	Leu	Ser	Arg	Leu	Leu	His	Asp		
				935					940					945			
tac	cac	aga	gac	ctc	tac	aat	cac	ctg	gag	gag	cac	gag	atc	ggc	ccc		3056
Tyr	His	Arg	Asp	Leu	Tyr	Asn	His	Leu	Glu	Glu	His	Glu	Ile	Gly	Pro		
				950					955					960			
agc	ctc	tac	gct	gcc	ccc	tgg	ttc	ctc	acc	atg	ttt	gcc	tca	cag	ttc		3104
Ser	Leu	Tyr	Ala	Ala	Pro	Trp	Phe	Leu	Thr	Met							

aca gag gtc ata ttt aaa gtg gct tta agt ctg ttg gga agc cat aag 3200  
 Thr Glu Val Ile Phe Lys Val Ala Leu Ser Leu Leu Gly Ser His Lys  
 995 1000 1005 1010  
 ccc ttg att ctg cag cat gaa aac cta gaa acc ata gtt gac ttt ata 3248  
 Pro Leu Ile Leu Gln His Glu Asn Leu Glu Thr Ile Val Asp Phe Ile  
 1015 1020 1025  
 aaa agc acg cta ccc aac ctt ggc ttg gta cag atg gaa aag acc atc 3296  
 Lys Ser Thr Leu Pro Asn Leu Gly Leu Val Gln Met Glu Lys Thr Ile  
 1030 1035 1040  
 aat cag gta ttt gaa atg gac atc gct aaa cag tta caa gct tat gaa 3344  
 Asn Gln Val Phe Glu Met Asp Ile Ala Lys Gln Leu Gln Ala Tyr Glu  
 1045 1050 1055  
 gtt gag tac cac gtc ctt caa gaa gaa ctt atc gat tcc tct cct ctc 3392  
 Val Glu Tyr His Val Leu Gln Glu Glu Leu Ile Asp Ser Ser Pro Leu  
 1060 1065 1070  
 agt gac aac caa aga atg gat aaa tta gag aaa acc aac agc agc tta 3440  
 Ser Asp Asn Gln Arg Met Asp Lys Leu Glu Lys Thr Asn Ser Ser Leu  
 1075 1080 1085 1090  
 cgc aaa cag aac ctt gac ctc ctt gaa cag ttg cag gtg gca aat ggt 3488  
 Arg Lys Gln Asn Leu Asp Leu Leu Glu Gln Leu Gln Val Ala Asn Gly  
 1095 1100 1105  
 agg atc caa agc ctt gag gcc acc att gag aag ctc ctg agc agt gag 3536  
 Arg Ile Gln Ser Leu Glu Ala Thr Ile Glu Lys Leu Leu Ser Ser Glu  
 1110 1115 1120  
 agc aag ctg aag cag gcc atg ctt acc tta gaa ctg gag cgg tgg gcc 3584  
 Ser Lys Leu Lys Gln Ala Met Leu Thr Leu Glu Leu Glu Arg Ser Ala  
 1125 1130 1135  
 ctg ctg cag acg gtg gag gag ctg cgg cgg cgg agc gca gag ccc agc 3632  
 Leu Leu Gln Thr Val Glu Glu Leu Arg Arg Arg Ser Ala Glu Pro Ser  
 1140 1145 1150  
 gac cgg gag cct gag tgc acg cag ccc gag ccc acg ggc gac tga 3677  
 Asp Arg Glu Pro Glu Cys Thr Gln Pro Glu Pro Thr Gly Asp \*  
 1155 1160 1165  
 cagctctgca ggagagattg caacaccatc ccacactgtc caggccttaa ctgagagggga 3737  
 cagaagacgc tggaaggaga gaaggaagcg ggaagtgtgc ttctcagggga ggaaaccggc 3797  
 ttgccagcaa gtagattctt acgaactcca acttgcaatt caggggggcat gtcccagtgt 3857  
 tttttttgtt gtttttagat actaaatcgt cccttctcca gtcttgatta ctgtacacag 3917  
 tagctttaga tggcgtggac gtgaataaat gcaacttatg ttttaaaaaa aaaaaaaaaa 3977  
 aaaaaa 3983

&lt;210&gt; 4

&lt;211&gt; 3988

T 05290 "T E 5290"

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; CDS

&lt;222&gt; 176..3730

&lt;220&gt;

&lt;221&gt; polyA\_signal

&lt;222&gt; 3947..3952

&lt;223&gt; AATAAA

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;222&gt; 1..458

&lt;223&gt; homology with Genset 5' EST in ref : A35235

&lt;400&gt; 4

ataataggca ctgaagacat gttaatggaa ggtggatttg tgattcagaa cctctagact	60
acctgggcga gtctttttaa atgtttctgc atatgaagtg tgtaaaatag attgcttgat	120
ccaaaacaga aaaacagtga taactgtttt gctgagttcc cagacccttc ccaag atg	178
Met	
1	
gaa cca ata aca ttc aca gca agg aaa cat ctg ctt cct aac gag gtc	226
Glu Pro Ile Thr Phe Thr Ala Arg Lys His Leu Leu Pro Asn Glu Val	
5 10 15	
tcg gtg gat ttt ggc ctg cag ctg gtg ggc tcc ctg cct gtg cat tcc	274
Ser Val Asp Phe Gly Leu Gln Leu Val Gly Ser Leu Pro Val His Ser	
20 25 30	
ctg acc acc atg ccc atg ctg ccc tgg gtt gtg gct gag gtg cga aga	322
Leu Thr Thr Met Pro Met Leu Pro Trp Val Val Ala Glu Val Arg Arg	
35 40 45	
ctc agc agg cag tcc acc aga aag gaa cct gta acc aag caa gtc cgg	370
Leu Ser Arg Gln Ser Thr Arg Lys Glu Pro Val Thr Lys Gln Val Arg	
50 55 60 65	
ctt tgc gtt tca ccc tct gga ctg aga tgt gaa cct gag cca ggg aga	418
Leu Cys Val Ser Pro Ser Gly Leu Arg Cys Glu Pro Glu Pro Gly Arg	
70 75 80	
agt caa cag tgg gat ccc ctg atc tat tcc agc atc ttt gag tgc aag	466
Ser Gln Gln Trp Asp Pro Leu Ile Tyr Ser Ser Ile Phe Glu Cys Lys	
85 90 95	
cct cag cgt gtt cac aaa ctg att cac aac agt cat gac cca agt tac	514

09694062604

Pro	Gln	Arg	Val	His	Lys	Leu	Ile	His	Asn	Ser	His	Asp	Pro	Ser	Tyr					
100								105								110				
ttt	gct	tgt	ctg	att	aag	gaa	gac	gct	gtc	cac	cgg	cag	agt	atc	tgc	562				
Phe	Ala	Cys	Leu	Ile	Lys	Glu	Asp	Ala	Val	His	Arg	Gln	Ser	Ile	Cys					
115								120								125				
tat	gtg	ttc	aaa	gcc	gat	gat	caa	aca	aaa	gtg	cct	gag	atc	atc	agc	610				
Tyr	Val	Phe	Lys	Ala	Asp	Asp	Gln	Thr	Lys	Val	Pro	Glu	Ile	Ile	Ser					
130								135								140				145
tcc	atc	cgt	cag	gcg	ggg	aag	atc	gcc	cgg	cag	gag	gag	ctg	cac	tgc	658				
Ser	Ile	Arg	Gln	Ala	Gly	Lys	Ile	Ala	Arg	Gln	Glu	Glu	Leu	His	Cys					
150												155				160				
ccg	tcc	gag	ttc	gac	gac	acg	ttt	tcc	aag	aag	ttc	gag	gtg	ctc	ttc	706				
Pro	Ser	Glu	Phe	Asp	Asp	Thr	Phe	Ser	Lys	Lys	Phe	Glu	Val	Leu	Phe					
165												170				175				
tgc	ggc	cgc	gtg	acg	gtg	gcg	cac	aag	aag	gct	ccg	ccg	gcc	ctg	atc	754				
Cys	Gly	Arg	Val	Thr	Val	Ala	His	Lys	Lys	Ala	Pro	Pro	Ala	Leu	Ile					
180												185				190				
gac	gag	tgc	atc	gag	aag	ttc	aat	cac	gtc	agc	ggc	agc	cgg	ggg	tcc	802				
Asp	Glu	Cys	Ile	Glu	Lys	Phe	Asn	His	Val	Ser	Gly	Ser	Arg	Gly	Ser					
195												200				205				
gag	agc	ccc	cgc	ccc	aac	ccg	ccc	cat	gcc	gcg	ccc	aca	ggg	agc	cag	850				
Glu	Ser	Pro	Arg	Pro	Asn	Pro	Pro	His	Ala	Ala	Pro	Thr	Gly	Ser	Gln					
210												215				220				225
gag	cct	gtg	cgc	agg	ccc	atg	cgc	aag	tcc	ttc	tcc	cag	ccc	ggc	ctg	898				
Glu	Pro	Val	Arg	Arg	Pro	Met	Arg	Lys	Ser	Phe	Ser	Gln	Pro	Gly	Leu					
230												235				240				
cgc	tcg	ctg	gcc	ttt	agg	aag	gag	ctg	cag	gat	ggg	ggc	ctc	cga	agc	946				
Arg	Ser	Leu	Ala	Phe	Arg	Lys	Glu	Leu	Gln	Asp	Gly	Gly	Leu	Arg	Ser					
245												250				255				
agc	ggc	ttc	ttc	agc	tcc	ttc	gag	gag	agc	gac	att	gag	aac	cac	ctc	994				
Ser	Gly	Phe	Phe	Ser	Ser	Phe	Glu	Glu	Ser	Asp	Ile	Glu	Asn	His	Leu					
260												265				270				
att	agc	gga	cac	aat	att	gtg	cag	ccc	aca	gat	atc	gag	gaa	aat	cga	1042				
Ile	Ser	Gly	His	Asn	Ile	Val	Gln	Pro	Thr	Asp	Ile	Glu	Glu	Asn	Arg					
275												280				285				
act	atg	ctc	ttc	acg	att	ggc	cag	tct	gaa	gtt	tac	ctc	atc	agt	cct	1090				
Thr	Met	Leu	Phe	Thr	Ile	Gly	Gln	Ser	Glu	Val	Tyr	Leu	Ile	Ser	Pro					
290												295				300				305
gac	acc	aaa	aaa	ata	gca	ttg	gag	aaa	aat	ttt	aag	gag	ata	tcc	ttt	1138				
Asp	Thr	Lys	Lys	Ile	Ala	Leu	Glu	Lys	Asn	Phe	Lys	Glu	Ile	Ser	Phe					
310												315				320				
tgc	tct	cag	ggc	atc	aga	cac	gtg	gac	cac	ttt	ggg	ttt	atc	tgt	cgg	1186				

Cys Ser Gln Gly Ile Arg His Val Asp His Phe Gly Phe Ile Cys Arg  
 325 330 335  
 gag tct tcc gga ggt ggc ggc ttt cat ttt gtc tgt tac gtg ttt cag 1234  
 Glu Ser Ser Gly Gly Gly Gly Phe His Phe Val Cys Tyr Val Phe Gln  
 340 345 350  
 tgc aca aat gag gct ctg gtt gat gaa att atg atg acc ctg aaa cag 1282  
 Cys Thr Asn Glu Ala Leu Val Asp Glu Ile Met Met Thr Leu Lys Gln  
 355 360 365  
 gcc ttc acg gtg gcc gca gtg cag cag aca gct aag gcg cca gcc cag 1330  
 Ala Phe Thr Val Ala Ala Val Gln Gln Thr Ala Lys Ala Pro Ala Gln  
 370 375 380 385  
 ctg tgt gag ggc tgc ccc ctg caa agc ctg cac aag ctc tgt gag agg 1378  
 Leu Cys Glu Gly Cys Pro Leu Gln Ser Leu His Lys Leu Cys Glu Arg  
 390 395 400  
 ata gag gga atg aat tct tcc aaa aca aaa cta gaa ctg caa aag cac 1426  
 Ile Glu Gly Met Asn Ser Ser Lys Thr Lys Leu Glu Leu Gln Lys His  
 405 410 415  
 ctg acg aca tta acc aat cag gag cag gcg act att ttt gaa gag gtt 1474  
 Leu Thr Thr Leu Thr Asn Gln Glu Gln Ala Thr Ile Phe Glu Glu Val  
 420 425 430  
 cag aaa ttg aga ccg aga aat gag cag cga gag aat gaa ttg att att 1522  
 Gln Lys Leu Arg Pro Arg Asn Glu Gln Arg Glu Asn Glu Leu Ile Ile  
 435 440 445  
 tct ttt ctg aga tgt tta tat gaa gag aaa cag aaa gaa cac atc cat 1570  
 Ser Phe Leu Arg Cys Leu Tyr Glu Glu Lys Gln Lys Glu His Ile His  
 450 455 460 465  
 att ggg gag atg aag cag aca tcg cag atg gca gca gag aat att gga 1618  
 Ile Gly Glu Met Lys Gln Thr Ser Gln Met Ala Ala Glu Asn Ile Gly  
 470 475 480  
 agt gaa tta cca ccc agt gcc act cga ttt agg cta gat atg ctg aaa 1666  
 Ser Glu Leu Pro Pro Ser Ala Thr Arg Phe Arg Leu Asp Met Leu Lys  
 485 490 495  
 aac aaa gca aag aga tct tta aca gag tct tta gaa agt att ttg tcc 1714  
 Asn Lys Ala Lys Arg Ser Leu Thr Glu Ser Leu Glu Ser Ile Leu Ser  
 500 505 510  
 cgg ggt aat aaa gcc aga ggc ctg cag gaa cac tcc atc agt gtg gat 1762  
 Arg Gly Asn Lys Ala Arg Gly Leu Gln Glu His Ser Ile Ser Val Asp  
 515 520 525  
 ctg gat agc tcc ctg tct agt aca tta agt aac acc agc aaa gag cca 1810  
 Leu Asp Ser Ser Leu Ser Ser Thr Leu Ser Asn Thr Ser Lys Glu Pro  
 530 535 540 545  
 tct gtg tgt gaa aag gag gcc ttg ccc atc tct gag agc tcc ttt aag 1858

09234 06504  
 05230 06230

Ser Val Cys Glu Lys Glu Ala Leu Pro Ile Ser Glu Ser Ser Phe Lys	
550 555 560	
ctc ctc ggc tcc tcg gag gac ctg tcc agt gac tcg gag agt cat ctc	1906
Leu Leu Gly Ser Ser Glu Asp Leu Ser Ser Asp Ser Glu Ser His Leu	
565 570 575	
cca gaa gag cca gct ccg ctg tcg ccc cag cag gcc ttc agg agg cga	1954
Pro Glu Glu Pro Ala Pro Leu Ser Pro Gln Gln Ala Phe Arg Arg Arg	
580 585 590	
gca aac acc ctg agt cac ttc ccc atc gaa tgc cag gaa cct cca caa	2002
Ala Asn Thr Leu Ser His Phe Pro Ile Glu Cys Gln Glu Pro Pro Gln	
595 600 605	
cct gcc cgg ggg tcc ccg ggg gtt tcg caa agg aaa ctt atg agg tat	2050
Pro Ala Arg Gly Ser Pro Gly Val Ser Gln Arg Lys Leu Met Arg Tyr	
610 615 620 625	
cac tca gtg agc aca gag acg cct cat gaa cga aag gac ttt gaa tcc	2098
His Ser Val Ser Thr Glu Thr Pro His Glu Arg Lys Asp Phe Glu Ser	
630 635 640	
aaa gca aac cat ctt ggt gat tct ggt ggg act cct gtg aag acc cgg	2146
Lys Ala Asn His Leu Gly Asp Ser Gly Gly Thr Pro Val Lys Thr Arg	
645 650 655	
agg cat tcc tgg agg cag cag ata ttc ctc cga gta gcc acc ccg cag	2194
Arg His Ser Trp Arg Gln Gln Ile Phe Leu Arg Val Ala Thr Pro Gln	
660 665 670	
aag gcg tgc gat tct tcc agc aga tat gaa gat tat tca gag ctg gga	2242
Lys Ala Cys Asp Ser Ser Ser Arg Tyr Glu Asp Tyr Ser Glu Leu Gly	
675 680 685	
gag ctt ccc cca cga, tct cct tta gaa cca gtt tgt gaa gat ggg ccc	2290
Glu Leu Pro Pro Arg Ser Pro Leu Glu Pro Val Cys Glu Asp Gly Pro	
690 695 700 705	
ttt ggc ccc cca cca gag gaa aag aaa agg aca tct cgt gag ctc cga	2338
Phe Gly Pro Pro Pro Glu Glu Lys Lys Arg Thr Ser Arg Glu Leu Arg	
710 715 720	
gag ctg tgg caa aag gct att ctt caa cag ata ctg ctg ctt aga atg	2386
Glu Leu Trp Gln Lys Ala Ile Leu Gln Gln Ile Leu Leu Leu Arg Met	
725 730 735	
gag aag gaa aat cag aag ctc caa gcc tct gaa aat gat ttg ctg aac	2434
Glu Lys Glu Asn Gln Lys Leu Gln Ala Ser Glu Asn Asp Leu Leu Asn	
740 745 750	
aag cgc ctg aag ctc gat tat gaa gaa att act ccc tgt ctt aaa gaa	2482
Lys Arg Leu Lys Leu Asp Tyr Glu Glu Ile Thr Pro Cys Leu Lys Glu	
755 760 765	
gta act aca gtg tgg gaa aag atg ctt agc act cca gga aga tca aaa	2530

FO9290:FE29260



Val Thr Thr Val Trp Glu Lys Met Leu Ser Thr Pro Gly Arg Ser Lys  
 770 775 780 785  
 att aag ttt gac atg gaa aaa atg cac tcg gct gtt ggg caa ggt gtg 2578  
 Ile Lys Phe Asp Met Glu Lys Met His Ser Ala Val Gly Gln Gly Val  
 790 795 800  
 cca cgt cat cac cga ggt gaa atc tgg aaa ttt cta gct gag caa ttc 2626  
 Pro Arg His His Arg Gly Glu Ile Trp Lys Phe Leu Ala Glu Gln Phe  
 805 810 815  
 cac ctt aaa cac cag ttt ccc agc aaa cag cag cca aag gat gtg cca 2674  
 His Leu Lys His Gln Phe Pro Ser Lys Gln Gln Pro Lys Asp Val Pro  
 820 825 830  
 tac aaa gaa ctc tta aag cag ctg act tcc cag cag cat gcg att ctt 2722  
 Tyr Lys Glu Leu Leu Lys Gln Leu Thr Ser Gln Gln His Ala Ile Leu  
 835 840 845  
 att gac ctt ggg cga acc ttt cct aca cac cca tac ttc tct gcc cag 2770  
 Ile Asp Leu Gly Arg Thr Phe Pro Thr His Pro Tyr Phe Ser Ala Gln  
 850 855 860 865  
 ctt gga gca gga cag cta tcg ctt tac aac att ttg aag gcc tac tca 2818  
 Leu Gly Ala Gly Gln Leu Ser Leu Tyr Asn Ile Leu Lys Ala Tyr Ser  
 870 875 880  
 ctt cta gac cag gaa gtg gga tat tgc caa ggt ctc agc ttt gta gca 2866  
 Leu Leu Asp Gln Glu Val Gly Tyr Cys Gln Gly Leu Ser Phe Val Ala  
 885 890 895  
 ggc att ttg ctt ctt cat atg agt gag gaa gag gcg ttt aaa atg ctc 2914  
 Gly Ile Leu Leu Leu His Met Ser Glu Glu Glu Ala Phe Lys Met Leu  
 900 905 910  
 aag ttt ctg atg ttt gac atg ggg ctg cgg aaa cag tat cgg cca gac 2962  
 Lys Phe Leu Met Phe Asp Met Gly Leu Arg Lys Gln Tyr Arg Pro Asp  
 915 920 925  
 atg att att tta cag atc cag atg tac cag ctc tcg agg ttg ctt cat 3010  
 Met Ile Ile Leu Gln Ile Gln Met Tyr Gln Leu Ser Arg Leu Leu His  
 930 935 940 945  
 gat tac cac aga gac ctc tac aat cac ctg gag gag cac gag atc ggc 3058  
 Asp Tyr His Arg Asp Leu Tyr Asn His Leu Glu Glu His Glu Ile Gly  
 950 955 960  
 ccc agc ctc tac gct gcc ccc tgg ttc ctc acc atg ttt gcc tca cag 3106  
 Pro Ser Leu Tyr Ala Ala Pro Trp Phe Leu Thr Met Phe Ala Ser Gln  
 965 970 975  
 ttc ccg ctg gga ttc gta gcc aga gtc ttt gat atg att ttt ctt cag 3154  
 Phe Pro Leu Gly Phe Val Ala Arg Val Phe Asp Met Ile Phe Leu Gln  
 980 985 990  
 gga aca gag gtc ata ttt aaa gtg gct tta agt ctg ttg gga agc cat 3202

```
<210> 5
<211> 1168
<212> PRT
```

&lt;213&gt; Homo sapiens

&lt;400&gt; 5

```

Met Glu Pro Ile Thr Phe Thr Ala Arg Lys His Leu Leu Pro Asn Glu
1           5           10           15
Val Ser Val Asp Phe Gly Leu Gln Leu Val Gly Ser Leu Pro Val His
           20           25           30
Ser Leu Thr Thr Met Pro Met Leu Pro Trp Val Val Ala Glu Val Arg
           35           40           45
Arg Leu Ser Arg Gln Ser Thr Arg Lys Glu Pro Val Thr Lys Gln Val
           50           55           60
Arg Leu Cys Val Ser Pro Ser Gly Leu Arg Cys Glu Pro Glu Pro Gly
65           70           75           80
Arg Ser Gln Gln Trp Asp Pro Leu Ile Tyr Ser Ser Ile Phe Glu Cys
           85           90           95
Lys Pro Gln Arg Val His Lys Leu Ile His Asn Ser His Asp Pro Ser
           100          105          110
Tyr Phe Ala Cys Leu Ile Lys Glu Asp Ala Val His Arg Gln Ser Ile
           115          120          125
Cys Tyr Val Phe Lys Ala Asp Asp Gln Thr Lys Val Pro Glu Ile Ile
           130          135          140
Ser Ser Ile Arg Gln Ala Gly Lys Ile Ala Arg Gln Glu Glu Leu His
145          150          155          160
Cys Pro Ser Glu Phe Asp Asp Thr Phe Ser Lys Lys Phe Glu Val Leu
           165          170          175
Phe Cys Gly Arg Val Thr Val Ala His Lys Lys Ala Pro Pro Ala Leu
           180          185          190
Ile Asp Glu Cys Ile Glu Lys Phe Asn His Val Ser Gly Ser Arg Gly
           195          200          205
Ser Glu Ser Pro Arg Pro Asn Pro Pro His Ala Ala Pro Thr Gly Ser
           210          215          220
Gln Glu Pro Val Arg Arg Pro Met Arg Lys Ser Phe Ser Gln Pro Gly
225          230          235          240
Leu Arg Ser Leu Ala Phe Arg Lys Glu Leu Gln Asp Gly Gly Leu Arg
           245          250          255
Ser Ser Gly Phe Phe Ser Ser Phe Glu Glu Ser Asp Ile Glu Asn His
           260          265          270
Leu Ile Ser Gly His Asn Ile Val Gln Pro Thr Asp Ile Glu Glu Asn
           275          280          285
Arg Thr Met Leu Phe Thr Ile Gly Gln Ser Glu Val Tyr Leu Ile Ser
           290          295          300
Pro Asp Thr Lys Lys Ile Ala Leu Glu Lys Asn Phe Lys Glu Ile Ser

```

T05230"TE29260



645	650	655
Arg Arg His Ser Trp Arg Gln Gln Ile Phe Leu Arg Val Ala Thr Pro		
660	665	670
Gln Lys Ala Cys Asp Ser Ser Ser Arg Tyr Glu Asp Tyr Ser Glu Leu		
675	680	685
Gly Glu Leu Pro Pro Arg Ser Pro Leu Glu Pro Val Cys Glu Asp Gly		
690	695	700
Pro Phe Gly Pro Pro Pro Glu Glu Lys Lys Arg Thr Ser Arg Glu Leu		
705	710	715
Arg Glu Leu Trp Gln Lys Ala Ile Leu Gln Gln Ile Leu Leu Leu Arg		
725	730	735
Met Glu Lys Glu Asn Gln Lys Leu Gln Ala Ser Glu Asn Asp Leu Leu		
740	745	750
Asn Lys Arg Leu Lys Leu Asp Tyr Glu Glu Ile Thr Pro Cys Leu Lys		
755	760	765
Glu Val Thr Thr Val Trp Glu Lys Met Leu Ser Thr Pro Gly Arg Ser		
770	775	780
Lys Ile Lys Phe Asp Met Glu Lys Met His Ser Ala Val Gly Gln Gly		
785	790	795
Val Pro Arg His His Arg Gly Glu Ile Trp Lys Phe Leu Ala Glu Gln		
805	810	815
Phe His Leu Lys His Gln Phe Pro Ser Lys Gln Gln Pro Lys Asp Val		
820	825	830
Pro Tyr Lys Glu Leu Leu Lys Gln Leu Thr Ser Gln Gln His Ala Ile		
835	840	845
Leu Ile Asp Leu Gly Arg Thr Phe Pro Thr His Pro Tyr Phe Ser Ala		
850	855	860
Gln Leu Gly Ala Gly Gln Leu Ser Leu Tyr Asn Ile Leu Lys Ala Tyr		
865	870	875
Ser Leu Leu Asp Gln Glu Val Gly Tyr Cys Gln Gly Leu Ser Phe Val		
885	890	895
Ala Gly Ile Leu Leu Leu His Met Ser Glu Glu Glu Ala Phe Lys Met		
900	905	910
Leu Lys Phe Leu Met Phe Asp Met Gly Leu Arg Lys Gln Tyr Arg Pro		
915	920	925
Asp Met Ile Ile Leu Gln Ile Gln Met Tyr Gln Leu Ser Arg Leu Leu		
930	935	940
His Asp Tyr His Arg Asp Leu Tyr Asn His Leu Glu Glu His Glu Ile		
945	950	955
Gly Pro Ser Leu Tyr Ala Ala Pro Trp Phe Leu Thr Met Phe Ala Ser		
965	970	975
Gln Phe Pro Leu Gly Phe Val Ala Arg Val Phe Asp Met Ile Phe Leu		

105290-11229260

980 985 990  
 Gln Gly Thr Glu Val Ile Phe Lys Val Ala Leu Ser Leu Leu Gly Ser  
 995 1000 1005  
 His Lys Pro Leu Ile Leu Gln His Glu Asn Leu Glu Thr Ile Val Asp  
 1010 1015 1020  
 Phe Ile Lys Ser Thr Leu Pro Asn Leu Gly Leu Val Gln Met Glu Lys  
 1025 1030 1035 1040  
 Thr Ile Asn Gln Val Phe Glu Met Asp Ile Ala Lys Gln Leu Gln Ala  
 1045 1050 1055  
 Tyr Glu Val Glu Tyr His Val Leu Gln Glu Glu Leu Ile Asp Ser Ser  
 1060 1065 1070  
 Pro Leu Ser Asp Asn Gln Arg Met Asp Lys Leu Glu Lys Thr Asn Ser  
 1075 1080 1085  
 Ser Leu Arg Lys Gln Asn Leu Asp Leu Leu Glu Gln Leu Gln Val Ala  
 1090 1095 1100  
 Asn Gly Arg Ile Gln Ser Leu Glu Ala Thr Ile Glu Lys Leu Leu Ser  
 1105 1110 1115 1120  
 Ser Glu Ser Lys Leu Lys Gln Ala Met Leu Thr Leu Glu Leu Glu Arg  
 1125 1130 1135  
 Ser Ala Leu Leu Gln Thr Val Glu Glu Leu Arg Arg Arg Ser Ala Glu  
 1140 1145 1150  
 Pro Ser Asp Arg Glu Pro Glu Cys Thr Gln Pro Glu Pro Thr Gly Asp  
 1155 1160 1165

&lt;210&gt; 6

&lt;211&gt; 18

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;221&gt; misc\_binding

&lt;222&gt; 1..18

&lt;223&gt; sequencing oligonucleotide PrimerPU

&lt;400&gt; 6

tgtaaaacga cggccagt

18

&lt;210&gt; 7

&lt;211&gt; 18

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

T05390" T0229260

